

CORRIGENDA

Association between variants at the GABA_Aβ2, GABA_Aα6 and GABA_Aγ2 gene cluster and alcohol dependence in a Scottish population

E-W Loh, I Smith, R Murray, M McLaughlin, S McNulty and D Ball. *Mol Psychiatry* 1999; **4**: 539–544

The sentence on page 540 of this article ‘A silent exonic gCt → gTt substitution (Ala → Ala) at nucleotide 1412 was found by SSCP analysis’ is incorrect. The

correct sentence should read ‘A silent exonic GCT → GCC substitution (Ala → Ala) at nucleotide 1412 was found by SSCP analysis’.

Family-based linkage disequilibrium mapping using SNP marker haplotypes: application to a potential locus for schizophrenia at chromosome 22q11

T Li, D Ball, J Zhao, RM Murray, X Liu, PC Sham and DA Collier. *Mol Psychiatry* 2000; **5**: 77–84

The sequence for the oligonucleotide primers used for SSCP analysis of the COMT promoter region, and genotyping of the *Hind*III polymorphism within, were given

incorrectly. The correct primer sequences are: forward 5'-TAG TAA CAG ACT GGG CAC GAA and reverse 5'-GTT CAA AGG GCA TTT ATC ATG.