# npg

# **ORIGINAL ARTICLE**

# Effect of PGX, a novel functional fibre supplement, on subjective ratings of appetite in overweight and obese women consuming a 3-day structured, low-calorie diet

V Kacinik<sup>1</sup>, M Lyon<sup>1,2</sup>, M Purnama<sup>1</sup>, RA Reimer<sup>3</sup>, R Gahler<sup>4</sup>, TJ Green<sup>2</sup> and S Wood<sup>2,5</sup>

Introduction: Dietary factors that help control perceived hunger might improve adherence to calorie-reduced diets.

**Objectives:** The objective of the study was to investigate the effect of supplementing a three-day, low-calorie diet with PolyGlycopleX (PGX), a highly viscous fibre, on subjective ratings of appetite compared with a placebo.

**Methods:** In a double-blind crossover design with a 3-week washout, 45 women (aged  $38\pm9$  years, body mass index  $29.9\pm2.8$  kg m<sup>-2</sup>) were randomised to consume a 1000-kcal per day diet for 3 days, supplemented with 5 g of PGX or placebo at each of breakfast, lunch and dinner. Subjective appetite was assessed using 100 mm visual analogue scales that were completed daily before, between and after consumption of meals.

**Results:** Thirty-five women completed the study. Consumption of PGX compared with placebo led to significantly lower mean area under the curve for hunger on day 3 (440.4 versus 375.4; P = 0.048), prospective consumption on day 3 (471.0 versus 401.8; P = 0.017) and the overall 3-day average (468.6 versus 420.2; P = 0.026). More specifically, on day 3 PGX significantly reduced total appetite, hunger, desire to eat and prospective consumption for 2.5 and 4.5 h after lunch and before dinner times, with hunger also being reduced 2.5 h after dinner (P < 0.05).

**Conclusion:** The results show that adding 5 g of PGX to meals during consumption of a low-calorie diet reduces subjective ratings of prospective consumption and increases the feelings of satiety, especially during afternoon and evening. This highly viscous polysaccharide may be a useful adjunct to weight-loss interventions involving significant caloric reductions. *Nutrition and Diabetes* (2011) 1, e22; doi:10.1038/nutd.2011.18; published online 12 December 2011

Keywords: viscous; soluble; fibre; appetite; PGX; PolyGlycopleX

# Introduction

Obesity increases the risk of type 2 diabetes, cardiovascular disease and other chronic diseases. 1,2 Weight reduction is associated with significant improvements in blood pressure, serum cholesterol levels and glycaemic control. Achieving negative energy balance is an important factor in determining the magnitude and rate of weight loss. Calorie-restricted diets, ideally combined with exercise and behaviour modification, are considered to be the initial treatment strategy for overweight and obese individuals. Low-calorie diets at a level of 1000–1200 kcal per day for most overweight men are

often recommended for weight loss.<sup>5</sup> With low-calorie diets, maintaining satiety is critical, as perceived hunger has been shown to be a significant predictor of failure to lose weight.<sup>6</sup> After weight loss is achieved, appetite often increases leading to refractory weight gain.<sup>7</sup> Identifying factors that suppress appetite is important for successful weight loss and maintenance.

Increased fibre intake has been associated with reduced energy intake and increased satiety, and may be an important factor in obesity management.<sup>8</sup> A meta-analysis of 22 studies concluded that consumption of 14 g of fibre per day resulted in a 10% decrease in energy intake and a 1.9-kg weight loss over 3.8 months.<sup>9</sup> Viscous soluble fibres, in particular, appear to increase satiety<sup>10–14</sup> and promote adherence to calorie-reduced diets.<sup>15,16</sup> Accordingly, consuming substantial quantities of viscous dietary fibre may provide significant therapeutic benefits under conditions of reduced energy intake. However, as the benefit of these fibres

<sup>&</sup>lt;sup>1</sup>Canadian Centre for Functional Medicine, Coquitlam, British Columbia, Canada; <sup>2</sup>Faculty of Land and Food Systems and Food, Nutrition and Health Program, University of British Columbia, Vancouver, British Columbia, Canada; <sup>3</sup>Faculty of Kinesiology and Department of Biochemistry and Molecular Biology, University of Calgary, Calgary, Alberta, Canada; <sup>4</sup>Factors Group Research & Development, Burnaby, British Columbia, Canada and <sup>5</sup>InovoBiologic, Calgary, Alberta, Canada



is related to viscosity,<sup>17</sup> their acceptance is often hampered by issues of palatability.<sup>18</sup>

Recently, PolyGlycopleX (PGX), a novel, highly viscous non-starch polysaccharide complex has been developed for human consumption. This soluble dietary fibre displays delayed viscosity for 15–30 min after ingestion, thus, allowing for a more palatable and easy-to-use functional fibre. In a recent randomised, placebo-controlled trial, PGX was shown to diminish hunger and reduce *ad libitum* food consumption compared with less viscous fibres (cellulose and glucomannan) given in meal-replacement drinks. <sup>19</sup> The objective of this study was to investigate the effect of supplementing a three-day low-calorie diet with PGX on subjective ratings of appetite compared with a placebo in overweight and obese women.

# Materials and methods

#### Subjects

Subjects were recruited through local newspaper advertisements. Women with a body mass index between 25 and 34.9 kg m<sup>-2</sup> and between 20 and 50 years of age were eligible to participate. Exclusion criteria included: pregnancy or lactation, smoking, consumption of more than two alcoholic drinks daily or nine alcoholic drinks weekly, recent participation in a calorie-reduced diet, the presence of chronic diseases, past history of gastrointestinal surgery, a change in body weight of more than 3 kg during the last 3 months, hypersensitivity to the study foods, taking medication or natural health products, which may affect gastrointestinal motility hunger or appetite. Furthermore excluded were those who scored 9 or more on the restraint scale; 27 or above for uncontrolled eating, or 18 or greater for emotional eating on the Three-Factor Eating Questionnaire-R18V2, validated for Canadian and US obese and non-obese individuals.<sup>20</sup> As lower palatability of foods may diminish the sensation of hunger and desire to eat, 21 a Food Preference Questionnaire was used to ensure that all the participants expressed acceptance of the foods included in the study. Those scoring less than 3 for more than 50% of the food items were excluded from the study. The University of British Columbia Research Ethics Committee approved the study and all subjects gave informed consent to participate.

# Study supplements

PGX ( $\alpha$ -D-glucurono- $\alpha$ -D-manno- $\beta$ -D-manno- $\beta$ -D-gluco), ( $\alpha$ -L-gulurono- $\beta$ -D mannurono),  $\beta$ -D-gluco- $\beta$ -D-mannan (Inovobiologic, Calgary, Alberta, Canada) is a novel functional fibre complex 22,23 manufactured by a proprietary process from three fibres (konjac (glucomannan), sodium alginate and xanthan gum) to form a highly viscous polysaccharide (higher viscosity than any currently known individual polysaccharide or fibre blend) with high water-holding and gel-forming properties. PGX has been shown to be safe and

well tolerated in rodents<sup>24</sup> and in humans.<sup>25</sup> Genotoxicity studies of PGX have shown no mutagenic effects using bacterial reverse mutation and mouse micronucleus assays.<sup>26</sup> PGX is 87.4% dietary fibre, of which 81.8% is soluble. It can be sprinkled on food before consumption or taken with soup or beverage before meals. Rice flour was used as a placebo and has been used in other clinical trials for its white colour, neutral taste and hypoallergenicity.<sup>27,28</sup> PGX and rice flour both consist of approximately similar carbohydrate content with carbohydrate in PGX being dietary fibre and in rice flour being starch. Boiling or other forms of moist heat processing is required for rice flour to be effectively digested. Without cooking, only about 1% of the starch in raw rice flour is digested.<sup>29</sup> Therefore, the carbohydrate in uncooked rice flour is essentially dietary fibre resulting in rice flour's caloric value that closely approximates PGX. Both study supplements were packaged with clear plastic scoops in white plastic containers containing 100 g and were indistinguishable from each other.

# Study design/protocol

The study was a prospective, randomised, double-blind, placebo-controlled crossover trial. The study had two experimental phases with a 3-week washout. Test periods were on the same days of the week (Tuesday-Thursday) and were conducted during the follicular phase (days 4–10) of the women's menstrual cycle to minimise hormonal influences on eating/feeding behaviour and appetite.<sup>30</sup> On Monday of the test week, subjects attended the clinic where their bodyweight was measured, study food was provided and they were randomised to either the test or placebo supplement. During the test period, subjects consumed the breakfasts, lunches and dinners provided immediately after the food was sprinkled with 5 g of supplement or placebo. Participants completed visual analogue scales (VAS) questionnaires at specified times (see below). To avoid a second meal effect and to minimise differences in glycogen stores, the night before consumption of the 3-day low-calorie diet, all subjects consumed the same evening meal. In addition, participants were instructed not to consume alcohol or engage in vigorous physical activity 48 h before consuming the low-calorie diet. On Friday, subjects returned to the clinic to have their compliance assessed, documents reviewed and to have any adverse effects recorded. To assess compliance, subjects kept a log of daily food and study product consumed and returned the product containers for weighing.

#### Low-calorie diet

A three-day structured low-calorie diet of 1000 kcal per day was provided with the intention that caloric restriction would accentuate feelings of hunger. The low-calorie diet was designed to provide participants with 65% of energy as carbohydrate, 10% protein, and 15% fat (Appendix 1). Commercially available frozen dinner entrees (President's



Choice Blue Menu, Loblaw, Ontario, Canada) were provided to participants to standardise caloric intake and macronutrient composition.

# Subjective appetite measures

Rating subjective appetite using VAS is a valid and reliable measure of appetite under both experimental and free-living conditions, especially when using a within-subject design, as used in the current study. 31,32 VAS ratings were made on a 100-mm scale with words anchored at either end in the following order: first, 'How strong is your desire to eat?' (Very strong—Very weak); second, 'How hungry do you feel?' (Not hungry at all—Very hungry); third, 'How full do you feel?' (Very full—Not at all full); and fourth, 'How much food do you think you could eat?' (A large amount-Nothing at all), which assesses prospective consumption. The palatability of the meals was also evaluated after meals by a fifth question asking 'How would you rate the overall palatability of the food? (Unpalatable—Very palatable). The times of rating were the same every day for both phases; however, to work with participants at various schedules, different start times in the morning were available. VAS were completed before (e.g., 0755 h) and after breakfast (e.g., 0830 h), in between the morning meals (e.g., 1030 h), before (e.g., 1225 h) and after (e.g., 1300 h) lunch, in between meals in the afternoon (e.g., 1500 h, 1700 h), before (e.g., 1725 h) and after (e.g., 1800 h) dinner and in the evening (e.g., 2000 h). Participants were provided with programmable alert wrist watches (CADEX Medication reminder watch, e-pill, LLC., Wellesley, MA, USA) to remind them when to eat and when to complete the questionnaires. Participants were allowed 20 min to consume the food, supplement and 500 ml of water, and were allowed to consume an additional 500 ml of water between meals. Participants were instructed not to consume water 30 min before completing premeal VAS. All food and water consumed during Phase 1 was recorded and the same amount was provided for Phase 2.

#### Statistical analyses

Changes in weight over each test period were tested for differences between treatment groups using the Wilcoxon rank sum test. For compliance, the *P*-values of differences between the two treatment groups within a single period were calculated from two-sample t-tests; the statistical test for the summary across both periods of data used a linear mixed model approach, which took into consideration that the same participants appeared once on each of the two treatments (due to the crossover design). For VAS score analyses, the area under the curve was analysed on the log scale to meet the distributional assumptions required for the statistical methods. The treatment effect was estimated from a linear mixed-effects model to take into consideration the crossover design of the study; the model was adjusted for phase, treatment, randomised sequence and a within-patient comparison was made to assess the treatment effect. With

the different study days expected to give differences in results, the analysis was carried out by days 1, 2, 3 and on the mean of days 1-3. To analyse VAS scores by time point, a similar linear mixed model was used incorporating terms in the model for treatment, phase, randomised group and time point, and then contrasts used to compare the treatment groups at the different time points. Statistical testing of the adverse event rates by treatment (either overall rates, treatment-related rates or rates for specific types of adverse events) was performed using generalised estimating equations logistic regression models. To evaluate whether there was a carryover effect in the second period by the randomised sequences, a standard statistical approach by Jones et al.33 was used to assess. For all statistical tests, the significance level was set at 0.05. All analyses were performed using SAS (Version 9.1.3, SAS Institute, Cary, NC, USA).

#### Results

In total, 51 individuals were recruited for the study of which 45 were randomised (Table 1). After Phase 1, five discontinued from the study. The reasons for withdrawal were adverse events (n=2), dislike of the frozen meals offered (n=1) and personal reasons unrelated to the study (n=2). Overall, 40 participants completed the entire study; however, only 35 participants were included in the per-protocol analyses as 2 participants deviated from the protocol and 3 reported feeling sick, which might affect subjective appetite ratings. Compliance was high for both groups, with 107% and 97% for the rice flour and PGX groups, respectively. There was no carryover effect in the second period by the randomised sequences.

Weight loss was similar between groups over the two phases (PGX 1.3 kg versus placebo 1.4 kg; P = 0.376). From the intent-to-treat analysis (n = 40), the PGX treatment resulted in a significantly lower mean area under the curve

Table 1 Subject characteristics

	Value (range)		
Age (years)	38.6 ± 9.1 (20–50)		
Weight (kg)	79.8 ± 9.3 (59.4–107.8)		
Body mass index (kg m <sup>-2</sup> )	29.9 ± 2.8 (25.0–34.8)		
Waist circumference (cm)	87.0 ± 7.4 (73.3–106.0)		
Three-factor eating questionnaire <sup>a</sup>			
Cognitive	$7.3 \pm 3.1 \ (4.0-25.0)$		
Uncontrolled eating	$21.2 \pm 4.6 (5.0-27.0)$		
Emotional eating	$14.3 \pm 3.8 \ (7.0-24.0)$		

Values are mean  $\pm$  s.d.; n = 45. The Three-Factor Eating Questionnaire evaluates some aspects of eating behaviours such as cognitive restraint (i.e., the voluntary control of eating), the tendency to lose control of food intake when faced with external cues, mood changes or disruptive events (i.e., disinhibition) and susceptibility to feelings of hunger. In this study, the Three-Factor Eating-R18V2 questionnaire was used as a means to exclude those who have potentially disordered eating motivation and to assess motivation of the participants eligible for the study.<sup>20</sup>




on day 3 for prospective consumption (P=0.018) and a trend towards significance for hunger (P = 0.059). After elimination of the protocol deviators (n=35), a similar pattern was seen with the differences between groups stronger with a mean percent change of prospective consumption at -14.7 (P = 0.017) and hunger at -14.8 on day 3 (P = 0.048, Table 2). There were no statistical differences in mean area under the curve for any of the scores for day 1 or 2, or the mean days 1-3, except for prospective consumption (P = 0.026). The majority of differences in subjective ratings between the placebo and PGX groups at various time points were detected on day 3. PGX supplementation significantly reduced total appetite, desire to eat, hunger, prospective consumption, and it was associated with increased fullness at 2.5 h after lunch (Figure 1); reduced total appetite, desire to eat, hunger and prospective consumption at 4.5 h after lunch (Figure 2) and reduced total appetite, hunger and prospective consumption before dinner (Figure 3). Hunger was also reduced at 2.5 h after dinner (P<0.05). For palatability, there was a statistically significant difference in mean palatability VAS scores by treatment for each day (P < 0.05), with PGX-supplemented meals scoring above 50 mm, but 5-10 mm below mean rice flour scores (data not shown).

Adverse events reported by both treatment groups are summarised in Table 3. There was no statistical difference between the two treatment groups in terms of the adverse events experience of the participants. Furthermore, the overall or treatment-related adverse events were similar for the two treatments.

# Discussion

This study demonstrates that adding 5 g of PGX to meals at the start of a low-calorie diet helps manage appetite by increasing satiety and decreasing prospective consumption (the amount one thinks one could eat). This effect was not statistically significant until day 3; however it resulted in reductions above 10% compared with the control, a




**Figure 1** Comparison of 2.5 h post breakfast, lunch and dinner mean hunger and prospective consumption scores of day 3 of the 1000-kcal calorie diet supplement with either 5 g of PGX or placebo at each meal. Values shown are mean  $\pm$  s.e. (n= 35). \*Significantly lower scores with PGX than with the placebo supplement (P<0.05).


Table 2 Daily and 3-day average comparison of subjective appetite measurements calculated as mean area under the curve in response to placebo or PGX for the per-protocol subject group

Score	Day	Placebo	PGX	% Change	P-value
Prospective consumption	1	457.2 ± 29.4	432.5 ± 27.8	-5.4	0.358
	2	477.9 ± 33.1	$427.1 \pm 29.6$	-10.6	0.079
	3	471 ± 32.5	$401.8 \pm 27.7$	-14.7	0.017
	Average days 1–3	$468.6 \pm 28.9$	$420.2 \pm 25.9$	-10.3	0.026
Desire	1	432.5 ± 29.4	$428.8 \pm 29.1$	-0.9	0.894
	2	$456.3 \pm 34.2$	$416.9 \pm 31.3$	-8.6	0.273
	3	$427.7 \pm 35$	$387.9 \pm 31.7$	-9.3	0.280
	Average days 1–3	$438.6 \pm 29.7$	$410.8 \pm 27.8$	-6.3	0.303
Fullness	1	598.3 ± 40.3	591.6 ± 39.9	-1.1	0.832
	2	584.5 ± 50.7	574.5 ± 49.8	-1.7	0.785
	3	614.1 ± 47.2	$632.3 \pm 48.6$	3	0.592
	Average days 1–3	$598.8 \pm 43.8$	$599 \pm 43.9$	0	0.996
Hunger	1	431.5 ± 28.6	418.3 ± 27.7	-3.1	0.651
	2	$434.9 \pm 33.6$	$407.6 \pm 31.5$	-6.3	0.429
	3	440.4 ± 32.9	$375.4 \pm 28$	-14.8	0.048
	Average days 1–3	$435.6 \pm 28.4$	$400 \pm 26.1$	-8.2	0.162
Total appetite	1	469.8 ± 27.3	458.7 ± 26.7	-2.4	0.675
	2	475.9 ± 31.8	$449.4 \pm 30.0$	-5.6	0.390
	3	464.5 ± 31.9	$414.3 \pm 28.4$	-10.8	0.106
	Average days 1-3	470 ± 27.9	$440.4 \pm 26.2$	-6.3	0.215

Abbreviation: PGX, PolyGlycopleX. Note: P-value is from a linear mixed-model fit on log scale of area under curve and adjusting for potential sequence and period effects in addition to treatment (significance level P<0.05). Values shown are mean  $\pm$  s.e.; n = 35.



**Figure 2** Comparison of the 4.5 h post lunch mean hunger and prospective consumption scores of day 3 of the 1000-kcal calorie diet supplement with either 5 g of PGX or placebo at each meal. Values shown are mean  $\pm$  s.e. (n= 35). \*Significantly lower scores with PGX than with the placebo supplement (P<0.05).



**Figure 3** Comparison of predinner mean hunger and prospective consumption scores of day 3 of the 1000-kcal calorie diet supplement with either 5 g of PGX or placebo at each meal. Values shown are mean  $\pm$  s.e. (n= 35). \*Significantly lower scores with PGX than with the placebo supplement (P<0.05).

magnitude which is considered to be of practical relevance.<sup>32</sup> With respect to time of day, PGX was found to exert its strongest effects in the afternoon and in the evening, reducing total appetite, hunger, desire to eat and prospective consumption. These results might be particularly helpful in managing obesity, given reports that food intake tends to be less in the morning and greater in the afternoon and highly evening in obese versus normal weight individuals.<sup>34</sup> One explanation for the effect seen with PGX on day 3 could be the increased fibre load. This particular test day had 7 g more dietary fibre and 9–10 g less protein than on day 1 or 2, resulting in a possible

synergistic effect and interaction between the PGX and the dietary fibre content of the meals. Another possibility may be the cumulative effects of PGX over time with the potential for upregulation of gut-satiety hormone gene expression (such as proglucagon) via fermentation and production of short-chain fatty acids in the large bowel.<sup>35</sup>

Soluble fibres that are fermentable by gut microbiota and produce significant quantities of short-chain fatty acids may contribute to appetite regulation by stimulating the release of gut-satiety hormones such as peptide YY (PYY) and glucagon like peptide-1.<sup>36</sup> In a study using Zucker diabetic rats, PGX consumption was associated with a significant increase in glucagon like peptide-1 levels.<sup>37</sup> In a 3-week human tolerance trial, supplementation of regular diets with 10 g of PGX per day was associated with higher fasting PYY levels than placebo control.<sup>38</sup> It has been reported that obese individuals have lower fasting and postprandial circulating PYY levels than lean individuals, <sup>39,40</sup> which may result in impaired satiety and may contribute to the development of obesity and may hinder weight loss efforts. To produce an equivalent stimulation of PYY sufficient to promote satiety, obese individuals need to consume a much greater caloric load than their lean counterparts.<sup>39</sup> An alternative to a greater caloric load could be a volumetric, lower energy dense load from supplementing a meal with PGX to augment PYY levels and stimulate satiety. Further investigation of a low-calorie diet supplemented with PGX on gut-satiety hormones is thus warranted.

Under conditions of fixed energy intake, such as a lowcalorie diet, viscous soluble fibres may also alter the viscosity of gastrointestinal content resulting in multiple effects on satiety. A magnetic resonance imaging study by Marciani et al.41 showed that a high-viscosity nutrient meal had the strongest additive effect in gastric distension and delayed gastric emptying compared with low-viscosity non-nutrient, low-viscosity nutrient and high-viscosity non-nutrient meals. Viscous soluble fibres also delay the absorption of nutrients by thickening the unstirred layer at the gut mucosal surface. This results in the transport of larger amounts of nutrients to the ileum and/or distributes nutrient absorption over a larger intestinal luminal surface area, leading to a more pronounced inhibitory feedback from the distal gut and activation of the ileal brake.<sup>42</sup> Other mediators of the ileal brake include the gut-satiety hormones PYY and glucagon like peptide-1 and vagal nerve stimulation.<sup>43</sup> The ileal brake is a distal to proximal negative feedback mechanism that inhibits gastric emptying, proximal gastrointestinal motility and secretion to control the transit of a meal through the gastrointestinal tract, which consequently optimises nutrient digestion and absorption.<sup>44</sup>

As a highly viscous soluble fibre, PGX may cause a delay in gastric emptying and/or prolong the transport of nutrients down the gastrointestinal tract as demonstrated by previous research on its effects on the glycaemic index (GI). The GI is a property of carbohydrate-containing food that describes the rise of blood glucose occurring after a meal. High GI



Table 3 Summary of the adverse events (AEs) by treatment allocation

Variable		Placebo	PGX	P-value
Number of participants		38	42	
Participants reporting adverse events		13 (34.2%)	12 (28.6%)	0.7
Total adverse events		25	28	
Participants reporting treatment-related events		8 (21.1%)	11 (26.2%)	0.34
Total treatment-related Events		19	24	
Total adverse events (by severity)	Unknown severity	0	0	
	Mild	5	2	
	Moderate	13	21	
	Severe	7	5	
Total treatment-related events (by severity)	Unknown Severity	0	0	
	Mild	4	2	
	Moderate	9	18	
	Severe	6	4	
Participants reporting AEs by type of AE	Constipation	3 (7.9%)	1 (2.4%)	0.15
	Diarrhea .	0 (0.0%)	3 (7.1%)	NA
	Bloating	4 (10.5%)	8 (19%)	0.07
	Flatulence	0 (0.0%)	2 (4.8%)	NA
	Headache	4 (10.5%)	1 (2.4%)	0.17
	Cramping	3 (7.9%)	3 (7.1%)	0.59
	Sick/unwell	4 (10.5%)	1 (2.4%)	0.08
	Other	3 (7.9%)	7 (16.7%)	0.06

Abbreviation: PGX, PolyGlycopleX. Parentheses denote the percentage from the total number of participants. Some *P*-values were not possible to compute due to no events being reported under one of the treatment groups. These are noted as 'NA' in the table.

foods are classified as being rapidly digested and absorbed into the bloodstream, resulting in a sharp rise and in a steep decline of blood glucose after consumption, whereas low GI foods are broken down more slowly by a slower rate of digestion and absorption and releasing glucose more gradually into the bloodstream. 45 When 2.5-5 g of PGX was sprinkled on high and moderate GI foods, it was consistently shown to be highly effective in significantly reducing their glycaemic index. 45,46 Influencing the digestive rate of foods and their blood glucose response may have an important effect on the perception of satiety. Arumugam et al.<sup>47</sup> examined the effects of high-GI versus low-GI meals on subjective satiety in overweight and obese women. They found that the appetite score was 35% higher at 4h after the high-glycaemic versus low glycaemic breakfast, and that at 3, 4 and 5 h after lunch, the appetite score was 30-44% higher in the high glycaemic condition.<sup>47</sup> Many other short-term studies (<1 day) have also reported a significant relationship between the glycaemic response and satiety using foods with varving glycaemic indices. 48-51 In the present study, the frozen meals were chosen to represent the average Western diet with convenient, processed higher GI foods and as such the effect of PGX supplementation on the study foods' glycaemic indices might be related to the 28%, 14%, 12% reduction of total appetite score at 2.5, 4.5, 5 h, respectively, after consuming the lunch on day 3. If however the study foods had a low GI, PGX may not have exerted as significant an effect on satiety.

This study was designed to alleviate some potentially confounding variables in appetite research. The timing of the test phases was scheduled according to participants' menstrual cycle, given that previous research has linked changes in gastrointestinal function, appetite and energy intake to the follicular and luteal phases of menstruation.<sup>30</sup> The energy intake, macronutrient composition, water consumption and eating schedule for all the meals over the three-diet days were controlled, demonstrating the effect of PGX on homeostatic hunger (physiological). Although the effect of PGX on ad libitum food intake cannot be concluded, prior research has shown that subjective ratings of appetite appear to be good indicators of motivation to eat and are predictors of actual food intake.<sup>31,52,53</sup> However, it should also be mentioned that appetite ratings have not always been found to correlate with ad libitum food intake,<sup>10</sup> as many factors influence food intake, including sensory hedonics, external factors, cognitive issues, body weight, genes and so on.<sup>54–56</sup>

To minimise the effect of palatability on appetite ratings, women who rated the palatability of the meals low were excluded. However, a significant difference in palatability between meals supplemented with PGX and rice flour was observed on all of the three-diet days. Previous research with PGX has not found any significant differences in palatability scores in comparisons with controls. 19,45 Study participants were advised to sprinkle PGX granules as they ate; however, the hot microwaved meals for lunch and dinner more than likely accelerated viscosity development, thereby affecting texture and palatability. This difference may be related more to texture than to any gastrointestinal discomforts experienced with PGX, as there was no statistical difference in adverse events reported between PGX and rice flour. Although palatability affects appetite, 57,58 this did not translate into significant differences in subjective appetite

scores for days 1 and 2, suggesting that the difference observed on day 3 may have not been due to the palatability of the PGX supplemented meal. When designing placebocontrolled trials with PGX, it is difficult to find a comparable inert substance and completely avoid any potentially confounding effect. Uncooked rice flour was found to closely approximate PGX in terms of flavour and texture, and it was thus judged to be a suitable placebo. However, the presence of resistant starches may have produced an increase in fermentation in the control group and dampened the difference in results between the two interventions.

As an initial investigation, this study was performed over 3 days because of the number of VAS questionnaires dispensed and strict adherence required for the food and meal schedule. It was conducted under free-living conditions so as to mimic people following a structured low-calorie diet in their regular lives. However, this inherently introduced some variability in the participants' environment, occupational and non-occupational activities, which could have resulted in diurnal variations in appetite. <sup>59</sup> Furthermore, the diet during the two test phases was self-administered and even though compliance of the study supplements was high, the possibility of noncompliance with the prescribed diets could not be completely avoided.

In summary, this study indicates that the supplementation of highly viscous PGX to meals could be a useful weight management aid during a low-calorie diet to help lessen feelings of hunger and to moderate food portions. Weight relapse after consuming a low-calorie diet is a common outcome of obesity interventions. Therefore, long-term studies investigating the effects of PGX on subjective appetite and the maintenance of reduced body weight would be warranted. In addition, it would be useful to examine the effects of PGX on biomarkers such as glucagon like peptide-1, PYY and ghrelin to better understand some of the physiological influences of PGX on satiety. By virtue of its multiple physiological effects on the mechanisms of satiety, PGX may be an effective weight-management aid for low-calorie diets.

# **Conflict of interest**

VK and MP are employees of the Canadian Centre for Functional Medicine where the study took place. ML is a consultant to the parent company of the sponsor. TG, RAR and SW receive consulting fees from InovoBiologic. RG is the owner of the Factors Group of Nutritional Companies and retains an interest in PGX. PGX, PolyGlycopleX are trademarks of InovoBiologic. All other marks are the property of their respective owners.

# **Acknowledgements**

We wish to thank the efforts of Ms Tracey Wood and Ms Suzana Damjanovic, whose support and organisational expertise were invaluable during the completion of this study, and Dr Natalie Kacinik for editorial assistance in the completion of this manuscript. We would also like to acknowledge support of the study and supply of PGX by InovoBiologic.

#### References

- 1 Flegal KM, Graubard BI, Williamson DF, Gail MH. Excess deaths associated with underweight, overweight, and obesity. *JAMA* 2005; **293**: 1861–1867.
- 2 Gregg EW, Cheng YJ, Cadwell BL, Imperatore G, Williams DE, Flegal KM et al. Secular trends in cardiovascular disease risk factors according to body mass index in US adults. JAMA 2005; 293: 1868–1874.
- 3 Blackburn GL. Effect of degree of weight loss on health benefits. *Obes Res* 1995; 3: 211S–216S.
- 4 Hill JO. Understanding and addressing the epidemic of obesity: an energy balance perspective. *Endocr Rev* 2006; 27: 750–761.
- 5 National Heart, Lung, and Blood Institute Expert Panel on the Identification, Evaluation, and Treatment of Overweight and Obesity in Adults. Executive summary of the clinical guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. *Arch Intern Med* 1998; 158: 1855–1867.
- 6 Womble LG, Williamson DA, Greenway FL, Redmann SM. Psychological and behavioural predictors of weight loss during drug treatment for obesity. *Int J Obes Relat Metab* 2001; 25: 340–345.
- 7 Doucet E, St-Pierre S, Almeras N, Tremblay A. Relation between appetite ratings before and after a standard meal and estimates of daily energy intake in obese and reduced obese individuals. *Appetite* 2003; **40**: 137–143.
- 8 Delzenne NM, Cani PD. A place for dietary fiber in the management of the metabolic syndrome. *Curr Opin Clin Nutr Metab Care* 2005; 8: 636–640.
- 9 Howarth NC, Saltzman E, Roberts SB. Dietary fiber and weight regulation. *Nutr Rev* 2001; **59**: 129–139.
- 10 Wanders AJ, van den Borne JJGC, de Graff C, Hulshof T, Jonathan MC, Kristensen M et al. Effects of dietary fibre on subjective appetite, energy intake and body weight: a systematic review of randomized controlled trial. Obes Rev 2011; 12: 724–739.
- 11 Mattes RD, Rothacker D. Beverage viscosity is inversely related to postprandial hunger in humans. *Physiol Behavior* 2001; 74: 551–557.
- 12 Delargy HJ, Burley VJ, O'Sullivan KR, Fletcher RJ, Blundell JE. Effects of different soluble: insoluble fiber ratios at breakfast on 24-h pattern of dietary intake and satiety. Eur J Clin Nutr 1995; 49: 754–766.
- 13 Delargy HJ, O'Sullivan KR, Fletcher RJ, Blundell JE. Effects of the amount and type of dietary fiber (soluble and insoluble) on short term control of appetite. *Int J Food Sci Nutr* 1997; **48**: 67–77.
- 14 Chow JM, Choe YS, Noss MJ, Robinson KJ, Dugle JE, Acosta SH *et al.* Effect of a viscous fiber-containing nutrition bar on satiety of patients with type 2 diabetes. *Diabetes Res Clin Pract* 2007; **76**: 335–340.
- 15 Astrup A, Vrist E, Quaade F. Dietary fiber added to very low calorie diet reduces hunger and alleviates constipation. *Int J Obes* 1990; 14: 105–112.
- 16 Pasman WJ, Saris WHM, Wauters MAJ, Westerterp-Plantenga MS. Effect of one week fiber supplementation on hunger and satiety ratings and energy intake. *Appetite* 1997; 29: 77–87.
- 17 Jenkins DJ, Wolever TM, Leeds AR, Gassull MA, Haisman P, Dilawari J *et al.* Dietary fibers, fiber analogues, and glucose tolerance: importance of viscosity. *Br Med J* 1978; 1: 1392–1394.
- 18 Ellis PR, Apling EC, Leeds AR, Bolster NR. Guar bread: acceptability and efficacy combined. Studies on blood glucose, serum insulin and satiety in normal subjects. *Br J Nutr* 1981; **46**: 267–276.
- 19 Vuksan V, Panahi S, Lyon M, Rogovik A, Jenkins A, Leiter L. Viscosity of fiber preloads affects food intake in adolescents. Nutr Metab Cardiovasc Dis 2009; 19: 498–503.
- 20 Cappelleri JC, Bushmakin AG, Gerber RA, Leidy NK, Sexton CC, Lowe MR et al. Psychometric analysis off the three-factor eating



- questionnaire-R21: results from a large diverse sample of obese and non-obese participants. *Int J Obes* 2009; 33: 611–620.
- 21 Blundell JE, Burley VJ. Satiation, satiety and the action of fiber on food intake. *Int J Obes* 1987; 11: 9–25.
- 22 Abdelhameed AS, Ang A, Morris GA, Smith I, Lawson C, Gahler R *et al.* An analytical ultracentrifuge study on ternary mixtures of konjac glucomannan supplemented with sodium alginate and xanthan gum. *Carbohydr Polym* 2010; **81**: 141–148.
- 23 Harding SE, Smith IH, Lawson CJ, Gahler RJ, Wood S. Studies on macromolecular interactions in ternary mixtures of konjac glucomannan, xanthan gum and sodium alginate. *J Carbpol* 2010; 10: 1016–1020.
- 24 Matulka R, Lyon M, Wood S, Marone P, Merkel D, Burdock G. The safety of PolyGlycopleX<sup>®</sup> (PGX<sup>®</sup>) as shown in a 90-day rodent feeding study. *Nutr J* 2009; **8**: 1–11.
- 25 Carabin I, Lyon M, Wood S, Pelletier X, Donazzolo Y, Burdock G. Supplementation of the diet with the functional fiber PolyGlycoplex<sup>®</sup> is well tolerated by healthy subjects in a clinical trial. *Nutr J* 2009; **8**: 9.
- 26 Marone P, Lyon M, Gahler R, Donath C, Hofman-Huther H, Wood S. Genotoxicity studies of PolyGlycopleX<sup>®</sup> (PGX<sup>®</sup>): a novel dietary fiber. *Int J Toxicol* 2009; **28**: 318–331.
- 27 Boon H, Clitheroe J, Forte T. Effects of greens+<sup>®</sup>: a randomized, controlled trial. *Can J Diet Prac Res* 2004; **65**: 66–71.
- 28 Bijkerk CJ, de Wit NJ, Muris JWM, Whorwell PJ, Knottnerus JA, Hoes AW. Soluble or insoluble fiber in irritable bowel syndrome in primary care? Randomised placebo controlled trial. *Br Med J* 2009; 339: b.3154.
- 29 Niba LL. Processing effects on susceptibility of starch to digestion in some dietary starch sources. *Int J Food Sci Nutr* 2003; **54**: 97–109.
- 30 Brennan IM, Feltrin KL, Nair NS, Hausken T, Little TJ, Gentilcore D et al. Effects of the phases of the menstrual cycle on gastric emptying, glycemia, plasma GLP-1 and insulin, and energy intake in healthy lean women. Am J Physiol Gastrointest Liver Physiol 2009; 297: G602–G610.
- 31 Stubbs RJ, Hughes DAQ, Johnstone AM, Rowley E, Reid C, Elia M *et al.* The use of visual analogue scales to assess motivation to eat in human subjects: a review of their reliability and validity with an evaluation of new hand-held computerized systems for temporal tracking of appetite ratings. *Br J Nutr* 2000; **84**: 405–415.
- 32 Blundell J, De Graaf C, Hulshof T, Jebb S, Livingstone B, Lluch A *et al.* Appetite control: methodological aspects of the evaluation of foods. *Obes Rev* 2010; **11**: 251–270.
- 33 Jones, B, Kenward, M. *Design and Analysis of Cross-Over Trials*, 2nd edn. Chapman & Hall/CRC: New York, USA 2003.
- 34 Bellisle F. Impact of the daily meal pattern on energy balance. *Food Nutr Res* 2004; **48**: 114–118.
- 35 Reimer RR, McBurney MI. Dietary fiber modulates intestinal proglucagon messenger ribonucleic acid and postprandial secretion of glucagon-like peptide-1 and insulin in rats. *Endocrinology* 1996; **137**: 3948–3956.
- 36 Sleeth ML, Thompson EL, Ford HE, Zac-Varghese SEK, Frost G. Free fatty acid receptor 2 and nutrient sensing: a proposed role for fibre, fermentable carbohydrates and short-chain fatty acids in appetite regulation. *Nutr Res Rev* 2010; 23: 135–145.
- 37 Grover GJ, Koetzner L, Wicks J, Gahler RJ, Lyon MR, Reimer RA *et al.* Effects of the soluble fiber complex PolyGlycopleX<sup>®</sup> (PGX<sup>®</sup>) on glycemic control, insulin secretion, and GLP-1 levels in Zucker diabetic rats. *Life Sci* 2011; **88**: 392–399.
- 38 Reimer RA, Pelletier X, Carabin IG, Lyon M, Gahler R, Parnell JA *et al.* Increased plasma PYY levels following supplementation with the functional fiber PolyGlycopleX in healthy adults. *Eur J Clin Nutr* 2010; 64: 1186–1191.
- 39 Le Roux CW, Batterham RL, Aylwin SJ, Patterson M, Borg CM, Wynne KJ et al. Attenuated peptide PYY release in obese subjects is associated with reduced satiety. Endocrinology 2006; 147: 3–8.

- 40 Battherham RL, Cohen MA, Ellis SM, Le Roux CW, Withers DJ, Frost GS et al. Inhibition of food intake in obese subjects by peptide YY3-36. New Engl J Med 2003; 349: 941–948.
- 41 Marciani L, Gowland PA, Spiller RC, Manoj P, Moore RJ, Young P *et al.* Effect of meal viscosity and nutrients on satiety, intragastric dilution, and emptying assessed by MRI. *Am J Physiol Gastrointest Liver Physiol* 2001; **280**: G1227–G1233.
- 42 Maljaars PW, Peters HP, Mela DJ, Masclee AA. Ileal brake: a sensible food target for appetite control. A review. *Physiol Behav* 2008: **95**: 271–281.
- 43 Van Citters GW, Lin HC. Ileal brake: neuropeptidergic control of intestinal transit. *Curr Gastroenterol Rep* 2006: 8: 367–373.
- 44 Jenkins DJ, Wolever TM, Taylor RH, Barker H, Fielden H, Baldwin JM *et al.* Glycemic index of foods: a physiological basis for carbohydrate exchange. *Am J Clin Nutr* 1981; 34: 362–366.
- 45 Jenkins AL, Kacinik V, Lyon M, Wolever TM. Effect of adding the novel fiber, PGX<sup>®</sup>, to commonly consumed foods on glycemic response, glycemic index and GRIP: a simple and effective strategy for reducing post prandial blood glucose levels—a randomized, controlled trial. *Nutr J* 2010; 22: 9:58.
- 46 Brand-Miller JC, Atkinson FS, Gahler RJ, Kacinik V, Lyon MR, Wood S. Effects of PGX, a novel functional fibre, on acute and delayed postprandial glycemia. Eur J Clin Nutr 2010; 64: 1488–1493.
- 47 Arumugam V, Lee J-S, Nowal JK, Pohle RJ, Nyrop JE, Leddy J *et al.* A high-glycemic meal pattern elicited increased subjective appetite sensations in overweight and obese women. *Appetite* 2008; 50: 215–222.
- 48 Dewan S, Gillett A, Mugarza JA, Dovey TM, Halford JC, Wilding JP. Effects of insulin-induced hypoglycemia on energy intake and food choices at a subsequent test meal. *Diabetes Metab Res Rev* 2004; **20**: 405–410.
- 49 Holt S, Brand Miller JC, Petoccz P. Interrelationship among postprandial satiety, glucose, and insulin responses and changes in subsequent food intake. *Eur J Clin Nutr* 1996; **50**: 788–797.
- 50 Anderson GH, Catherine NL, Woodend DM. Inverse association between the effect of carbohydrates on blood glucose and subsequent short-term food intake in young men. *Am J Clin Nutr* 2002; **76**: 1023–1030.
- 51 Bornet FRJ, Jardy-Gennetier A-E, Jacquet N, Stowell J. Glycemic response to foods: impact on satiety and long-term weight regulation. *Appetite* 2007; **49**: 535–553.
- 52 Flint A, Raben A, Blundell JE, Astrup A. Reproducibility, power and validity of visual analogue scales in assessment of appetite sensations in single test meal studies. *Int J Obes* 2000; **24**: 38–48.
- 53 De Graaf C. The validity of appetite ratings. Appetite 1993; 21: 156–160.
- 54 De Castro J. Independence of genetic influences on body size, daily intake, and meal patterns in humans. *Psychol Behav* 1993; 54: 633–639.
- 55 Jéquier E, Tappy L. Regulation of body weight in humans. *Psychol Rev* 1999; **79**: 451–480.
- 56 Herman CP, Ostovich JM, Polivy J. Effects of attentional focus on subjective hunger ratings. *Appetite* 1999; 33: 181–193.
- 57 Drewnowski A. Energy density, palatability, and satiety: implications for weight control. *Nutr Rev* 1998; **56**: 347–353.
- 58 Rogers PJ, Schutz HG. Influence of palatability on subsequent hunger and food intake: a retrospective replication. *Appetite* 1992; **19**: 155–156.
- 59 Womble LG, Wadden TA, Chandler JM, Martin AR. Agreement between weekly vs. daily assessment of appetite. *Appetite* 2003; **40**: 131–135.



This work is licensed under the Creative Commons Attribution-NonCommercial-No

Derivative Works 3.0 Unported License. To view a copy of this license, visit http://creativecommons.org/licenses/by-nc-nd/3.0/

Appendix 1 The nutritional composition of the three-day low calorie diet

	Quantity	Calories	Carbohydrate (g)	Fibre (g)	Protein (g)	Fat(g)
Day 1		1004	156	7	55	16
Breakfast		244	38		8	5
Blue Menu granola cereal	0.5 cup	165	30	3	3	2
Milk 2%	150 ml	77	7		5	3
Tea or Coffee brewed	1 cup	2	0.7		0	0
Lunch	·	340	58		20	3
Blue Menu Chicken Bangkok	1 pkg	230	32	2	19	3
Juice, Orange	1 cup	110	26		2	0
Dinner	·	420	60		27	8
Blue Menu Chinese Sweet & Sour Chicken	1 pkg	350	60	2	22	2
Marble cheese stick	1	70	0		5	6
Day 2		1002	157	7	54	15
Breakfast		249	41		9	2
Maple Brown Sugar Oatmeal	1 pkg	170	33	3	4	2
Milk 2%	150 ml	77	7		5	0
Tea or Coffee brewed	1 cup	2	0.7		0	0
Lunch	·	333	57		22	2
Blue Menu Ginger Glazed Chicken	1 pkg	220	32	2	19	2
Blue Menu Multigrain Pretzels	30 g	113	25	1	3	0.3
Dinner	•	420	59		23	11
Blue Menu Thai Sweet Chili Lemon Grass Chicken	1 pkg	350	59	1	18	5
Marble cheese stick	1 piece	70	0		5	6
Day 3		1005	166	14	45	20
Breakfast		246	45		9	5
Blue Menu Oatmeal Bagel	1	160	30	5	7	2
Blue Menu Old-Fashioned Peanut Butter	1 tsp	33	1	0.3	1	3
Tea or Coffee brewed	1 cup	2	0.7		0	0
Banana, fresh, medium	1/2	53	13	1.5	1	0.2
Lunch		346	58		14	6
Blue Menu Roasted Vegetable Lasagna	1 pkg	240	33	3	12	6
Banana, fresh, medium	1/2	53	13	1.5	1	0.2
Carrots, baby, fresh	1 cup	53	12	2.5	1	0
Dinner		413	63		22	9
Blue Menu Indian Butter Chicken	1 pkg	370	52	2	21	9
Orange, medium	0.5	43	11	2	1	0

Abbreviations: pkg, package; tsp, teaspoon.