

CORRECTION



Correction: Comparison of two families with and without ataxia harboring novel variants in *PRKCG*

Yui Tada, Kodai Kume, Soma Noguchi, Tomoko Sekiya, Kazuto Nishinaka, Hiroshi Ishiguchi, Jinsoo Koh, Seiji Emori, Yoshiaki Nakayama , Takashi Kurashige , Yuishin Izumi, Hidefumi Ito, Norio Sakai and Hideshi Kawakami 

© The Author(s), under exclusive licence to The Japan Society of Human Genetics 2022

Journal of Human Genetics (2022) 67:621; <https://doi.org/10.1038/s10038-022-01064-7>

Correction to: *Journal of Human Genetics* <https://doi.org/10.1038/s10038-022-01057-6>, published online 28 June 2022

In this article, refs. [9–12] were incorrect. They should have been as below.

REFERENCES

9. Adachi N, Kobayashi T, Takahashi H, Kawasaki T, Shirai Y, Ueyama T, et al. Enzymological analysis of mutant protein kinase Cgamma causing spinocerebellar

ataxia type 14 and dysfunction in Ca²⁺ homeostasis. *J Biol Chem.* 2008;283:19854–63.

10. Wong MMK, Hoekstra SD, Vowles J, Watson LM, Fuller G, Németh AH, et al. Neurodegeneration in SCA14 is associated with increased PKCγ kinase activity, mislocalization and aggregation. *Acta Neuropathol Commun.* 2018;6:99.

11. Matsuda Y, Morino H, Miyamoto R, Kurashige T, Kume K, Mizuno N, et al. Biallelic mutation of HSD17B4 induces middle age-onset spinocerebellar ataxia. *Neurol Genet.* 2020;6:e396.

12. Kume K, Morino H, Miyamoto R, Matsuda Y, Ohsawa R, Kanaya Y, et al. Middle-age-onset cerebellar ataxia caused by a homozygous TWNK variant: a case report. *BMC Med Genet.* 2020;21:68.