



CORRECTION

Correction: Prefrontal GABA levels, hippocampal resting perfusion and the risk of psychosis

Gemma Modinos^{1,2}, Fatma Şimşek¹, Matilda Azis¹, Matthijs Bosson³, Ilaria Bonoldi¹, Carly Samson¹, Beverly Quinn⁴, Jesus Perez^{4,5,6}, Matthew R. Broome^{7,8}, Fernando Zelaya², David J. Lythgoe², Oliver D. Howes¹, James M. Stone^{1,2}, Anthony A. Grace⁹, Paul Allen^{1,10} and Philip McGuire¹

Neuropsychopharmacology (2018) 43:2660; <https://doi.org/10.1038/s41386-018-0118-5>

Correction to: *Neuropsychopharmacology*; <https://doi.org/10.1038/s41386-017-0004-6>; published online 30 January 2018.

This article was originally published under NPG's License to Publish, but has now been made available under a [CC BY 4.0] license. The PDF and HTML versions of the paper have been modified accordingly.

¹Department of Psychosis Studies, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK; ²Department of Neuroimaging, Institute of Psychiatry, Psychology & Neuroscience, King's College London, London, UK; ³Department of Psychiatry, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, Netherlands; ⁴CAMEO Early Intervention in Psychosis Service, Cambridgeshire and Peterborough NHS Foundation Trust, Cambridge, UK; ⁵Department of Psychiatry, University of Cambridge, Cambridge, UK; ⁶Department of Neuroscience, Instituto de Investigacion Biomedica de Salamanca (IBSAL), University of Salamanca, Salamanca, Spain; ⁷Department of Psychiatry, University of Oxford, Oxford, UK; ⁸Oxford Health NHS Foundation Trust, Oxford, UK; ⁹Department of Neuroscience, Psychiatry and Psychology, University of Pittsburgh, Pittsburgh, PA, USA and ¹⁰Department of Psychology, University of Roehampton, Roehampton, UK
Correspondence: Gemma Modinos (gemma.modinos@kcl.ac.uk)