

CORRECTION OPEN



Correction to: Identification of tRF-29-79MP9P9NH525 as a biomarker and tumor suppressor of gastric cancer via regulating KIF14/AKT pathway

Jiaxin Ge, Ji Dai, Haoqiang Ji, Jie Guo , Xiaoban Shen, Desen Sun, Qiang Chen, Pan Chen , Guoliang Ye, Junming Guo and Shuangshuang Zhang

© The Author(s) 2025

Cell Death Discovery (2025)11:353; <https://doi.org/10.1038/s41420-025-02614-6>

Correction to: *Cell Death Discovery* <https://doi.org/10.1038/s41420-025-02514-9>, published online 15 May 2025

In this article on page 4, lines 17 and 18 of Table 1: The correct number of “NO” situation in the high group is “18”; the correct number of “N1 & N2&N3” situation in the high group is “41”; the correct number of “NO” situation in the low group is “19”; the correct number of “N1&N2&N3” situation in the low group is “41”. The percentage and the total number (n) in the column 2 have been corrected accordingly.

Line 21 of Table 1: The correct percentage of low group is “10.00”.

Line 25 of Table 1: The correct percentage of low group is “36.67”.

Line 30 of Table 1: The correct percentage of low group is “20.00”.

Line 44 of Table 1: The correct content is “>30”.

Table 1. The correlation between the amount of tRF-29-79MP9P9NH525 in tissues and clinicopathological features of advanced gastric cancer patients.

Characteristics	n (%)	High (%)	Low (%)	P-value
All cases	119	59	60	
Gender				>0.9999
Male	71 (59.66)	35 (59.32)	36 (60.00)	
Female	48 (40.34)	24 (40.68)	24 (40.00)	
Age (y)				0.1627
≥60	83 (69.75)	45 (76.27)	38 (63.33)	
<60	36 (30.25)	14 (23.73)	22 (36.67)	
Differentiation				0.0107
Well	9 (7.56)	5 (8.48)	4 (6.67)	
Moderate	68 (57.14)	41 (69.49)	27 (45.00)	
Poor	42 (35.30)	13 (22.03)	29 (48.33)	
Lymphatic node metastasis				>0.9999
N0	37 (31.09)	18 (30.51)	19 (31.67)	
N1 & N2 & N3	82 (68.91)	41 (69.49)	41 (68.33)	
Invasion				0.1563
T2	39 (32.77)	20 (33.90)	19 (31.67)	
T3	7 (5.88)	1 (1.69)	6 (10.00)	
T4	73 (61.35)	38 (64.41)	35 (58.33)	

Published online: 30 July 2025

Official journal of CDDpress

Table 1. continued

Characteristics	n (%)	High (%)	Low (%)	P-value
Tumor size (cm)				0.0278
≥5	63 (52.94)	25 (42.37)	38 (63.33)	
<5	56 (47.06)	34 (57.63)	22 (36.67)	
Distal metastasis				0.3644
Yes	4 (3.36)	3 (5.08)	1 (1.67)	
No	115 (96.64)	56 (94.92)	59 (98.33)	
TNM stage				0.6373
IB	25 (21.00)	13 (22.03)	12 (20.00)	
IIA & IIB	24 (20.17)	10 (16.95)	14 (23.33)	
IIIA & IIIB & IIIC	66 (55.46)	33 (55.93)	33 (55.00)	
IV	4 (3.36)	3 (5.09)	1 (1.67)	
Vessel invasion				>0.9999
Yes	85 (71.43)	42 (71.19)	43 (71.67)	
No	34 (28.57)	17 (28.81)	17 (28.33)	
Lymphatic invasion				0.6883
Yes	84 (70.59)	43 (72.88)	41 (68.33)	
No	35 (29.41)	16 (27.12)	19 (31.67)	
Nerve invasion				0.0877
Yes	76 (63.87)	33 (55.93)	43 (71.67)	
No	43 (36.13)	26 (44.07)	17 (28.33)	
Ki-67 (%)				0.3689
>30	94 (78.99)	49 (83.05)	45 (75.00)	
≤30	25 (21.01)	10 (16.95)	15 (25.00)	

The original article has been corrected.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2025