

CORRECTION



Correction to: New molecular evidence on genome relationships and chromosome identification in fescue (*Festuca*) and ryegrass (*Lolium*)

I. Pašakinskienė, K. Anamthawat-Jönsson, M. W. Humphreys, V. Paplauskiene and R. N. Jones

© The Author(s), under exclusive licence to The Genetics Society 2022

Heredity (2022) 128:139–140; <https://doi.org/10.1038/s41437-021-00488-9>

Correction to: *Heredity* <https://doi.org/10.1046/j.1365-2540.1998.00446.x>, published online 1 December 1998

In Fig. 2 and Fig. 3 of this article are appear twisted and upside down the figures. The correct figures are given below.




Fig. 2 Genomic Southern hybridization using the ECL method. **(a)** Ethidium bromide-stained gels showing *Bam*HI digests of *Lolium multiflorum* (*Lm*), *F. arundinacea* (*Fa*), *F. pratensis* (*Fp*) and *F. glaucescens* (*Fg*), 1 µg per lane. **(b)** Luminograph showing hybridization of a genomic *Lm* probe to the same digests **(a)** of *Lm*, *Fa*, *Fp* and *Fg*; ECL detection 35-min. The DNA size maker is given in kb.




Fig. 3 Southern blot hybridization patterns of rDNA pTa7I to *L. multiflorum* (*Lm*), *F. arundinacea* (*Fa*), *F. pratensis* (*Fp*) and *F. glaucescens* (*Fg*), 1 µg per lane. Digests 2–5 with *Eco*RI, 6–9 *Dra*I, 10–13 *Bam*HI, track 1 for marker. ECL detection 5 min. The DNA size maker is given in kb.