

ARTICLE


<https://doi.org/10.1038/s41467-022-29911-3>

OPEN

Prospective individual patient data meta-analysis of two randomized trials on convalescent plasma for COVID-19 outpatients

Pere Millat-Martinez^{1,2,62}, Arvind Gharbharan^{3,62}, Andrea Alemany^{1,4,5,62}, Casper Rokx³, Corine Geurtsvankessel⁶, Grigorios Papageorgiou⁷, Nan van Geloven⁸, Carlijn Jordans³, Geert Groeneveld⁹, Francis Swaneveld¹⁰, Ellen van der Schoot¹¹, Marc Corbacho-Monné^{1,5,12}, Dan Ouchi^{1,13}, Francini Piccolo Ferreira¹⁴, Pierre Malchair¹⁵, Sebastian Videla^{1,16,17}, Vanesa García García¹⁵, Anna Ruiz-Comellas^{18,19,20}, Anna Ramírez-Morros¹⁸, Joana Rodriguez Codina²¹, Rosa Amado Simon²¹, Joan-Ramon Grifols^{4,22}, Julian Blanco²³, Ignacio Blanco^{4,24}, Jordi Ara^{4,25}, Quique Bassat^{2,26,27,28,29}, Bonaventura Clotet^{1,4,20,23}, Bàrbara Baro², Andrea Troxel³⁰, Jaap Jan Zwaginga^{31,32,63}, Oriol Mitjà^{1,4,20,33,63}, Bart J. A. Rijnders^{3,63}✉, CoV-Early study group* & COnV-ert study group*

Data on convalescent plasma (CP) treatment in COVID-19 outpatients are scarce. We aimed to assess whether CP administered during the first week of symptoms reduced the disease progression or risk of hospitalization of outpatients. Two multicenter, double-blind randomized trials (NCT04621123, NCT04589949) were merged with data pooling starting when <20% of recruitment target was achieved. A Bayesian-adaptive individual patient data meta-analysis was implemented. Outpatients aged ≥ 50 years and symptomatic for ≤ 7 days were included. The intervention consisted of 200–300 mL of CP with a predefined minimum level of antibodies. Primary endpoints were a 5-point disease severity scale and a composite of hospitalization or death by 28 days. Amongst the 797 patients included, 390 received CP and 392 placebo; they had a median age of 58 years, 1 comorbidity, 5 days symptoms and 93% had negative IgG antibody-test. Seventy-four patients were hospitalized, 6 required mechanical ventilation and 3 died. The odds ratio (OR) of CP for improved disease severity scale was 0.936 (credible interval (CI) 0.667–1.311); OR for hospitalization or death was 0.919 (CI 0.592–1.416). CP effect on hospital admission or death was largest in patients with ≤ 5 days of symptoms (OR 0.658, 95%CI 0.394–1.085). CP did not decrease the time to full symptom resolution.

Trial registration: Clinicaltrials.gov NCT04621123 and NCT04589949. Registration: NCT04621123 and NCT04589949 on <https://www.clinicaltrials.gov>

A full list of author affiliations appears at the end of the paper.

The unprecedented pace and amount of research on the pathogenesis of severe acute respiratory syndrome coronavirus 2 (SARS-CoV-2) led to the availability of mortality-reducing therapies within a year after the start of the coronavirus disease 2019 (COVID-19) pandemic^{1–3}. For non-hospitalized COVID-19 patients, only anti-SARS-CoV-2 monoclonal antibodies have emerged as a treatment that reduces hospital admission but only when given in the first week of illness. However, they are typically unavailable to middle and low-income countries^{4–7}.

Convalescent plasma (CP) from COVID-19 recovered patients contains polyclonal anti-SARS-CoV-2 antibodies, can be collected in large quantities at relatively low costs and was used as a therapeutic strategy in previous viral outbreaks^{8,9}. So far, randomized trials were unable to generate convincing evidence in support of CP for hospitalized patients with COVID-19^{10–18}. However, because an autologous SARS-CoV-2 antibody response typically precedes hospital admission, CP is more likely to be beneficial when it is administered very early after symptom onset¹⁹. Indeed, the only evidence from a randomized trial in favor of CP for COVID-19 comes from a small study in which elderly outpatients received CP in the first 72 h after symptom onset²⁰. In a more recent trial, CP did not reduce the risk of disease progression of COVID-19 in patients with early disease (≤ 7 days). However, in this trial, patients were recruited at emergency rooms and were, therefore, more likely to manifest severe symptoms²¹. This approach resulted in a trial profile of patients with moderate or late-stage disease, opposed to what was intended in the design. Hence, whether early treatment with CP improves the outcome of outpatients with COVID-19 remains an important question.

As soon as effective vaccines against COVID-19 became available in high-income countries, they were prioritized for individuals at higher risk for a poorer COVID-19 outcome. Because studies on CP for outpatients with COVID-19 focus on these high-risk populations as well, a high vaccination uptake will reduce the number of COVID-19 patients eligible for these studies. More importantly, the risk for a severe outcome will be small when patients become infected despite vaccination. Therefore, we anticipated that vaccination would slow down recruitment, reduce the number of events in the recruited patients and result in individual studies being underpowered. In light of the uncertainty for achieving recruitment goals, real-time pooling of individual patient data from ongoing clinical trials was proposed as a tool for providing timely data to respond to the public health crisis²². With this in mind, we initiated the Continuous Monitoring of Pooled International trials of convalescent plasma for COVID-19 patients at home Consortium (COMPILE_{home}), which provided a platform to pool individual patient data continuously and in real-time from randomized clinical trials (RCTs) on CP for outpatients with COVID-19²². This COMPILE_{home} consortium prospectively pooled and monitored the data from 2 double-blind RCTs, the CoV-Early (NCT04589949) and the CONV-ert (NCT04621123) studies, to assess the effectiveness of high-titer CP for COVID-19 outpatients.

Results

Trials profile. The search for trials resulted in 35 identified studies, thirty-one of which did not meet the selection criteria of the consortium (Supplementary Fig. 1). Of the four remaining studies, one study team opted to abstain from pooling data while another never responded to repeated emails and calls, resulting in two trials included in the pooled analysis: The CONV-ert study (NCT04621123) and the CoV-Early study (NCT04589949). The CONV-ert study received approval from the Institutional Review

Board of the Hospital Germans Trias I Pujol (reference PI 20-313) and the CoV-Early study received approval from the Institutional Review Board of the Erasmus Medical Center Rotterdam (reference MEC-2020-0682). Briefly, the CONV-ert study randomized outpatients at 4 sites in Catalunya (Spain) aged ≥ 50 years with ≤ 7 days of symptoms to one unit (200–300 mL) of CP or sterile 0.9% saline solution, both covered with opaque tubular bags for blinding investigators and patients. The CONV-ert study joined the consortium when 65 of 474 planned patients were enrolled. CoV-Early enrolled outpatients at 10 sites aged ≥ 50 years with ≤ 7 days of symptoms and at least one additional risk factor for severe COVID-19 to receive either one unit (300 mL) of CP or non-convalescent plasma (donated before 01/2020) masked to investigators and patients. It had randomized 150 of the 690 planned patients when they joined the consortium. Details about the allocation concealment, blinding and selection of CP donors in both trials can be found in Supplementary Table 2 and the study protocols.

The CONV-ert study selected the CP after being screened for high anti-SARS-CoV-2 IgG titers with ELISA (EUROIMMUN ratio ≥ 6), according to guidelines, and supplied by the regional blood bank (Banc de Sang i Teixits de Catalunya—BST); and the CoV-Early study selected the convalescent plasma based on a plaque reduction neutralization test (PRNT) 50 titer of 1:160 or higher. The two trials used a different assay to measure the titer of SARS-CoV-2 neutralizing antibodies. Therefore, a panel of 15 plasma samples was provided for comparison by the Support-E consortium, aimed at harmonizing CP evaluation in Europe²³. These results confirmed the linearity of both assays and allowed conversion of all neutralizing antibody titers into international units (IU/mL). The median neutralizing antibody titer in the plasma units was 1:386 (IQR 1:233–1:707) IU/mL, which is twice the median titer we previously observed in Dutch CP donors¹⁹. More details are described in the online Supplementary Data (page 13), in Supplementary Tables 1, 2 and in the individual study protocols.

Study patients and recruitment. Between November 2020 and July 2021, the CoV-Early and CONV-ert study teams contacted approximately 4450 outpatients with a positive SARS-CoV-2 PCR or an antigen test. The majority of exclusions occurred for one of the following reasons: few remaining or clearly improving symptoms, no comorbidities, >7 days of symptoms, unable to come to study site or declined to participate. The online supplement provides more information about the recruitment procedures of each trial.

The rapid uptake of COVID-19 vaccination in Europe, which significantly affected recruitment rate in both studies (Supplementary Fig. 2) and the authorization of specific anti-SARS-CoV-2 monoclonal antibodies for high-risk outpatients resulted in early trial termination (CONV-ert on 8th of June and CoV-Early on 13th of July 2021) following recommendations of their DSMBs. By that time, 797 participants had been enrolled and 782 of them had received the allocated intervention and could be pooled for the analysis (Fig. 1).

Patients included in the analysis had a median age of 58 years (IQR 53–64), a median of 5 days (IQR 4–6) from symptom onset, and a median of 1 comorbidity (IQR 0–2). According to the baseline assessment, 688 patients (93%) had a negative result for serum IgG anti-SARS-CoV-2 S-protein, and 21 had completed their COVID-19 vaccination. 14 participants had received one of 2 doses of a mRNA vaccine at the time of inclusion. Baseline characteristics were comparable between both study arms (Table 1).




Fig. 1 CONSORT flow diagram. Figure shows the CONSORT flow diagram of the COMPILEhome patients. 833 patients were screened at a study site and 782 were included for analysis.

Table 1 Baseline characteristics.

Characteristic	Total (n = 782)	CP ^a (n = 390)	Control (n = 392)
Male sex—no. (%)	522 (66.8%)	267 (68.5%)	255 (65.1%)
Age—median (IQR)	58 (53–64)	58 (53–64)	58 (54–65)
50–60 y	428	222	206
61–70 y	217	103	114
>70 y	82	36	46
O ₂ saturation—median (IQR) ^b	97 (96–98)	97 (96–98)	97 (96–98)
Severe immunodeficiency—no. (%)	13 (1.7%)	5 (1.3%)	8 (2.1%)
Number of comorbidities—median (IQR) ^c	1 (0–2)	1 (0–2)	1 (0–2)
0	225	111	114
1	349	171	178
2–3	192	100	92
>3	15	7	8
Days since first symptoms—median (IQR)	5 (4–6)	5 (4–6)	5 (4–6)
Positive antibody status at baseline—no. (%)	53 (7.0%)	28 (7.7%)	24 (6.4%)

^aConvalescent plasma.

^bBaseline oxygen saturation without supplementary oxygen.

^cObesity, cardiac disease, lung disease, neurological disease, diabetes, chronic renal failure, cancer and/or liver disease. See the Supplementary Appendix for additional details of the comorbidities.

Primary endpoints. Table 2 and Fig. 2 show the distribution of patients across the five categories of the disease severity scale. The overall estimated OR for patients treated with CP was 0.936 (posterior mean, 95% credible interval 0.667–1.311) with a 64.9% posterior probability of benefit (OR < 1). Hospital admission or

death occurred in 34 of 390 (8.7%) patients treated with CP and in 40 of 392 (10.2%) patients in the control arm with an OR of 0.919 (posterior mean, 95% credible interval 0.592–1.416) and a 64.3% posterior probability of benefit. Although being included in the CONV-ert trial was associated with a poorer overall outcome,

Table 2 Distribution of the outcome of the patients in the 28 days after inclusion across the 5-points disease severity scale.			
Worst disease severity score	Total (n = 782)	CP ^a (n = 390)	Control (n = 392)
Recovered before day 8 after transfusion—no. (%) ^b	143 (18.3%)	74 (19.0%)	69 (17.6%)
Continued symptoms after day 7—no. (%) ^c	565 (72.3%)	282 (72.3%)	283 (72.2%)
Admitted to hospital but no invasive ventilation needed—no. (%)	65 (8.3%)	31 (7.9%)	34 (8.7%)
Admitted to hospital and invasive ventilation needed—no. (%)	6 (0.8%)	2 (0.5%)	4 (1.0%)
Death—no. (%)	3 (0.4%)	1 (0.3%)	2 (0.5%)

^aConvalescent plasma.
^bRecovered with no symptoms related to COVID-19 within 7 days after inclusion.
^cContinued symptoms attributable to COVID-19.




Fig. 2 Distribution for COVID-19 severity at 28 days. CP Convalescent plasma. Figure shows the distribution of the outcome of the patients in the 28 days after inclusion across the 5-point disease severity scale: 1 = recovered before day 8 after transfusion, 2 = continued symptoms after day 7, 3 = hospital admission, 4 = invasive ventilation, 5 = death. Moving from lighter to darker shading represents increasing scores on the severity scale. The darker shade includes point 4 and 5 of the scale (invasive ventilation or death).

the effect of CP was similar in both trials. This increased risk for patients in CONV-ert was independent of age, sex, and the number of comorbidities. The results of all covariates included in the primary analysis can be found in the online supplement (Supplementary Figs. 5, 6).

Secondary endpoints. No differences between CP and control patients regarding time to complete resolution of COVID-19 symptoms was seen (log-rank $p = 0.66$, Fig. 3). The effect size of CP on the binary outcome of hospital admission or death was larger in patients with ≤ 5 days of symptoms (OR 0.658, 95% CI 0.394–1.085) compared to those with >5 days (OR 1.427, 95% CI 0.789–2.580) and comparable results were observed for the ordinal outcome (OR 0.720, 95% CI 0.486–1.064 in the early treated group, Supplementary Figs. 7, 8).

Finally, the ORs for patients who received CP with neutralizing antibody titers above or below the median titer were nearly identical (Supplementary Fig. 9). Also, no notable difference was

observed when patients with IgG anti-SARS-CoV-2 antibodies detected at baseline were excluded (OR 0.880, 95% CI 0.590–1.310 for the binary outcome, OR 0.892, 95% CI 0.643–1.236 for ordinal outcome, Supplementary Figs. 7, 8).

Safety. The intervention was well-tolerated. 89 serious adverse events (SAE) were reported, 4 were considered related to the plasma transfusion (3 in the control arm). Three patients could leave the hospital <24 h after transfusion while the fourth was hospitalized for 5 days 1 week after the CP transfusion and diagnosed with thrombophlebitis at the infusion site and a pulmonary embolism (Table 3).

Discussion

In this analysis of 782 patients with COVID-19 randomized to high-titer CP or placebo within 7 days of disease onset, treatment with CP did not prevent COVID-19 progression, hospitalization, or other clinical outcomes. Our results agree with those by Korley




Fig. 3 Time to full symptom resolution up to day 28 (end of follow-up). CP Convalescent plasma. Log-rank test $p = 0.66$. The dotted error bars represent the 95% CI.

Table 3 Serious adverse events ^a .			
SAE category	Total	CP ^b	Control
(Prolongation of) hospital admission—no. ^c	80	37	43
Death—no.	3	1	2
Serious transfusion related adverse event—no. ^d	4	1	3
Life threatening transfusion reaction—no. ^e	2	0	2
Other AE	2	1	1

^aSerious adverse events (SAE) were registered in all patients that signed the informed consent form ($n = 797$) regardless of being transfused or not.
^bConvalescent plasma.
^cWhen a patient is hospitalized more than once, each admission is counted separately.
^dAny transfusion reaction associated with a plasma transfusion that was considered as a SAE.
^e2 patients with anaphylaxis very soon after discharge that required urgent therapy by paramedics.

et al. in patients of the same age and symptom duration but with probably more severe symptoms as they were recruited at emergency rooms in the USA²¹. These findings differ from those of a smaller trial that used CP within 72 h of symptom onset in much older patients (≥ 75 years)²⁰. We explored signs of efficacy in various subgroups most likely to benefit from CP. The only subgroup in our study that we found that could potentially benefit from CP was the subgroup with ≤ 5 days from the onset of symptoms (OR 0.70, CI 0.47–1.03). The potential effect of CP when administered early after disease onset has been suggested by other authors²⁴, and could explain the results reported by Libster et al. study²⁰. However, in our study this was a secondary endpoint and the confidence interval was wide, so confirmation in other studies is needed. Regarding the safety parameters of this strategy, our study shows no major concerns, with only four SAEs related to the plasma infusion; these findings are in line with those described in previous studies²⁵.

Our study has several strengths. It is the largest of its kind, studying the effect of CP for high-risk outpatients with COVID-19 early after initiation of symptoms. The fact that 93% of all

patients were SARS-CoV-2 antibody negative at the time of inclusion confirms that they were recruited in the early stage of the disease. Pooling of the data from both studies was pre-planned and initiated before any interim analyses were performed and when both studies were early in their recruitment. Both teams remained fully blinded as the (interim) analyses were done by an unblinded statistical team that shared the results with the DSMB on a regular basis. The COMPILEhome study used the same primary endpoint as the CoV-Early trial, and therefore we did not perform a separate sample size calculation. As our assumptions about the outcome across the ordinal scale were somewhat different than anticipated in the original sample size calculation (fewer hospitalizations and deaths in particular), we repeated the calculation of the effect size that our study was powered for post-hoc. This showed that our study still had 80% power to detect an odds ratio of 0.65 for the primary endpoint, very close to the original power calculation. We, therefore, consider our results methodologically sound.

Several limitations should be mentioned. Although we only included patients aged ≥ 50 , and most of them also had

comorbidities, the hospital admission and death rates were relatively low at 9.5%. Therefore, the study was not powered to exclude a small overall treatment effect on these endpoints. However, administering CP to infectious and symptomatic outpatients is complex and labor-intensive. Hence, we think that CP's clinical role is significantly diminished if unable to establish something greater than "a small effect" because it ceases to be practical. The contribution of the individual comorbidities to COVID-19 risk in our study should be interpreted cautiously because, owing to the lack of consensus regarding the relative relevance of each of them, we summed them in a non-weighted fashion. As vaccination uptake progressed in patients aged 50 or older and monoclonal antibody-based therapy with proven effectiveness in high-risk outpatients became available, the recruitment dropped dramatically as of June 2021. This resulted in the recommendation by the individual and COMPILE_{home} DSMBs that further enrollment was unlikely to change the results, and both studies were discontinued. Regarding the advent of the SARS-CoV-2 variants that may be less susceptible to antibodies induced by the original SARS-CoV-2 virus or the alpha variant, it is reassuring that >95% of the patients in both countries were included at a time when the delta variant was still rare (<5%) (Supplementary Figs. 3, 4). The last limitation of our study (and all studies on CP for COVID-19 so far) is the lack of a proper phase 2 dose-finding study. In a recent study, we administered 600 mL of CP to 25 SARS-CoV-2 antibody-negative B-cell depleted patients diagnosed with COVID-19²⁶. While all seroconverted immediately after transfusion, the median virus neutralization titer only rose to 1:40. This is 4 times lower than the median titer in immunocompetent convalescent COVID-19 patients and up to 100 times lower than titers observed after treatment with monoclonal antibodies^{7,19}. Therefore, we postulate that the range of neutralizing antibody titers present in the 200–300 mL of plasma we used may well have been too low. That underdosing may partially explain our findings is also suggested by a study in which human CP with a neutralizing antibody titer of 1:320 did not prevent disease in hamsters while a titer of 1:2560 did²⁷. Hence, we recommend that any future study on CP for COVID-19 should use donors at the upper extreme end of antibody titers (e.g., >1:2560 IU). Although, this was virtually impossible in 2020, this should no longer be difficult now as plasma donors recently vaccinated or boosted with a mRNA SARS-CoV-2 vaccine can be selected.

Last but not least, a recent preprint publication by Sullivan et al. described the results of the CSSC04 study in which 1181 outpatients received one unit of convalescent or control plasma. In this trial, CP lowered the risk of hospital admission or death from 6.3 to 2.9%, $p = 0.004$ ²⁸. Therefore, the limited impact on hospital admission or death in our study should be interpreted in the context of this trial.

In conclusion, treatment of COVID-19 with CP in the first 7 days after symptom onset did not improve the outcome. Proper dose-finding studies should be conducted, preferentially in patients with ≤5 days of symptoms before future phase 3 studies on CP are initiated.

Methods

Overview of study design and research partners. Beginning in November 2020, we systematically searched for RCTs recruiting outpatients that compared treatment with CP with a blinded or unblinded control arm in the European (<https://www.clinicaltrialsregister.eu/>) and American (www.clinicaltrials.gov) trial register. Search terms were convalescent plasma, COVID-19, phase 2 or phase 3, adult, and recruiting or not recruiting. Studies were selected if they were RCTs on outpatients, if their inclusion criteria were confined to patients who had symptoms less than 7 days, and if they had a planned sample size of at least 100 participants of age 50 or older. Investigators of qualifying trials were contacted and informed about COMPILE_{home} and invited to collaborate in the study.

The full COMPILE_{home} protocol is available as an online supplement. The study was designed as a Bayesian-adaptive individual patient data meta-analysis of ongoing clinical trials. Prior to the start of pooling, the study teams agreed upon a minimal set of data required to analyze the primary and secondary endpoints was agreed upon by the study teams. Each trial provided updated data every 6 weeks. The pooled data were monitored by 2 unblinded statisticians and a data and safety monitoring board (DSMB) every 6 weeks using a pre-established stopping guideline for efficacy. At each interim analysis, a posterior distribution of the treatment effect was estimated.

Study patients and selection criteria. Although the exact inclusion and exclusion criteria could vary across the trials, all the subjects had to fulfill the following criteria; (1) Participant of a trial that joined the COMPILE_{home} consortium, (2) Confirmed COVID-19 diagnosis by a diagnostic PCR or antigen test of <7 days, (3) Neither hospitalized nor at the emergency room department of a hospital before or at the time of randomization, (4) Symptomatic with illness onset ≤7 days at the time of screening for the study defined by a physician with a complete clinical history, and (5) Age 50 or older. Trials had to be approved by the institutional review boards, and competent authorities of the countries involved, and all patients gave written informed consent.

Intervention. To qualify for COMPILE_{home}, participants randomly assigned to the experimental group had to receive an infusion of ABO-compatible CP with high antibody titers as determined via a semiquantitative antibody test against the spike protein or a virus neutralization assay. Only trials in which the participants were masked for the intervention were included.

Outcomes. Two primary efficacy outcome variables were selected. The first primary endpoint incorporated the speed of recovery as well as the risk of hospital admission, ICU admission or death in a 5-point ordinal scale. It was defined as the highest score on a 5-point ordinal disease severity scale within the 28 days after randomization. A patient scored 1 if he/she recovered quickly (i.e., fully recovered within seven days after transfusion), 2 when continued symptoms attributable to COVID-19 were present on day seven, 3 when admission to a hospital was required at any point within 28 days, 4 when invasive ventilation was required at any point within 28 days, and 5 when the patient had died at any point within 28 days. This means that the best outcome (ordinal scale score of 1) is given when a patient is fully recovered before day 8 and was never hospitalized nor died in the 28 days after transfusion, while a patient who recovered after day 7 but was never hospitalized nor died in the 28 days scored a 2 on the scale. The second primary endpoint was the occurrence of hospitalization or death within 28 days. Secondary endpoints were time to full symptom resolution (assessed by the blinded study team during a telephone contact on day 7, day 14, and day 28) and the safety of CP in outpatients with COVID-19. Pre-planned subgroup analyses assessed the efficacy of the 2 primary outcomes in the following subgroups: (1) days since disease onset (1–5 or >5 days), (2) level of neutralizing antibody anti-SARS-CoV-2 titers in transfused plasma, and (3) Negative serum anti-SARS-CoV-2 IgG status (Trimeric Spike antibody test, Liaison, Diasorin, Saluggia, Italy).

Statistical analysis. The first primary endpoint was analyzed with a Bayesian proportional odds model with normally distributed priors. The model included a main treatment effect shared among the trials (using a skeptical (i.e., conservative) standard deviation of 0.4), main trial effects (using standard deviation 0.5 for the prior distribution), and trial by treatment interactions (using a standard deviation of 0.14 for the prior distribution). The following covariates were included with a standard deviation of 0.5 for the prior distribution of their effects: age, sex, number of comorbidities (0–9), oxygen saturation at baseline (in %), immunocompromised state (Y/N) and duration of time (in days) since COVID-19 symptom onset (Supplementary Table 1). The second primary endpoint was analyzed with a Bayesian logistic model with a similar specification.

The use of the Bayesian framework and stopping rules enables continuous monitoring of the accrued data, and allowed for real-time decisions without penalties for multiple data looks associated with the classic frequentist approach. The results of each interim analysis were reported to the unblinded DSMB. The process and pre-specified thresholds for efficacy are described in detail in the protocol. The full statistical analysis plan is available as an online supplement.

The number of studies and patients included in COMPILE_{home} was not restricted and there was no pre-determined minimum or maximum sample size. The monitoring was planned to continue until the DSMB determined that there was sufficient evidence to recommend stopping the study. This situation could be achieved when the predefined stopping thresholds signaled efficacy or when the included studies had finished enrollment or any future recruitment was very unlikely to change the conclusion.

Ethical approval. The study was reviewed and approved by the institutional review boards of the Erasmus University Medical Center. The study was done according to the Helsinki Declaration of the World Medical Association. Written informed consent was obtained from every patient or legal representative. The COMPILE_{home} DSMB consisted of the chair (an infectious diseases physician), the

unblinded statisticians from the individual studies and another infectious diseases specialist. They reviewed the pooled dataset on a regular basis as described in the COMPILEhome study protocol and recommended the study team regarding the further conduct of the study. Findings are reported according to the CONSORT (Consolidated Standards of Reporting Trials) statement. For the CONV-ert study, the protocol was approved by the Ethics Committee at Hospital Germans Trias i Pujol (number PI 20-313) and the institutional review boards of participating centers. For the CoV-Early study, the protocol was approved by the medical ethical review board of the Erasmus Medical Center (METC-2020-0682).

Reporting summary. Further information on research design is available in the Nature Research Reporting Summary linked to this article.

Data availability

The source data and generated data are available in the Supplementary Data files.

Code availability

The codes generated during the current study are available from the corresponding author on reasonable request.

Received: 23 November 2021; Accepted: 7 April 2022;

Published online: 11 May 2022

References

- Horby, P. et al. Dexamethasone in hospitalized patients with Covid-19. *N. Engl. J. Med.* **384**, 693–704 (2021).
- Brown, M. J., Alazawi, W. & Kanoni, S. Interleukin-6 receptor antagonists in critically ill patients with Covid-19. *N. Engl. J. Med.* **385**, 1147 (2021).
- RECOVERY Collaborative Group. Tocilizumab in patients admitted to hospital with COVID-19 (RECOVERY): a randomised, controlled, open-label, platform trial. *Lancet* **397**, 1637–1645 (2021).
- Dougan, M. et al. Bamlanivimab plus etesevimab in mild or moderate Covid-19. *N. Engl. J. Med.* **385**(Oct), 1382–1392 (2021).
- Gottlieb, R. L. et al. Effect of bamlanivimab as monotherapy or in combination with etesevimab on viral load in patients with mild to moderate COVID-19: a randomized clinical trial. *Jama* **325**, 632–644 (2021).
- Weinreich, D. M. et al. REGN-COV2, a neutralizing antibody cocktail, in outpatients with Covid-19. *N. Engl. J. Med.* **384**, 238–251 (2020).
- Weinreich D. M. et al. REGEN-COV antibody cocktail clinical outcomes study in Covid-19 outpatients. *medRxiv* <https://doi.org/10.1101/2021.05.19.21257469> (2021).
- Garraud, O. et al. Plasma therapy against infectious pathogens, as of yesterday, today and tomorrow. *Transfus. Clin. Biol.* **23**, 39–44 (2016).
- Mair-Jenkins, J. et al. The effectiveness of convalescent plasma and hyperimmune immunoglobulin for the treatment of severe acute respiratory infections of viral etiology: a systematic review and exploratory meta-analysis. *J. Infect. Dis.* **211**, 80–90 (2015).
- Estcourt, L. J. et al. Effect of convalescent plasma on organ support-free days in critically ill patients with COVID-19: a randomized clinical trial. *Jama* **326**, 1690–1702 (2021).
- Simonovich, V. A. et al. A randomized trial of convalescent plasma in Covid-19 severe pneumonia. *N. Engl. J. Med.* **384**, 619–629 (2021).
- Agarwal, A. et al. Convalescent plasma in the management of moderate covid-19 in adults in India: open label phase II multicentre randomised controlled trial (PLACID Trial). *Bmj* **371**, m3939 (2020).
- Li, L. et al. Effect of convalescent plasma therapy on time to clinical improvement in patients with severe and life-threatening COVID-19: a randomized clinical trial. *Jama* **324**, 460–470 (2020).
- O'Donnell M. R. et al. A randomized double-blind controlled trial of convalescent plasma in adults with severe COVID-19. *J. Clin. Investig.* **131** (2021).
- Avendaño-Solà C. et al. Convalescent plasma for COVID-19: a multicenter, randomized clinical trial: *medRxiv* <https://doi.org/10.1101/2020.08.26.20182444> (2020).
- Janiaud, P. et al. Association of convalescent plasma treatment with clinical outcomes in patients with COVID-19: a systematic review and meta-analysis. *Jama* **325**, 1185–1195 (2021).
- RECOVERY Collaborative Group. Convalescent plasma in patients admitted to hospital with COVID-19 (RECOVERY): a randomised controlled, open-label, platform trial. *Lancet* **397**, 2049–2059(2021).
- Bégin, P. et al. Convalescent plasma for hospitalized patients with COVID-19: an open-label, randomized controlled trial. *Nat. Med.* **27**(Nov), 2012–2024 (2021).
- Gharbharan, A. et al. Effects of potent neutralizing antibodies from convalescent plasma in patients hospitalized for severe SARS-CoV-2 infection. *Nat. Commun.* **12**, 3189 (2021).
- Libster, R. et al. Early high-titer plasma therapy to prevent severe Covid-19 in older adults. *N. Engl. J. Med.* **384**, 610–618 (2021).
- Korley, F. K. et al. Early convalescent plasma for high-risk outpatients with Covid-19. *N. Engl. J. Med.* **385**(Nov), 1951–1960 (2021).
- Petkova, E., Antman, E. M. & Troxel, A. B. Pooling data from individual clinical trials in the COVID-19 era. *Jama* **324**, 543–545 (2020).
- Nguyen D. et al. SARS-CoV-2 neutralising antibody testing in Europe: towards harmonisation of neutralising antibody titres for better use of convalescent plasma and comparability of trial data. *Euro Surveill.* **26** (2021).
- Arnold Egloff S. A. et al. Convalescent plasma associates with reduced mortality and improved clinical trajectory in patients hospitalized with COVID-19. *J. Clin. Investig.* **131** (2021).
- Joyner, M. J. et al. Safety update: COVID-19 convalescent plasma in 20,000 hospitalized patients. *Mayo Clin. Proc.* **95**, 1888–1897 (2020).
- Gharbharan A. et al. Effects of treatment of COVID-19 with convalescent plasma in 25 B-cell depleted patients. *Clin. Infect. Dis.* **74**, 1271–1274 (2022).
- Haagmans, B. L. et al. SARS-CoV-2 neutralizing human antibodies protect against lower respiratory tract disease in a Hamster Model. *J. Infect. Dis.* **223**, 2020–2028 (2021).
- Sullivan D. J. et al. Randomized controlled trial of early outpatient COVID-19 treatment with high-titer convalescent plasma. *medRxiv* <https://doi.org/10.1101/2021.12.10.21267485> (2021).

Acknowledgements

For the CoV-Early study group. The DSMB members for COMPILEhome (Jan Nouwen, Andrea Troxel, Grigorios Papageorgiou, David Boulware, Josep Puig). All GGD contact tracers who informed potential study candidates. All infectiologists at Erasmus MC who worked even harder during COVID times to let the study team focus on the COVID trials (Adam Anas, Hannelore Bax, Mariana de Mendonça-Melo, Els van Nood, Jan Nouwen, Karin Schurink, Lennert Slobbe, Dorine de Vries-Sluijs) and the ER physicians and colleagues from the department of internal medicine who referred patients and facilitated patient recruitment. The medical students who contacted the patients (Romée Land, Eva Pruijt, Silje Taal, Liselotte Jeletich, Willem Sebrechts, Femke de Vries, Tia Rijlaarsdam). The many research assistants, research nurses and other participants who helped in facilitating the study (Maartje Wagemaker, Marita Tjaauw Joe Kim—Amadmoestar, Diane Struik, Denise Heida-Peters, Sabine Harinck, Marjo van der Poel, Greetje van Asselt, Danielle Orij—Westerhof, Marloes Romeijn, Marlies Bouterse, Pepita de Vries, Dewi Dubbelaar, Cynthia Oud, Jo Anne den Ouden, Milly Haverkort). Stichting Hemato-Oncologie voor Volwassenen Nederland (HOVON) for building the eCRF and providing real-time support with the eCRF on very short notice (Henk Hofwegen, Ronnie van der Holt, Mirjam Stomp, Marleen Luten, Monique Steijart). Professor Eva Petkova at NYU Langone who co-designed the COMPILE concept. All the persons involved in the blinding and distribution of plasma products at the sites. Last but certainly not least, all patients that participated in the trial, the thousands of plasma donors and the COVID test centers who informed patients across the Netherlands about the study. This study was made possible by a research grant from ZONMW, the Netherlands (10430062010001). Sanquin Blood Supply provided convalescent plasma free of charge for study sites in the Netherlands. **For the CONV-ert study group.** The trial was sponsored by the Fight AIDS and Infectious Diseases Foundation with funding from the pharmaceutical company Grifols S.A. and the crowdfunding campaign YoMeCorono (www.yomecorono.com). The study received support of the Hospital Universitari Germans Trias i Pujol, and Banc de Sang i Teixits de Catalunya (BST). We thank all the plasma donors and the participants in this study and all the effort they made to attend the visits for follow-up. We thank Gerard Carot-Sans for providing medical writing support with manuscript preparation and Roser Escrig for her support in the study design and medical writing assistance with the study documentation. We also thank Laia Bertran, Mireia Clua, Jordi Mitjà, Claudia Laporte, Sergi Gavilan, Miquel Àngel Rodríguez, Joan Mercado and Enric Nieto for the operational and financial management of the project. We thank the personnel from the Fight Aids and Infectious Diseases Foundation for their support in administration, human resources and supply chain management. We thank the independent DSMB for their time and dedication: Cinta Hierro (Catalan Institute of Oncology, Badalona, Spain), Natalia Tovar (Hospital Clinic, Barcelona, Spain), Binh Ngo (University of Southern California, Los Angeles, US), David Boulware (University of Minnesota, Minneapolis, US), Robin Mogg (Bill and Melinda Gates Research Institute, Seattle, US). ISGlobal receives support from the Spanish Ministry of Science and Innovation through the “Centro de Excelencia Severo Ochoa 2019–2023” Program (CEX2018-000806-S), and support from the Generalitat de Catalunya through the CERCA Program. CISM is supported by the Government of Mozambique and the Spanish Agency for International Development (AECID). BB was supported by a Beatriu de Pinós postdoctoral fellow granted by the Government of Catalonia’s Secretariat for Universities and Research, and by Marie Skłodowska-Curie Actions COFUND Program

(BP3, 801370). EP was supported by a doctoral grant from National Agency for Research and Development of Chile (ANID): 72180406. OM was supported by the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program. ZONMW, the Netherlands, grant number 10430062010001. SUP-PORT-E, grant number 101015756. YoMeCorono, www.yomecorono.com. The Fight AIDS and Infectious Diseases Foundation with funding from the pharmaceutical company Grifols S.A. The funders had no role in the design or conduct of the study; collection, management, analysis, or interpretation of the data; preparation, review, or approval of the manuscript; or decision to submit the manuscript for publication.

Author contributions

B.J.A.R. conducted the search for studies that could potentially enter the COMPILEhome consortium. B.J.A.R. and OM agreed on the terms for the COMPILEhome consortium. B.J.A.R., O.M., P.M.M., A.G., A.A., C.R., M.C.M., B.B., A.T., and J.J.Z. conceived and designed the ConV-ert and CoV-Early studies. The COMPILEhome protocol was written by B.J.A.R. and G.P. G.P. and NvG wrote the statistical analysis plan and did the statistical analysis. P.M.M., A.G., A.A., C.R., C.G., G.P., N.v.G., C.J., G.G., F.S., E.v.d.S., M.C.M., D.O., F.P.F., P.M., S.V., V.G.G., A.R.C., A.R.M., J.R.C., R.A.S., J.R.G., J.B., I.B., J.A., Q.B., B.C., B.B., A.T., J.J.Z., O.M., B.J.A.R. acquired and interpreted the data. P.M.M., A.G., A.A., O.M., G.P., and B.J.A.R. drafted the paper. P.M.M., A.G., A.A., C.R., C.G., G.P., N.v.G., C.J., G.G., F.S., E.v.d.S., M.C.M., D.O., F.P.F., P.M., S.V., V.G.G., A.R.C., A.R.M., J.R.C., R.A.S., J.R.G., J.B., I.B., J.A., Q.B., B.C., B.B., A.T., J.J.Z., O.M., B.J.A.R. critically revised the paper for important intellectual content. O.M. and B.J.A.R. were responsible for the final decision to submit for publication. All authors have seen and approved the paper.

Competing interests

The authors declare no competing interests.

Additional information

Supplementary information The online version contains supplementary material available at <https://doi.org/10.1038/s41467-022-29911-3>.

Correspondence and requests for materials should be addressed to Bart J. A. Rijnders.

Peer review information *Nature Communications* thanks Waldo H Belloso, Dean Follmann and the other, anonymous, reviewer(s) for their contribution to the peer review of this work. Peer reviewer reports are available.

Reprints and permission information is available at <http://www.nature.com/reprints>

Publisher's note Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2022, corrected publication 2024

¹Fight AIDS and Infectious Diseases Foundation, Badalona, Spain. ²ISGlobal, Hospital Clínic, Universitat de Barcelona, Barcelona, Spain. ³Department of Internal Medicine, Section of Infectious Diseases and department of Medical Microbiology and Infectious Diseases, Erasmus MC, University Medical Center, Rotterdam, The Netherlands. ⁴Hospital Universitari Germans Trias i Pujol, Badalona, Spain. ⁵Facultat de Medicina-Universitat de Barcelona, Barcelona, Spain. ⁶Department of Viroscience, Erasmus MC, Rotterdam, The Netherlands. ⁷Department of Biostatistics, Erasmus MC, University Medical Center, Rotterdam, The Netherlands. ⁸Department of Biomedical Data Sciences, Section of Medical Statistics, Leiden University Medical Center, Leiden, The Netherlands. ⁹Department of Infectious Diseases and Acute Internal Medicine, Leiden University Medical Center, Leiden, The Netherlands. ¹⁰Unit of Transfusion Medicine, Sanquin Blood Supply, Amsterdam, The Netherlands. ¹¹Department of Experimental Immunohematology, Sanquin Research, Amsterdam, The Netherlands. ¹²Hospital Universitari Parc Taulí I3PT, Sabadell, Spain. ¹³Universitat Autònoma de Barcelona, Barcelona, Spain. ¹⁴Bioclever-CRO, Barcelona, Spain. ¹⁵Emergency Department, Bellvitge University Hospital, L'Hospitalet de Llobregat, Barcelona, Spain. ¹⁶Clinical Research Support Unit (HUB-IDIBELL: Bellvitge University Hospital & Bellvitge Biomedical Research Institute), Bellvitge University Hospital, L'Hospitalet de Llobregat, Barcelona, Spain. ¹⁷Pharmacology Unit, Department of Pathology and Experimental Therapeutics, School of Medicine and 33 Health Sciences, IDIBELL, University of Barcelona, L'Hospitalet de Llobregat, Barcelona, Spain. ¹⁸Unitat de Suport a la Recerca de la Catalunya Central, Fundació Institut Universitari per a la recerca a l'Atenció Primària de Salut Jordi Gol i Gurina, Sant Fruitós de Bages, Spain. ¹⁹Health Promotion in Rural Areas Research Group, Gerència Territorial de la Catalunya Central, Institut Català de la Salut, Sant Fruitós de Bages, Spain. ²⁰Universitat de Vic—Universitat Central de Catalunya (UVIC-UCC), Vic, Spain. ²¹Salut Catalunya Central, Hospital de Berga, Berga, Spain. ²²Blood Bank Department—Banc de Sang i Teixits (BST), Barcelona, Spain. ²³IrsiCaixa AIDS Research Institute, Germans Trias i Pujol Research Institute (IGTP), Badalona, Spain. ²⁴Metropolitana Nord Laboratory, Institut Català de la Salut, Badalona, Spain. ²⁵Gerència Territorial Metropolitana Nord, Institut Català de la Salut, Barcelona, Spain. ²⁶Centro de Investigación em Saúde de Manhica (CISM), Maputo, Mozambique. ²⁷Pg. Lluís Companys 23, ICREA, Barcelona, Spain. ²⁸Pediatrics Department, Hospital Sant Joan de Déu, Universitat de Barcelona, Esplugues, Spain. ²⁹Consorcio de Investigación Biomédica en Red de Epidemiología y Salud Pública (CIBERESP), Madrid, Spain. ³⁰Department of Population Health, NYU Grossman School of Medicine, New York, NY, USA. ³¹Department of Haematology, Leiden University Medical Centre, Leiden, The Netherlands. ³²CCTR, Sanquin Blood Supply, Amsterdam, The Netherlands. ³³Lihir Medical Centre—InternationalSOS, Lihir Island, Papua New Guinea. ⁶²These authors contributed equally: Pere Millat-Martinez, Arvind Gharbharan, Andrea Alemany. ⁶³These authors jointly supervised this work: Jaap Jan Zwaginga, Oriol Mitjà, Bart J.A. Rijnders. *Lists of authors and their affiliations appear at the end of the paper. ⁶²email: b.rijnders@erasmusmc.nl

CoV-Early study group

Arvind Gharbharan³⁴, Casper Rokx³⁴, Carlijn Jordans³⁴, Corine Geurtsvankessel³⁴, Grigorios Papageorgiou³⁴, Bart Rijnders³⁴, Peter Katsikis³⁴, Yvonne Müller³⁴, Marion Koopmans³⁴, Susanne Bogers³⁴, Jelle Miedema³⁴, Henk Russcher³⁴, Cees Scherpenisse³⁴, Rene van Engen³⁴, Ayten Karisli³⁴, Hannelore Götz³⁵, Jelle Struik³⁶, Lotte Rokx-Niemantsverdriet³⁶, Nan van Geloven³⁷, Geert Groeneveld³⁷, Jaap Jan Zwaginga³⁷, Lisa Zwaginga³⁷, Josine Oud³⁷, Romy Meier³⁷, Erik van Zwet³⁷, Simon Mooijaart³⁷, Arjan Albersen³⁷, Francis Swaneveld³⁸,

Ellen van der Schoot³⁸, Hans Vrieling³⁸, Leo van de Watering³⁸, Boris Hogema³⁸, Peter van Wijngaarden³⁹, Ronald van Etten³⁹, Adriaan van Gammeren³⁹, Nanda Maas³⁹, Betty van Ginneken³⁹, Jan den Hollander⁴⁰, Jose Verstijnen⁴⁰, Juliette van den Berg – Rahman⁴⁰, Faiz Karim⁴¹, Siepke Hiddema⁴¹, Kim van Elst⁴¹, Elena van Leeuwen-Segarceanu⁴², Annette Reitsma⁴², Karin Molenkamp⁴², Robert Soetekouw⁴³, Caterina Band⁴³, José de Droog⁴³, Jolanda Lammers⁴⁴, Lonneke Buitenhuis⁴⁴, Douwe Postma⁴⁵, David Koster⁴⁵, Michaël Lukens⁴⁵, Thea Scholtens⁴⁵, Maartje van den Boomgaard⁴⁵, Machiel Vonk⁴⁵, Linda Kampschreur⁴⁶, Marit van Vonderen⁴⁶, Loes Vrolijk⁴⁶, Chantal Reusken⁴⁷, Johan Reimerink⁴⁷ & Heli Harvala⁴⁸

³⁴Erasmus MC, Rotterdam, The Netherlands. ³⁵GGD Rotterdam, Rotterdam, The Netherlands. ³⁶Gezondheidscentrum Mathenesserlaan, Rotterdam, The Netherlands. ³⁷Leids Universitair Medisch Centrum, Leiden, The Netherlands. ³⁸Sanquin Blood Supply, Amsterdam, The Netherlands. ³⁹Amphia Hospital, Breda, The Netherlands. ⁴⁰Maasstad ziekenhuis, Rotterdam, The Netherlands. ⁴¹Groene hart hospital, Gouda, The Netherlands. ⁴²St Antonius hospital, Nieuwegein, The Netherlands. ⁴³Spaarne Gasthuis, Hoofddorp, The Netherlands. ⁴⁴Isala Hospital, Zwolle, The Netherlands. ⁴⁵Universitair medisch centrum Groningen, Groningen, The Netherlands. ⁴⁶Medical Center Leeuwarden, Leeuwarden, The Netherlands. ⁴⁷Centrum voor Infectieziektebestrijding, RIVM, Bilthoven, The Netherlands. ⁴⁸NHS Blood and Transplant, London, UK.

CONV-ert study group

Andrea Alemany⁴⁹, Marc Corbacho-Monné⁴⁹, Dan Ouchi⁴⁹, Bonaventura Clotet⁴⁹, Oriol Mitjà⁴⁹, Gèlia Costes⁴⁹, Mar Capdevila-Jáuregui⁴⁹, Pamela Torrano-Soler⁴⁹, Alba San José⁴⁹, Zahida Jiménez⁴⁹, Ferran Ramírez-Viaplana⁴⁹, Susana Ferrer⁴⁹, Mireia Gallardo⁴⁹, Maria Ubals⁴⁹, Camila González-Beiras⁴⁹, Martí Vall-Mayans⁴⁹, Miquel Angel Rodríguez-Arias⁴⁹, Clara Suñer⁴⁹, Jordi Puig⁴⁹, Aroa Nieto⁴⁹, Ivan Galvan-Femenia⁴⁹, Xavier Comas-Leon⁴⁹, Pere Millat-Martínez⁵⁰, Quique Bassat⁵⁰, Bàrbara Baro⁵⁰, Ignacio Blanco⁵¹, Jordi Ara⁵¹, Glòria Bonet Papell⁵¹, Maria Delgado Capel⁵¹, Beatriz Díez Sánchez⁵¹, Maria Pons Barber⁵¹, Cristian Gonzalez Ruiz⁵¹, Laura Navarrete Gonzalez⁵¹, David González García⁵¹, Ainhoa Vivero Larraza⁵¹, Victor Carceles Peiró⁵¹, Clàudia Roquer López⁵¹, Magí Ferrer⁵¹, Pierre Malchair⁵², Sebastian Videla⁵², Vanesa García García⁵², Carlota Gudiol⁵², Aurema Otero⁵², Jose Carlos Ruibal Suarez⁵², Alvaro Zarauza Pellejero⁵², Ferran Llopis Roca⁵², Orlando Rodríguez Cortez⁵², Pablo Casares Gonzalez⁵², Gemma Arcos Vila⁵², Begoña Flores Aguilera⁵², Graciela Rodríguez-Sevilla⁵², Macarena Dastis Arias⁵², Anna Ruiz-Comellas⁵³, Anna Ramírez-Morros⁵³, Judit Roca Font⁵³, Katherine M. Carrasco Matos⁵³, Glòria Saüch Valmaña⁵³, Carla Vidal Obradors⁵³, Joana Rodríguez Codina⁵⁴, Rosa Amado Simon⁵⁴, Silvia Tarres García⁵⁴, Margarida Curriu Sabatès⁵⁴, Raquel Nieto Rodríguez⁵⁴, Joan-Ramon Grifols⁵⁵, Anna Millan⁵⁵, Enric Contreras⁵⁵, Àgueda Ancochea⁵⁵, Rosa Línio⁵⁵, Miriam Fornos⁵⁵, Natàlia Casamitjana⁵⁵, Eva Alonso⁵⁵, Núria Martinez⁵⁵, Laura Analía Maglio⁵⁵, Laura Comellas Fernandez⁵⁵, Nadia Garcia⁵⁵, Luis Hernández⁵⁵, María Isabel González⁵⁵, Anna Bravo⁵⁵, Yolanda García⁵⁵, Núria Prat⁵⁶, Joaquim Verdaguer⁵⁶, Thatiana Vértiz Guidotti⁵⁶, Sergio Benavent⁵⁶, Andrea Sofia Bianco⁵⁶, Ney Nicanor Briones Zambrano⁵⁶, Maria Viozquez Meza⁵⁶, Anna Forcada⁵⁷, Josep Vidal-Alaball⁵⁷, Montserrat Giménez⁵⁸, Alexa París⁵⁸, Gema Fernández Rivas⁵⁸, Cristina Casañ Lopez⁵⁸, Águeda Hernández⁵⁸, Antoni E. Bordoy⁵⁸, Victoria González Soler⁵⁸, Julian Blanco⁵⁹, Edwards Pradenas⁵⁹, Silvia Marfil⁵⁹, Benjamin Trinité⁵⁹, Francini Piccolo Ferreira⁶⁰, Mireia Bonet⁶⁰, Jordi Cantoni⁶⁰ & Michael Marks⁶¹

⁴⁹Fight AIDS and Infectious Diseases Foundation, Badalona, Spain. ⁵⁰Barcelona Institute for Global Health, ISGlobal, Barcelona, Spain. ⁵¹Hospital Universitari Germans Trias i Pujol (HUGTIP), Badalona, Spain. ⁵²Hospital Universitari de Bellvitge, L'Hospitalet de Llobregat, Barcelona, Spain. ⁵³CUAP Manresa, Manresa, Spain. ⁵⁴Hospital Comarcal de Sant Bernabé, Berga, Spain. ⁵⁵Blood Bank Department—Banc de Sang i Teixits (BST), Barcelona, Spain. ⁵⁶Gerència Territorial Metropolitana Nord, Badalona, Spain. ⁵⁷Gerència Territorial Catalunya Central, Sant Fruitós de Bages, Spain. ⁵⁸Metropolitana Nord Laboratory, Badalona, Spain. ⁵⁹IrsiCaixa AIDS Research Institute, Badalona, Spain. ⁶⁰Bioclever-CRO, Barcelona, Spain. ⁶¹London School of Hygiene and Tropical Medicine, London, UK.