

# SCIENTIFIC REPORTS



OPEN

## Improving protein content and quality by over-expressing artificially synthetic fusion proteins with high lysine and threonine constituent in rice plants

Shu-Ye Jiang, Ali Ma, Lifen Xie & Srinivasan Ramachandran

Received: 26 April 2016  
Accepted: 13 September 2016  
Published: 28 September 2016

Rice grains are rich in starch but low in protein with very low level of both lysine and threonine. Thus, it is important to further improve protein quality and quantity, especially to increase lysine and threonine content in rice grains. We artificially synthesized two new genes by fusing endogenous rice genes with lysine (K)/threonine (T) motif (TKTKK) coding sequences. They were designated as *TKTKK1* and *TKTKK2* and their encoded proteins consist of 73.1% and 83.5% of lysine/threonine, respectively. These two genes were under the control of 35S promoter and were independently introduced into the rice genome to generate transgenic plants. Our data showed that overexpression of *TKTKK1* generated stable proteins with expected molecular weight and the transgenic rice seeds significantly increased lysine, threonine, total amino acids and crude protein content by 33.87%, 21.21%, 19.43% and 20.45%, respectively when compared with wild type control; significant improvement was also observed in transgenic rice seeds overexpressing *TKTKK2*. However, limited improvement in protein quality and quantity was observed in transgenic seeds carrying tandom array of these two new genes. Our data provide the basis and alternative strategy on further improving protein quality and quantity in other crops or vegetable plants by synthetic biology.

Rice grains are rich in carbohydrates (nearly 90%) but low in protein (around 10%). Its protein contains low level of lysine and threonine, which are two of nine essential amino acids for humanity and many animals as they are lack of enzymatic machinery for *de novo* synthesis of these amino acids<sup>1</sup>. Thus, increasing lysine and threonine content in rice grains will have a significant social and economic impact.

Various attempts to improve the content of protein and essential amino acids such as lysine and threonine have been carried out. At the early stage, natural or artificial mutants were employed to improve lysine content through traditional breeding. For example, the high lysine maize mutant *opaque2*<sup>2</sup> was used as a parent line to develop the 'quality protein maize' with nearly doubled lysine content in seeds<sup>3</sup>. In rice, higher lysine plants (14%) were regenerated from calli subjected to inhibitory levels of lysine plus threonine<sup>4</sup>.

The second strategy to increase essential amino acids is by modifying biosynthetic and catabolic fluxes<sup>5–8</sup>. This strategy is successful for improving free lysine, threonine and methionine in some plants including tobacco<sup>9,10</sup>, canola<sup>11</sup>, soybean<sup>11</sup> and Arabidopsis<sup>12,13</sup>. However, in maize, the accumulation was mainly observed in embryo but not in endosperm<sup>14–16</sup>. In rice and barley, the expression of the bacterial *DHPS* only slightly increased the content of free lysine<sup>17,18</sup>. To improve the lysine and threonine content in these crops, silencing of *LKR/SDH* by RNA interference (RNAi) was carried out in maize, which dramatically increased lysine content in seeds<sup>14,19</sup>. In rice, free lysine level could be increased up to ~12-fold in leaves and ~60-fold in seeds by over-expression *AK* and *DHPS* and silencing *LKR/SDH* by RNAi<sup>8</sup>.

The third strategy is to generate transgenic plants by over-expressing genes encoding the proteins with higher ratios of essential amino acids. At least 3 lysine-rich genes have been characterized for improving lysine content. Expression of these genes could increase the lysine accumulation up to 10–65% in maize<sup>20–23</sup>. In rice, endogenous

Rice Functional Genomics Group, Temasek Life Sciences Laboratory, 1 Research Link, National University of Singapore 117604, Singapore. Correspondence and requests for materials should be addressed to S.R. (email: sri@tll.org.sg)

genes *RLRH1* and *RLRH2* were characterized, which encode proteins with 14.7% and 20.6% of lysine in their amino acid composition, respectively<sup>24</sup>. They were used for improving lysine content in rice and transgenic plants showed up to 35% increase in its content<sup>24</sup>. Besides these naturally evolved lysine-rich encoding genes, artificially synthetic or modified genes were also used for lysine, methionine, or cysteine enrichment in potato<sup>25</sup>, tobacco<sup>26</sup> or soybean<sup>27</sup>.

In addition to the above mentioned strategies, genetic manipulation of seed storage proteins (SSPs) has also been employed to improve protein quality and quantity in crops<sup>28–30</sup>. Generally, various strategies have been employed to improve protein content and quantity, especially to increase the essential amino acid content. However, reports showed that these strategies were accompanied by various disadvantages<sup>7,24</sup>. Possible healthy risk has been reported on the application of high free lysine transgenic maize<sup>31</sup>. In this study, we have developed the efficient strategy to further improve protein content and quantity, especially for lysine and threonine content in rice leaves and mature seeds.

## Results

### Designing of the synthetic genes by fusing lysine-threonine coding motifs to rice endogenous genes.

To synthesize a high lysine and threonine coding gene, we selected two endogenous rice genes as templates. We first selected the rice gene *LOC\_Os12g16880*, which encodes a seed storage protein with a putative function in impeding the digestion of plant starch and proteins. The lysine-threonine coding motif was designed as 'TKTKKTKKTKKTKKTKKTKKTKKTKKTKKTKKTKKTKR' (T, lysine; K, threonine; R, arginine; S, serine). The gene *LOC\_Os12g16880* was fused with the fragment encoding 16 times of the lysine-threonine motifs (top panel in Fig. 1a). The synthetic gene was designated as *TKTKK1*, which encodes a 78.7 kDa protein, consisting of 48.1% lysine and 25.0% threonine. The synthetic gene was under the control of 35S promoter and the resulted construct was named as *pTKTKK1*. Similar to *pTKTKK1*, the gene *LOC\_Os08g03579* (encoding an unknown protein) was fused with the same lysine-threonine motifs, resulting in another gene designated as *TKTKK2*, which encodes a 68.7 kDa protein, consisting of 55.5% lysine and 28.0% threonine. Accordingly, the resulted construct was named as *pTKTKK2* (middle panel in Fig. 1a). In the third construct *pTKTKK3*, both genes *TKTKK1* and *TKTKK2* were tandemly arrayed under the 35S promoter with opposite orientation (bottom panel in Fig. 1a). We designated these transgenic rice lines generated from these three constructs as *35S::TKTKK1*, *35S::TKTKK2* and *35S::TKTKK3*, respectively.


The gene *LOC\_Os12g16880* was mainly expressed in endosperm (Fig. 1b; Supplementary Table S1). Less expression abundance was observed in ovary followed by embryo (Fig. 1b). Very low expression level was detected in the remaining tissues. Thus, we selected the endosperm-preferred gene so that its protein might show more stable storage in seeds as usual. On the contrary, the second gene *LOC\_Os08g03579* was totally different from the first one in their expression patterns with very high expression abundance in all tested tissues (Fig. 1c; Supplementary Table S1). As this gene was also highly expressed in endosperm, its protein was supposed to be naturally stored in seeds.

### Molecular characterization of transgenic plants carrying different constructs.

We have generated and planted a total of 40 T0 transgenic lines from 60 independent hygromycin resistant calli for the construct *pTKTKK1*. DNA samples extracted from 18 T0 lines were submitted to Southern blot hybridization (Fig. 2a). The blotting data showed that a total of 11 lines might contain single copy of T-DNA insertion as indicated by red triangle. They were lines 2, 5, 9, 14, 28, 21, 12, 22, 20, 11, and 31. The quantitative real-time reverse transcription PCR (qRT-PCR) analysis showed that all of the analyzed lines exhibited at least 1.5-fold higher expression level when compared with wild type (WT) plant (Fig. 2b; Supplementary Table S2). The line 23 showed the highest expression abundance (664.9-fold) but contained three copies of T-DNA insertion (Fig. 2a,b). Finally, we selected three independent lines 9, 14 and 21 for further investigation. These lines showed relatively high expression level for the gene *TKTKK1* and contained single copy of T-DNA insertion.

For the construct *pTKTKK2*, a total of 60 T0 lines were regenerated from 80 independent transgenic calli. Based on the qRT-PCR analysis (Fig. 3a,b; Supplementary Table S3), the *TKTKK2* gene showed higher level of expression in all transgenic lines when compared with its endogenous gene *LOC\_Os08g03579*. However, the relative expression abundance was significantly lower than those lines from *pTKTKK1* (Fig. 2b). This should be due to that the expression level of calibrator gene *LOC\_Os08g03579* in WT leaves is much higher than that of the gene *LOC\_Os12g16880* (Fig. 1b,c). We selected the top 12 lines based on their expression abundance for Southern blot hybridization (Fig. 3c). The analysis showed that most of the lines have 2–4 copies of T-DNA insertion and only two lines (21 and 46) contained single copy of T-DNA integration as indicated by red stars (Fig. 3c), which were selected for further investigation.

For the construct *pTKTKK3*, we have also generated and planted a total of 60 T0 lines regenerated from 85 independent transgenic calli. As two TKTKK-motif coding genes were overexpressed in these transgenic plants, we measured the expression level of these two genes including *TKTKK1* (Fig. 4a; Supplementary Table S4) and *TKTKK2* (Fig. 4b; Supplementary Table S4). Among the 60 transgenic lines, expression data from 5 of these lines were not qualified for further analysis, thus, only the expression data from the remaining 55 lines were presented in this study. For the *TKTKK1* gene, some of the 55 analyzed lines showed the similar expression level to the endogenous gene *LOC\_Os12g16880* (1.29-fold for Line 27) and others showed higher (up to 54.45-fold for Line 35) expression level. Similar results were observed for the *TKTKK2* gene (Fig. 4b). Generally, expression level of either *TKTKK1* or *TKTKK2* in *35S::TKTKK3* transgenic plants was lower than that in *35S::TKTKK1* or *35S::TKTKK2* plants. For example, for the *35S::TKTKK1* plants, many lines showed more than 100-fold higher expression than the endogenous gene *LOC\_Os12g16880*, significantly higher than that in the *35S::TKTKK3* plants




**Figure 1. Construction of binary vectors by fusing endogenous rice genes with TKTKK coding motifs.** (a) Construction of three binary vectors for improving lysine/threonine content. Top and middle panels showed the vectors *pTKTKK1* and *pTKTKK2*, which were constructed by fusing a rice gene *LOC\_Os12g16880* or *LOC\_Os08g03579* with 16-fold TKTKK coding motifs TKTKKTKTKKTKTKKTKTKKTKTKKTKTKTRS, respectively. Bottom panel showed the binary vector *pTKTKK3*, which was constructed by tandem arraying both *TKTKK1* and *TKTKK2* in inverse order. (b,c) show the expression patterns of both genes *LOC\_Os12g16880* and *LOC\_Os08g03579*, respectively. Expression data were achieved from NCBI GEO dataset with accession number GSE21396. Processed signal intensity, which was converted from processed raw data, was used to estimate the expression abundance in each gene among 11 tissues from different developmental stages. The prefix “*LOC\_*” in each gene locus name was omitted for convenience. Tissue names in (b,c) were labelled in each column.

(Figs 2b and 4a,b). The data suggested the slight co-suppression between *TKTKK1* and *TKTKK2* when they were both overexpressed under the control of 35S promoter.

Based on the expression data of *TKTKK1* and *TKTKK2*, we have selected top 17 lines for T-DNA copy number detection by Southern blot hybridization (Fig. 4c). Most of transgenic lines with relatively higher expression level for these two genes contained two or more copies of T-DNA insertion. We have detected a total of 6 lines with single copy of T-DNA insertion and these lines showed similar expression level for both genes *TKTKK1* and *TKTKK2*.

### Synthetic fusion proteins were stably expressed in the 35S::TKTKK1 transgenic plants.

As the synthetic genes encode proteins with 16 times of repeated motifs “TKTKKTKTKKTKTKKTKTKKTKTKKTKTKTRS”; one might argue whether they could be translated and be stable in rice seeds. We first tested the protein stability in *E. coli* cells. The synthetic gene *TKTKK1* was fused with *GLUTATHIONE S-TRANSFERASE*




**Figure 2. Molecular characterization of 35S::TKTKK1 transgenic plants.** (a) Copy number detection of T-DNA insertion in transgenic plants by Southern blot hybridization. DNA samples from a total of 18 transgenic plants were restricted by *EcoRV* and then transferred onto nylon membrane for hybridization using the *HPT* probe. The red triangle indicated the lines with single copy of T-DNA insertion. (b) Bar diagrams showing expression patterns of the *TKTKK1* gene in the 18 independent transgenic plants by qRT-PCR. The mRNA relative amount (Y axis) was calculated according to the description in Materials and methods. The amplification of an *eEF-1a* gene was used as an internal control to normalize the data. The red arrows in (a) indicated the lines with single copy of T-DNA insertion by Southern blotting analysis. The red stars “\*” in (a, b) indicated the lines with single copy of T-DNA insertion and with relatively higher expression level. These three independent lines were selected for further investigation.

(GST) by sub-cloning into the *pGEX-6P-1* vector. The resulted plasmid was used for *E. coli* transformation followed by protein extraction and western blot hybridization. The blotting result using GST antibody showed that the fusion protein could be stably expressed in the *E. coli* cells with expected molecular weight (78.7 kDa for TKTKK1 and 28 kDa for GST, Fig. 4d). Furthermore, we also detected the protein stability in transgenic rice seeds by Western blot hybridization. The result showed that the synthetic fusion proteins could be detected in all 35S::TKTKK1 transgenic rice seeds (Fig. 4e). Although mRNA transcript signal could be detected in the 35S::TKTKK2 or 35S::TKTKK3 transgenic plants, very faint signal was detected in 35S::TKTKK2 and no protein signal was detected in 35S::TKTKK3 transgenic seeds (Fig. 4e). The data suggested that the stable expression of synthetic proteins might be dependent on the fused endogenous genes and used constructs.

**Preliminary phenotyping and genotyping in T1 generation of transgenic plants.** As expression level varied with a large range for the 35S::TKTKK1 transgenic lines, we selected two lines (9 and 23) for testing lysine and protein content. The line 9 showed the middle level of expression abundance with single copy of T-DNA insertion while the line 23 showed the highest expression level with three copies of T-DNA insertion (Fig. 2). Both lines contained significantly higher level of lysine content in two-month-old vegetative stage of plants (Fig. 5a; Supplementary Table S5). The lysine content was increased by 46.2% for line 9 and by 38.5% for line 23. These two lines showed no difference in lysine content by statistical analysis although they exhibited significant difference in their expression level. The crude protein content was enhanced by 27.1% and 25.4% for lines 9 and 23, respectively when compared with the WT plants (Fig. 5b; Supplementary Table S5). Thus, our preliminary data showed that one of our synthetic genes should play a role in improving lysine and protein content.

Thermal asymmetric interlaced PCR (TAIL-PCR)<sup>32</sup> was employed to amplify T-DNA flanking sequence tags (FSTs) for single copy of T-DNA insertion lines. A total of 5 independent lines were subjected to FST analysis (Fig. 5c). For the 35S::TKTKK1 transgenic line 9, T-DNA was inserted into the 8,255,730<sup>th</sup> bp of chromosome 10. The T-DNA was inserted into the first exon of the gene *LOC\_Os10g16560*, which was annotated to encode a retrotransposon. For line 14, T-DNA was inserted into the 6,303,477<sup>th</sup> bp of chromosome 1 and no gene was tagged. For line 21, T-DNA was inserted into the 1,238,504<sup>th</sup> bp of chromosome 12 and no annotated gene was tagged. In the 35S::TKTKK2 line 46, T-DNA was inserted into the 11204023<sup>rd</sup> bp of chromosome 10 and no gene was




**Figure 3. Molecular characterization of 35S::TKTKK2 transgenic plants.** (a,b) show the expression patterns of the *TKTKK2* gene in 60 transgenic plants by qRT-PCR. The mRNA relative amount (Y axis) was calculated as shown in Materials and methods. The *eEF-1a* gene was used as an internal control to normalize the data as shown in Fig. 2b. (c) Copy number detection of T-DNA insertion in transgenic plants by Southern blot hybridization. DNA samples from top 12 transgenic plants in expression were restricted by *EcoRV* and then transferred into nylon membrane for hybridization using the *HPT* probe. The red star “\*” in (b,c) indicated the line with high level of expression signal and with single copy of T-DNA insertion, which was selected for further analysis.


tagged by the T-DNA insertion. In the line 5 carrying the construct *pTKTKK3*, the T-DNA was inserted into the 2690708<sup>th</sup> bp of chromosome 12. Similarly, no gene was tagged as the T-DNA was inserted into the non-coding region. Thus, in these lines, T-DNA was inserted into either non-coding region or a retrotransposon region, which minimized the phenotypic variation from T-DNA mutagenesis.

Based on T-DNA tagging positions in the rice genome, we designed three primer sets to identify the genotypes (homozygote, heterozygote and WT) at T-DNA insertion locus for each line as explained in the figure (Fig. 5d,e). Based on the PCR analysis, the T-DNA locus in each line was segregated at the ratio 3:1 by  $\chi^2$  test at  $p < 0.01$  for all 5 population from 5 independent lines (Fig. 5f). Thus, seeds were harvested from homozygous transgenic plants for further phenotype investigation. All DNA samples were also subjected to another set of PCR using primers designed from the selection marker gene *HPT* (encoding hygromycin phosphotransferase). The experiment showed that the marker gene was detected in all heterozygotes and homozygotes and no signal was detected in the WT genotype. The data further confirmed that only single copy of T-DNA was integrated into the rice genome in each line.

**Fusing *TKTKK* coding motifs to a rice endogenous gene significantly increased lysine, threonine and crude protein content in rice seeds.** Seeds harvested from homozygous T2 transgenic plants in each of 5 lines were subjected to measuring protein content and quality. We have tested lysine, threonine, total amino acids and crude protein content in 3 independent 35S::TKTKK1 lines. For 35S::TKTKK2 lines, only two




**Figure 4. Molecular characterization of 35S::TKTKK3 transgenic plants and western blot hybridization.** (a,b) show the bar diagrams of expression patterns of both genes *TKTKK1* and *TKTKK2* in 55 35S::TKTKK3 transgenic plants by qRT-PCR, respectively. (c) Southern blot hybridization for detecting copy numbers of T-DNA insertion. DNA samples (line numbers were labelled on the top of the panel) were digested by the restriction enzyme *EcoRV* and were then transferred into nylon member for hybridization with the *HPT* probe. The red stars “\*” indicated the lines with single copy number of T-DNA insertion. (d,e) Western blot hybridization using crude proteins from *E. coli* (d) and T2 rice grains (e), respectively. The “M” in (d,e) indicates the protein marker used for the hybridization. The numbers in (d) indicate the crude proteins extracted from *E. coli* carrying the plasmid *pGEX-6P-1* (1 and 2) and *pGEX-6P-1::TKTKK1* (3 and 4). The white arrows indicated the GST expression in the *E. coli* carrying the plasmid *pGEX-6P-1*. The green arrows indicated the GST fusion protein expression in the *E. coli* carrying *pGEX-6P-1::TKTKK1*. The numbers in (e) indicate the different transgenic lines used for crude proteins extraction from rice seeds. Three lines (1, line 9, 2, line 14 and 3, line 21) were from 35S::TKTKK1. Line 46 (indicated by 4) and Line 5 (indicated by 5) were from 35S::TKTKK2 and 35S::TKTKK3, respectively. The number 6 indicates WT and 7 indicates negative control. The green arrows indicated the stable expression of TKTKK1 protein in lines 9, 14 and 21 by Western blot hybridization.



**Figure 5. Preliminary analysis of lysine and crude protein and FST analysis.** (a,b) Lysine and crude protein content, respectively, in T0 generation of transgenic plants carrying the construct 35S::TKTKK1. (c) Characterization of FSTs of T-DNA insertion. A total of 5 independent lines from 35S::TKTKK1, 35S::TKTKK2 or 35S::TKTKK3 were selected for FST analysis. NA, not available. (d) Schematic diagram for PCR genotyping T1 or T2 generation of transgenic plants. Forward (F) and reverse (R) primers were designed according to the FSTs. Left border reverse (LBR) and right border forward (RBF) primers were designed according to T-DNA border sequence. (e) Identification of three different genotypes including heterozygotes, homozygotes and WT in T1 or T2 transgenic plants based on PCR results. The symbol “+” indicated that PCR fragment could be amplified using corresponding primer sets and “-” indicated no PCR product. (f) Genotyping of five populations from 5 T1 generations of transgenic plants and their  $\chi^2$  test.

independent lines with single copy of T-DNA insertion was generated and we selected line 46 for further analysis as the expression level of *TKTKK2* in this line is lower than another one (Fig. 3). For the 35S::TKTKK3 lines, a



**Figure 6. Expression analysis of synthetic genes and measurement of amino acid and crude protein contents in transgenic rice seeds.** (a) The qRT-PCR analysis of synthetic genes in transgenic seeds from 3 constructs. (b–e) shows the content (percentage) of lysine (b), threonine (c), total amino acids (d) and crude protein (e) in 5 independent transgenic plants. Matured seeds were dried at 37°C for 3 days and were then subjected to amino acid and protein measurement. Asterisks “\*” and “\*\*” indicate significant differences at  $P < 0.05$  and  $P < 0.01$ , respectively.

total of 8 independent lines have single copy of T-DNA insertion and they showed similar expression level for both *TKTKK1* and *TKTKK2* (Fig. 4) and we randomly selected line 5 for further investigation.

We first analysed the expression abundance of targeted genes in transgenic seeds in these 5 independent lines (Fig. 6a; Supplementary Table S6). The qRT-PCR data showed that four lines including 9, 14 and 21 from *35S::TKTKK1* and 46 from *35S::TKTKK2* exhibited at least 3-fold higher expression level when compared with corresponding endogenous genes (Fig. 6a). However, low expression level was detected for the line 5 from *35S::TKTKK3*. We then further investigated the protein and lysine/threonine content in these lines. Our data showed that lysine content was increased by 16.13–33.87% in 3 independent *35S::TKTKK1* lines when compared with WT, statistically higher than WT (Fig. 6b; Supplementary Table S6). For the *35S::TKTKK2* line 46, 12.90% increase was detected with statistically higher than WT (Fig. 6b). However, for the *35S::TKTKK3* line 5, only 6.45% more lysine content was detected with no statistical difference when compared with WT (Fig. 6b). On the other hand, threonine content was improved by 12.12–21.21% in 3 *35S::TKTKK1* lines when compared with WT (Fig. 6c; Supplementary Table S6). For the *35S::TKTKK2* line 46, threonine was enhanced by 13.63%, significantly

higher than WT (Fig. 6c). In the 35S::TKTKK3 line 5, only 9.09% increase in threonine content was measured, with no statistical difference when compared with WT (Fig. 6c). For total amino acid analyses, three lines from 35S::TKTKK1 showed 17.16–19.43% increase, significantly higher than WT (Fig. 6d; Supplementary Table S6). For the 35S::TKTKK2 line 46, its total amino acid content showed up to 14.05% increase and is statistically higher than WT. However, the 35S::TKTKK3 line 5 showed no significant difference although its content was increased by 9.38% (Fig. 6d). We then compared the crude protein content of all these 5 independent transgenic seeds with that in WT seeds (Fig. 6e; Supplementary Table S6). Similarly, in all three 35S::TKTKK1 lines, protein content was increased by 16.03–20.45%. The 35S::TKTKK2 line 46 showed significant increase by 12.09% and another line from 35S::TKTKK3 showed no significant difference. In general, both 35S::TKTKK1 and 35S::TKTKK2 significantly increased the content of lysine, threonine, total amino acids and crude protein while 35S::TKTKK3 showed limited improvement in protein content and quality.

### Overexpression of TKTKK1 or TKTKK2 might not affect normal growth and development in rice.


Generally, the line 46 from 35S::TKTKK2 exhibited shorter height when compared with the remaining plants (Fig. 7a). At the mature stage, normally filled grains (full seeds) were less in lines 9 and 14 (Fig. 7b). Detail measurement showed that all three transgenic plants from 35S::TKTKK1 showed similar height to WT plants (Fig. 7c; Supplementary Table S7). Similar result was observed in 35S::TKTKK3 (Fig. 7c). However, the line 46 from 35S::TKTKK2 showed shorter height when compared with WT plants (Fig. 7c). We calculated till number per plant in both WT and transgenic plants and found that no significant difference was observed when compared with WT plants (Fig. 7d; Supplementary Table S7). We then surveyed the average seeding rate and showed that both line 21 from 35S::TKTKK1 and line 5 from 35S::TKTKK3 exhibited similar seeding rate to that in WT plants (Fig. 7e; Supplementary Table S7). However, the remaining 3 plants from either 35S::TKTKK1 or 35S::TKTKK2 showed significant lower seeding rate when compared with WT plants (Fig. 7e). Finally, we measured the average grain yield per plant and found that only the line 9 showed lower grain yield per plant (Fig. 7f; Supplementary Table S7). The remaining 4 transgenic lines showed no significant difference when compared with WT plants (Fig. 7f). In this study we have generated at least one transgenic plant (line 21), which showed normal growth and development but improved protein quality and quantity. All in all, although some transgenic plants showed shorter plant height, lower seeding rate or grain yield, these changed phenotypes were not related to the overexpression of any of synthetic fusion genes (Fig. 7).

### Discussion

Generally, content of essential amino acids in an organism could be improved by increasing either protein-bound or free amino acids. Although free essential amino acids could be increased by around 60-fold in rice seeds<sup>8</sup>, free amino acids pool is small compared to the protein-bound amino acids<sup>33</sup> and thus, limiting the net accumulation of essential amino acids. Additionally, free lysine is known to react on heating with sugars to form chemical compounds called N $\epsilon$ -(Carboxymethyl) lysine, which is one of well-characterized advanced glycation end products (AGEs)<sup>31</sup>. AGEs are a diverse group of highly oxidant compounds that are linked to numerous diseases, including diabetes, Alzheimer's disease and cancers<sup>34,35</sup>. The transgenic maize variety LY038 with 50-fold higher levels of free lysine in the maize kernel by expressing the bacterial *DHPS* has been approved for animal food use in Japan, S. Korea, Canada, Australia, New Zealand and the US (<https://www.isaaa.org/gmaprovaldatabase/event/default.asp?EventID=146>). However, Monsanto has withdrawn its application due to its potential food safety (<http://www.independentsciencenews.org/news/transgenic-corn-ly038-withdrawn/>). All these data imply the importance for us to further improve the protein-bound essential amino acids. However, currently employed strategies to improve protein-bound essential amino acids have various disadvantages such as protein instability for synthetic genes, protein allergy in seeds, and other accompanied agronomic traits including low seeding set, low yield etc. In this study, we used the rice endogenous genes as templates to design new genes by fusing *TKTKK* coding motifs to further improve both lysine and threonine content. Our data showed that the expressed proteins could stably exist in either *E. coli* cells or transgenic rice seeds (Fig. 4d,e). Expression of such fusion proteins has limited effect on other agronomic traits and our data showed no relationship between changed agronomic traits and T-DNA integration (Fig. 7). Thus, our study provides an alternative strategy to further improve protein-bound essential amino acids.

Artificially modified genes have been used to improve protein quality as such a strategy is a straightforward molecular improvement of amino acid constituent<sup>36</sup>. Many genes have been modified including these genes encoding  $\alpha$ -zein,  $\gamma$ -zein,  $\beta$ -phaseolin, 2S albumin, Braizil nut 2S etc<sup>36,37</sup>. However, major challenge to this strategy is the instability of modified proteins<sup>36,37</sup>. In this study, the original protein sequences were not modified and were fused with lysine- and threonine-rich motifs. Thus, the employed strategy might minimize the change in protein stability. On the other hand, some of seed proteins are responsible for allergy. We submitted both genes *Os12g16880* and *Os08g03579* as well as TKTKK motif sequences for BLAST searches against the allergy database (<http://www.allergenonline.org/index.shtml>). The results showed that no sequence homology was found in the database. We have also used AllerHunter (<http://tiger.dbs.nus.edu.sg/AllerHunter/running.html>) and Allerdicator (<http://allerdicator.vbi.vt.edu/predict/>) for allergen prediction of TKTKK1 and TKTKK2. No potential allergen was predicted. Thus, both employed rice genes and the designed TKTKK motifs could be used to synthesize high lysine/threonine coding genes without potential risk for the production of allergy proteins.

The application of synthetic biology on improving protein quality has been carried out long time ago. For example, several genes have been designed according to an alpha-helical coiled-coil structure and these genes encoded high lysine proteins<sup>26</sup>. As a result, lysine content in seeds was increased in these transgenic tobacco seeds carrying one of these genes. However, lysine content was increased by less than 20%<sup>26</sup>, which might be due to that these genes encoded high lysine proteins with very small molecular weight (3–7 kDa). In this study, lysine/threonine-rich fragments consist of 73% and 83% of total amino acids with molecular weight at 78.7 kDa



**Figure 7. Investigation on agronomical traits in transgenic plants.** (a) Phenotypic characterization of rice plants at the grain filling stage in WT and 5 independent transgenic lines. (b) Panicles at the mature stage in WT and 5 independent transgenic lines. (c–f) shows average plant height (cm), average tiller numbers per plant, average seeding rate, and average grain yield per plant, respectively, in WT and 5 independent transgenic lines. Only the productive tillers that could produce spikes and seeds were scored in this investigation. The seeding rate was determined using the ratio of normally filled seeds among total panicle grains. In (a–f), lines 9, 14 and 21 carried the *35S::TKTKK1* construct; line 46 and line 5 were from the constructs *35S::TKTKK2* and *35S::TKTKK3*, respectively.

and 68.7 kDa in both *TKTKK1* and *TKTKK2* genes, respectively. These synthetic genes encode proteins with similar structure, which forms alpha-helices followed by coiled-coil tails (Supplementary Figure S1). Although the predicted structures are similar for these two proteins, *TKTKK1* showed more stability and contributed more efficiently to the improvement of protein quantity (Fig. 6). Thus, more experiments should be carried out to

test which endogenous genes should be employed to form TKTKK fusion proteins to synthesize stable proteins. Currently, we fused 16 times of TKTKKTKTKKTKTKKTKTKKTKTKKTKTKKTR motifs to an endogenous rice protein. We may also need to figure out the optimized motif numbers to achieve the highest level of protein improvement.

Evidence has shown that over-expression of high lysine coding genes could significantly increase lysine content in transgenic plants<sup>21,23,38,39</sup>. Similarly, over-expression of genes encoding proteins with higher percentage of threonine or other amino acids could also increase the level of threonine or other amino acids<sup>7,40–42</sup>. However, in plants, limited genes are available that encode high lysine or threonine proteins. One of widely employed high lysine coding genes is *SB401* from maize, which encodes a protein consisting of 16.7% of lysine<sup>38</sup>. In rice, our genome-wide survey showed that only 11 genes encoded proteins (with >150 amino acid long) containing more than 20% lysine. However, these protein sequences consist of only 0.72–7.89% threonine. On the other hand, we have detected only 8 genes encoding proteins (no less than 150 amino acids) with more than 15% threonine. No gene encodes a protein with more than 15% lysine and 15% threonine in the rice genome. Thus, high lysine and/or threonine genes should be artificially modified or synthesized so that the lysine /threonine content could occupy higher percentage. However, even in artificially modified genes, the percentage of lysine or threonine among the total amino acids was still not high due to the instability of the modified proteins. For example, in the synthetic gene *CP 3-5*, it encodes only 31% lysine and 20% methionine<sup>26</sup>. In this study, we increased lysine and threonine percentages to 48.1% and 25.0%, respectively, by fusing a TKTKK coding motifs to endogenous rice genes. The synthetic genes could be stably expressed in the rice genome and generated stable proteins in rice seeds as detected by Western blot hybridization (Fig. 4e). As a result, we successfully generated high lysine plants and seeds. We have over-expressed two genes encoding proteins with molecular weight at 78.7 and 68.7 kDa, respectively. More experiments should be carried out to demonstrate whether the molecular weight or motif length of synthetic proteins might affect the protein stability or lysine/threonine content in transgenic plants or seeds.

Although the efficient methods to increase free lysine or threonine through metabolic pathway have been reported, the strategies might not be commercially used to improve essential amino acid content due to (1) low total free amino acid content compared with protein-bound amino acids and (2) possible healthy risk from free amino acids. Thus, further improvement of protein-bound essential amino acids has been put in the first choice. Our study showed that artificially fusing TKTKK coding sequences to an endogenous rice gene and then over-expressing it in the rice genome could significantly increase both crude protein and essential amino acid content. Our research might provide an alternative way to improve protein content and quantity. Our data showed that employed rice genes might also affected the accumulation of TKTKK-fused protein. Therefore, higher accumulation of the fused protein through this strategy should be achieved by selecting and optimizing the endogenous rice genes which were used to fuse with TKTKK motif coding sequences. Currently, we have tested only two rice genes *LOC\_Os08g03579* and *LOC\_12g16880*. These two genes showed very high expression level in the rice plants (Fig. 1). As a result, this might affect the expression level of *TKTKK1* or *TKTKK2* and subsequent accumulation of their fused proteins. Therefore, selection of seed-specific genes with low expression level should be an alternative way to further improve lysine and threonine content through fusing with TKTKK coding sequences.

Increase of free lysine content has been achieved by transgenic expression of bacterial lysine feedback-insensitive *DHPS* genes and this method has been used for many species as shown in the Introduction. However, some limitations were found in maize<sup>14–16</sup>, rice and barley<sup>17,18</sup>. Thus, it is still not a universal method. On the other hand, modified high lysine genes were used for lysine improvement in limited species. For example, a modified gene encoding a lysine-rich zein was expressed normally in maize<sup>39</sup>. However, the modified protein was abnormally localized on cell wall instead of endoplasmic reticulum<sup>43</sup>. Current data showed that no universal method can be used for improving amino acid content. We have developed an efficient strategy to improve lysine, threonine and crude protein content in rice. Two genes have been used to fuse with TKTKK coding motifs for improving protein content and quality. However, the seed-specific gene *LOC\_Os12g16880* showed higher efficiency. Thus, in order for us to apply this method to other species, seed-specific endogenous genes in targeted species might be selected to fuse with the TKTKK coding motifs for transgenic expression. As we used endogenous genes for fusion expression and their proteins should be more stable, thus, providing a universal strategy for improving lysine, threonine and crude protein content in any crop species and even in vegetable plants.

## Methods

**Plant materials and growth conditions.** Rice variety Nipponbare (*Oryza sativa*) was used for all experiments. Rice seeds were germinated and were then transferred into soil pots. Plants were grown in greenhouse under natural sunlight and temperature conditions. The mature seeds were also used to induce calli for *Agrobacterium*-mediated genetic transformation.

**Selection of candidate endogenous genes and expression analysis of both *LOC\_Os12g16880* and *LOC\_Os08g03579*.** The selection standard of endogenous genes for fusing expression with TKTKK motifs is based on gene size, expression and their putative functions. We selected these genes encoding proteins with 100–200 amino acids long so that the fused proteins are no more than 100 kDa in molecular weight. We have identified 12,978 annotated genes encoding such length of proteins based on the Rice Genome Annotation Project database. We surveyed the expression profiling of these genes based on the microarray dataset with the NCBI GEO accession number GSE21396. Finally, we selected one gene *LOC\_Os12g16880*, which showed seed-specific expression and encodes a seed storage protein. Another gene *LOC\_Os08g03579* showed high expression abundance in multiple tissues including seeds and encodes an expressed protein with unknown function, which might reduce the negative effect of this gene on plant growth.



**Investigation of morphological traits of transgenic plants.** Besides the measurement of amino acids and crude protein in vegetative stage of plants and mature grains, we have also investigated the morphological traits of transgenic plants by comparing with WT plants. Plant height, tiller number, seeding rate and grain yield were evaluated according to the standard evaluation system for rice (IRRI, 2002). A total of 5 independent homozygous T3 lines were used for the trait investigation. Around 40 individuals in each line were planted in each replicate for all of the survey of morphological traits. Three biological replicates were carried out and the difference was tested by statistical analysis.

## References

- Galili, G. *et al.* Production of transgenic plants containing elevated levels of lysine and threonine. *Biochem. Soc. Trans.* **22**, 921–925 (1994).
- Mertz, E. T., Bates, L. S. & Nelson, O. E. Mutant gene that changes protein composition and increases lysine content of maize endosperm. *Science*. **145**, 279–280 (1964).
- Gibbon, B. C. & Larkins, B. A. Molecular genetic approaches to developing quality protein maize. *Trends Genet.* **21**, 227–233 (2005).
- Schaeffer, G. W. & Sharpe, F. T. Increased lysine and seed storage protein in rice plants recovered from calli selected with inhibitory levels of lysine plus threonine and S-(2-aminoethyl) cysteine. *Plant Physiol.* **84**, 509–515 (1987).
- Galili, G., Tang, G., Zhu, X. & Gakiere, B. Lysine catabolism: a stress and development super-regulated metabolic pathway. *Curr. Opin. Plant Biol.* **4**, 261–266 (2001).
- Galili, G. New insights into the regulation and functional significance of lysine metabolism in plants. *Annu. Rev. Plant Biol.* **53**, 27–43 (2002).
- Galili, G. & Amir, R. Fortifying plants with the essential amino acids lysine and methionine to improve nutritional quality. *Plant Biotechnol. J.* **11**, 211–222 (2013).
- Long, X., Liu, Q., Chan, M., Wang, Q. & Sun, S. S. Metabolic engineering and profiling of rice with increased lysine. *Plant Biotechnol. J.* **11**, 490–501 (2013).
- Shaul, O. & Galili, G. Increased lysine synthesis in transgenic tobacco plants expressing a bacterial dihydrodipicolinate synthase in their chloroplasts. *Plant J.* **2**, 203–209 (1992).
- Shaul, O. & Galili, G. Concerted regulation of lysine and threonine synthesis in tobacco plants expressing bacterial feedback-insensitive aspartate kinase and dihydrodipicolinate synthase. *Plant Mol. Biol.* **23**, 759–768 (1993).
- Falco, S. C. *et al.* Transgenic canola and soybean seeds with increased lysine. *Biotechnology*, **13**, 577–582 (1995).
- Ben-Tzvi Tzchori, I., Perl, A. & Galili, G. Lysine and threonine metabolism are subject to complex patterns of regulation in Arabidopsis. *Plant Mol. Biol.* **32**, 727–734 (1996).
- Zhu, X. & Galili, G. Increased lysine synthesis coupled with a knockout of its catabolism synergistically boosts lysine content and also transregulates the metabolism of other amino acids in Arabidopsis seeds. *Plant Cell.* **15**, 845–853 (2003).
- Frizzi, A. *et al.* Modifying lysine biosynthesis and catabolism in corn with a single bifunctional expression/silencing transgene cassette. *Plant Biotechnol. J.* **6**, 13–21 (2008).
- Huang, S. *et al.* High-lysine corn produced by the combination of enhanced lysine biosynthesis and reduced zein accumulation. *Plant Biotechnol. J.* **3**, 555–569 (2005).
- Mazur, B., Krebbers, E. & Tingey, S. Gene discovery and product development for grain quality traits. *Science*. **285**, 372–375 (1999).
- Lee, S. I. *et al.* Constitutive and seed-specific expression of a maize lysine-feedbackinsensitive dihydrodipicolinate synthase gene leads to increased free lysine levels in rice seeds. *Mol. Breeding*. **8**, 75–84 (2001).
- Brinch-Pedersen, H., Galili, G., Knudsen, S. & Holm, P. B. Engineering of the aspartate family biosynthetic pathway in barley (*Hordeum vulgare* L.) by transformation with heterologous genes encoding feed-back-insensitive aspartate kinase and dihydrodipicolinate synthase. *Plant Mol. Biol.* **32**, 611–620 (1996).
- Houmard, N. M. *et al.* High-lysine corn generated by endosperm-specific suppression of lysine catabolism using RNAi. *Plant Biotechnol. J.* **5**, 605–614 (2007).
- Lang, Z., Zhao, Q., Yu, J., Zhu, D. & Ao, G. Cloning of potato *SBgLR* gene and its intron splicing in transgenic maize. *Plant Sci.* **166**, 1227–1233 (2004).
- Yu, J. *et al.* Seed-specific expression of the lysine-rich protein gene *sb401* significantly increases both lysine and total protein content in maize seeds. *Food Nutr. Bull.* **26**, 427–431 (2005).
- Wang, M. *et al.* Improved nutritive quality and salt resistance in transgenic maize by simultaneously overexpression of a natural lysine-rich protein gene, *SBgLR*, and an ERF transcription factor gene, *TSRF1*. *Int. J. Mol. Sci.* **14**, 9459–9474 (2013).
- Yue, J., Li, C., Zhao, Q., Zhu, D. & Yu, J. Seed-specific expression of a lysine-rich protein gene, *GhLRP*, from cotton significantly increases the lysine content in maize seeds. *Int. J. Mol. Sci.* **15**, 5350–5365 (2014).
- Wong, H. W., Liu, Q. & Sun, S. S. Biofortification of rice with lysine using endogenous histones. *Plant Mol. Biol.* **87**, 235–248 (2015).
- Yang, M. S. *et al.* Expression of a synthetic gene for improved protein quality in transformed potato plants. *Plant Sci.* **64**, 99–111 (1989).
- Keeler, S. J. *et al.* Expression of de novo high-lysine alpha-helical coiled-coil proteins may significantly increase the accumulated levels of lysine in mature seeds of transgenic tobacco plants. *Plant Mol. Biol.* **34**, 15–29 (1997).
- Zhang, Y. *et al.* Improved protein quality in transgenic soybean expressing a *de novo* synthetic protein, MB-16. *Transgenic Res.* **23**, 455–467 (2014).
- Coleman, C. E. *et al.* Expression of a mutant alpha-zein creates the floury2 phenotype in transgenic maize. *Proc. Natl. Acad. Sci. USA* **94**, 7094–7097 (1997).
- Kim, H. J., Lee, J. Y., Yoon, U. H., Lim, S. H. & Kim, Y. M. Effects of reduced prolamins on seed storage protein composition and the nutritional quality of rice. *Int. J. Mol. Sci.* **14**, 17073–17084 (2013).
- Kawakatsu, T., Hirose, S., Yasuda, H. & Takaiwa, F. Reducing rice seed storage protein accumulation leads to changes in nutrient quality and storage organelle formation. *Plant Physiol.* **154**, 1842–1854 (2010).
- Fu, M. X. *et al.* The advanced glycation end product, Nεpsilon-(carboxymethyl) lysine, is a product of both lipid peroxidation and glycoxidation reactions. *J. Biol. Chem.* **271**, 9982–9986 (1996).
- Liu, Y. G. & Whittier, R. F. Thermal asymmetric interlaced PCR: automatable amplification and sequencing of insert end fragments from P1 and YAC clones for chromosome walking. *Genomics*. **25**, 674–681 (1995).
- Amir, R. & Tabe, L. Molecular approaches to improving plant methionine content. In Pawan, K. J., Jaiwal, K. & Rana, P. S. eds, *plant genetic engineering: metabolic engineering and molecular farming ii*. Vol 8. Studium Press LLC, Houston, pp 1–26 (2006).
- Vistoli, G. *et al.* Advanced glycoxidation and lipoxidation end products (AGEs and ALEs): an overview of their mechanisms of formation. *Free Radic Res* **47**, 3–27 (2013).
- Turner, D. P. Advanced glycation end-products: a biological consequence of lifestyle contributing to cancer disparity. *Cancer Res.* **75**, 1925–1929 (2015).
- Sun, S. S. M. & Liu, Q. Transgenic approaches to improve the nutritional quality of plant proteins. *In Vitro Cell Dev. Biol.* – Plant, **40**, 155–162 (2004).

37. Beauregard, M. & Hefford, M. A. Enhancement of essential amino acid contents in crops by genetic engineering and protein design. *Plant Biotechnol. J.* **4**, 561–574 (2006).
38. Liu, J., Seul, U. & Thompson, R. Cloning and characterization of a pollen-specific cDNA encoding a glutamic-acid-rich protein (GARP) from potato *Solanum berthaultii*. *Plant Mol. Biol.* **33**, 291–300 (1997).
39. Torrent, M., Alvarez, I., Geli, M. I., Dalcol, I. & Ludevid, D. Lysine-rich modified gamma-zeins accumulate in protein bodies of transiently transformed maize endosperms. *Plant Mol. Biol.* **34**, 139–149 (1997).
40. Altenbach, S. B. *et al.* Accumulation of a Brazil nut albumin in seeds of transgenic canola results in enhanced levels of seed protein methionine. *Plant Mol. Bio.* **18**, 235–245 (1992).
41. Dinkins, R. D. *et al.* Increased sulfur amino acids in soybean plants overexpressing the maize 15 kD zein protein. *In Vitro Cell Dev. Biol. Plant.* **37**, 742–747 (2001).
42. Lee, T. T. *et al.* Enhanced methionine and cysteine levels in transgenic rice seeds by the accumulation of sesame 2S albumin. *Biosci. Biotechnol. Biochem.* **67**, 1699–1705 (2003).
43. Alvarez, I., Geli, M. I., Pimentel, E., Ludevid, D. & Torrent, M. Lysine-rich gamma-zeins are secreted in transgenic Arabidopsis plants. *Planta.* **205**, 420–427 (1998).
44. Wang, G., Wang, G., Zhang, X., Wang, F. & Song, R. Isolation of high quality RNA from cereal seeds containing high levels of starch. *Phytochem. Anal.* **23**, 159–163 (2012).
45. Hiei, Y., Ohta, S., Komari, T. & Kumashiro, T. Efficient transformation of rice (*Oryza sativa* L) mediated by Agrobacterium and sequence analysis of the boundaries of the T-DNA. *Plant J.* **6**, 271–282 (1994).
46. Jiang, S. Y. *et al.* Ds insertion mutagenesis as an efficient tool to produce diverse variations for rice breeding. *Plant Mol Biol.* **65**, 385–402 (2007).

## Acknowledgements

This research is supported by Singapore Economic Development Board Research Incentive Scheme for Companies (COY-15-RISC/120014 S12-1146-RISC) and the Singapore National Research Foundation under the Competitive Research Programme Funding Scheme (CRP award no. NRF-CRP7-2010-02).

## Author Contributions

S.R. supervised the study; S.Y.J. conceived of the study, analyzed the data and wrote the article; A.M. carried out molecular and phenotypic characterization of transgenic plants. L.X. generated transgenic plants. S.R. supervised and complemented the writing. All authors read and approved the final manuscript.

## Additional Information

**Supplementary information** accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Jiang, S.-Y. *et al.* Improving protein content and quality by over-expressing artificially synthetic fusion proteins with high lysine and threonine constituent in rice plants. *Sci. Rep.* **6**, 34427; doi: 10.1038/srep34427 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2016