

SCIENTIFIC REPORTS

OPEN

Method for detection of hydrogen peroxide in HT22 cells

Dagmara Jacewicz¹, Kamila Siedlecka-Kroplewska², Joanna Drzeżdżon¹,
Agnieszka Piotrowska¹, Dariusz Wyrzykowski¹, Aleksandra Tesmar¹, Krzysztof Żamojć¹ &
Lech Chmurzyński¹

Received: 05 December 2016


Accepted: 02 March 2017

Published: 30 March 2017

We have proposed a new method which can be applied in assessing the intracellular production of hydrogen peroxide. Using this assay we have examined the hydrogen peroxide generation during the L-glutamate induced oxidative stress in the HT22 hippocampal cells. The detection of hydrogen peroxide is based on two crucial reagents *cis*-[Cr(C₂O₄)(pm)(OH)₂]⁺ (pm denotes pyridoxamine) and 2-ketobutyrate. The results obtained indicate that the presented method can be a promising tool to detect hydrogen peroxide in biological samples, particularly in cellular experimental models.

Reactive oxygen species (ROS) are generated in cells under normal conditions e.g. during aerobic respiration¹. Electrons transferred in the mitochondrial electron transport chain may escape and interact with molecular oxygen to form superoxide anion (O₂⁻)¹. Superoxide dismutase (SOD) catalyzes dismutation of O₂⁻ resulting in the generation of hydrogen peroxide (H₂O₂), which in turn can react with Fe²⁺ in the Fenton reaction to form hydroxyl radicals (HO⁻)²⁻⁴. When the cellular antioxidant defense mechanisms fail to effectively eliminate ROS the oxidative stress appears. Superoxide anion, hydroxyl radical or hydrogen peroxide lead to irreversible damage of lipids, proteins and DNA, resulting in dysfunction of cell organelles^{5,6}. The extraordinary high rate of respiration in some organs such as the brain makes them particularly susceptible to ROS mediated injury. ROS and induced by them oxidative damage are believed to contribute to the development of many diseases such as cancer, arteriosclerosis, diabetes and neurodegenerative diseases⁷⁻¹⁰. Establishing the role of oxidative stress requires the ability to measure its mediators accurately¹¹. Therefore, there is a need to develop and improve sensitive and specific methods to detect and evaluate the level of reactive oxygen species in biological samples of different origin. A method of protecting the neuronal cells HT22 against the oxidative stress generated by hydrogen peroxide has been investigated and described by Jia and co-workers¹². This method is based on cytoprotection properties of endogenous cannabinoid anandamide. Moreover, recently it has been found that the Homer1a - protein regulating calcium pathways are involved in the glutamate-induced HT22 cell death¹³. Homer1a plays the cytoprotective role in the prevention of the glutamate-induced oxidative injury. In most cases existing methods to detect hydrogen peroxide in biological samples are based on the horseradish peroxidase (HRP) mediated reaction¹⁴. These methods are sensitive but have also disadvantages because HRP can react with many cellular substrates. Fluorescent probes used for the detection of H₂O₂ under biological conditions in addition to the advantages have some limitations. For example some probes require long response times (30–60 min) or an increase in the luminescence is also observed in the presence of the citrate and phosphate buffer¹⁵. Moreover, the autofluorescence of cells is a problem in the flow cytometry technique or fluorescence microscopy for the detection of H₂O₂. The purpose of this study was to design a method for the hydrogen peroxide detection which is characterized by a high sensitivity, moreover the autofluorescence of samples is eliminated and the results obtained by this method can be presented in concentration units.

The new method we propose is based on the following steps: (a) the nonenzymatic reaction of 2-ketobutyrate with H₂O₂ resulting in the CO₂ release and (b) the interaction of the coordination compound *cis*-[Cr(C₂O₄)(pm)(OH)₂]⁺ with CO₂. 2-ketobutyrate reacts with H₂O₂ with 1:1 stoichiometry (1). The amount of H₂O₂ in this reaction is equivalent with the CO₂ released. 2-ketoacids such as 2-ketobutyrate or pyruvate can react nonenzymatically with H₂O₂, yielding CO₂, H₂O and the corresponding one carbon atom shorter carboxylic acid. This reaction was first reported by Holleman¹⁶.



¹Faculty of Chemistry, University of Gdansk, Wita Stwosza 63, 80-308 Gdansk, Poland. ²Department of Histology, Medical University of Gdansk, Debinki 1, 80-211, Gdansk, Poland. Correspondence and requests for materials should be addressed to D.J. (email: dagmara.jacewicz@ug.edu.pl)




Figure 1. Structure of $cis-[Cr(C_2O_4)(pm)(OH_2)_2]^+$.

Thus, based on the CO_2 measurement it is possible to assess the level of H_2O_2 . The coordination complex of chromium $[Cr(C_2O_4)(pm)(OH_2)_2]^+$, where pm denotes pyridoxamine, was shown to effectively catch CO_2 ¹⁷. Previously we successfully used $cis-[Cr(C_2O_4)(pm)(OH_2)_2]^+$ as a molecular biosensor to detect CO_2 in osteosarcoma 143B cells treated with hydrogen peroxide¹⁸. In this study we demonstrate an improved version of this method based on both $cis-[Cr(C_2O_4)(pm)(OH_2)_2]^+$ and 2-ketobutyrate.

Results

To study the generation of hydrogen peroxide in a cellular model we used hippocampal HT22 cells treated with 5 mM L-glutamate for 24 h. Glutamate belongs to neurotransmitters of the central nervous system. However, at high (mM) concentration it was found to induce death of neurons¹⁹. 5 mM glutamate was reported to induce oxidative stress in HT22 cells resulting in cell death²⁰. Our results also showed that the viability of HT22 cells treated with 5 mM L-glutamate for 24 h dramatically decreases (publication in press). On the basis of our proposed hydrogen peroxide detection method, we measured CO_2 concentration after adding of 2-ketobutyrate. 2-ketobutyrate reacts with H_2O_2 . The amount of CO_2 released in this reaction reflects the amount of H_2O_2 . In addition to cell lysates, the CO_2 level was assessed in supernatants as well as in cell free culture medium – all of them potentially contained H_2O_2 . It is important to note that they differed in chemical composition. In order to determine the most effective concentration of 2-ketobutyrate required to scavenge H_2O_2 we tested its following concentrations: 0.5 mM, 1 mM, 2 mM, 3 mM, 4 mM, 5 mM, 6 mM, 7 mM, 8 mM, 9 mM, 10 mM, 11 mM, 12 mM, 13 mM, 14 mM and 15 mM.

In order to assess the CO_2 concentration in samples we used a specific molecular biosensor $cis-[Cr(C_2O_4)(pm)(OH_2)_2]^+$ and the spectrophotometric stopped-flow technique. The structure of the synthesized complex of chromium(III), $cis-[Cr(C_2O_4)(pm)(OH_2)_2]^+$ ion, is shown in Fig. 1²¹.

It turned out that the coordination ion, $cis-[Cr(C_2O_4)(pm)(OH_2)_2]^+$, could be successfully applied in the case of the detection of carbon dioxide generated in the reaction of decarboxylation of 2-ketobutyrate caused by H_2O_2 . This reaction between the Cr(III) complex ion with pyridoxamine and the carbon dioxide was observed between 340–700 nm by using spectrophotometric stopped-flow method. The concentration of carbon dioxide has been identified based on the method which was used and described in the paper²².

As shown in Fig. 2 the CO_2 levels in lysates of cells incubated in the presence or absence of 5 mM L-glutamate, both measured without the addition of 2-ketobutyrate, were $0.025 \mu\text{mol mg}^{-1}$ protein (± 0.006) and $0.006 \mu\text{mol mg}^{-1}$ protein (± 0.002), respectively. We found that after addition of 0.5–10 mM 2-ketobutyrate to the lysates of L-glutamate-treated HT22 cells, the CO_2 level gradually increased (Fig. 2). Noteworthy, at 2-ketobutyrate concentrations ranging from 10 mM to 15 mM the CO_2 level was very similar (ranging from $0.14 \mu\text{mol mg}^{-1}$ protein to $0.15 \mu\text{mol mg}^{-1}$ protein). Moreover, the highest CO_2 level was then observed. CO_2 was also detected in lysates of cells incubated in the absence of 5 mM L-glutamate. However, its level was much lower than that in the presence of L-glutamate.

Intracellularly produced H_2O_2 has the ability to penetrate biological membranes and affect neighbouring cells²³. In order to better evaluate and understand the chemical environment of cells in our experimental model, we assessed the CO_2 level in supernatants. The results revealed that the CO_2 levels in supernatants separated from cells incubated in the presence or absence of 5 mM L-glutamate, both measured without the addition of 2-ketobutyrate, were $1.315 \mu\text{M}$ (± 0.220) and $0.949 \mu\text{M}$ (± 0.114), respectively (Fig. 3). We found that the CO_2 level gradually increased depending on the concentration of 2-ketobutyrate added (Fig. 3). In the case of supernatants separated from cells incubated in the presence of L-glutamate, at 2-ketobutyrate concentrations ranging from 10 mM to 15 mM the CO_2 level was very similar and the highest (ranging from $7.15 \mu\text{M}$ to $7.6 \mu\text{M}$). Moreover, the concentration of CO_2 detected in supernatants separated from cells incubated in the absence of L-glutamate was then much lower than that in the presence of L-glutamate.

The oxidation of chemical components of the culture medium may be a source of hydrogen peroxide generation^{24,25}. The source of hydrogen peroxide in the cell-free DMEM medium, used in this study, in the absence of




Figure 2. CO₂ assessment in cell lysates. HT22 cells were incubated for 24 h in the presence or absence of 5 mM sodium L-glutamate. After treatment, cells were lysed. Before the CO₂ measurement sodium 2-ketobutyrate was added (final concentrations: 0.5–15 mM, respectively). The CO₂ level was determined using *cis*-[Cr(C₂O₄)₂(pm)(OH₂)₂]⁺ and a stopped-flow technique. Data are presented as mean ± SD of three independent experiments. *p < 0.05, ***p < 0.001, statistically significant differences between samples incubated in the presence and samples incubated in the absence of 5 mM L-glutamate.




Figure 3. CO₂ assessment in supernatants. HT22 cells were incubated for 24 h in the presence or absence of 5 mM sodium L-glutamate. After treatment, supernatants were separated from cells. Before the CO₂ measurement sodium 2-ketobutyrate was added (final concentrations: 0.5–15 mM, respectively). The CO₂ level was determined using *cis*-[Cr(C₂O₄)₂(pm)(OH₂)₂]⁺ and a stopped-flow technique. Data are presented as mean ± SD of three independent experiments. *p < 0.05, statistically significant differences between samples incubated in the presence and samples incubated in the absence of 5 mM L-glutamate.

L-glutamate could be the oxidation of some chemical components of the culture medium. It has been reported that thiol compounds such as cysteine are unstable in culture media and can oxidize to generate hydrogen peroxide²⁴. Noteworthy, cysteine is a constituent of the DMEM medium. There is also evidence that riboflavin may be a source of the photogeneration of ROS in cell culture media²⁵. It has been shown that it is possible to generate superoxide anion and hydrogen peroxide as a product of its dismutation. Taking it into consideration, we have examined the CO₂ level in the cell-free complete culture medium. The CO₂ concentration in medium samples incubated in the presence or absence of 5 mM L-glutamate, both measured without the addition of 2-ketobutyrate, were 1.153 μM (±0.206) and 0.998 μM (±0.068), respectively (Fig. 4). We found that upon addition of 2-ketobutyrate at increasing concentrations, the CO₂ level gradually increased (Fig. 4). The increase in CO₂ concentration was observed in medium samples incubated in the presence or absence of 5 mM L-glutamate. Noteworthy, in the presence of L-glutamate after adding of 2-ketobutyrate at concentrations ranging from 12 mM to 15 mM the CO₂ level was the highest and nearly constant (about 4.5 μM). In the absence of L-glutamate, at 2-ketobutyrate concentrations ranging from 6 mM to 15 mM only slight changes in the CO₂ concentration appeared (ranging from 3.6 μM to 3.8 μM).

Discussion

In this study, we demonstrate a new method of hydrogen peroxide detection. It can be applied to assess hydrogen peroxide level in different biological samples, e.g. cell lysates, supernatants or cell-free culture medium.

Our results suggest that the 10 mM–15 mM concentration range of 2-ketobutyrate can be used in the proposed method to effectively detect and assess the CO₂ concentration in lysates of 5 mM L-glutamate-treated HT22 cells. At this concentration range the CO₂ level was nearly the same. The main source of CO₂ in cell lysates was then the reaction of 2-ketobutyrate with intracellularly produced H₂O₂. In order to find out whether the cell metabolism may affect the CO₂ level, we incubated HT22 cells in the absence of 5 mM L-glutamate. We assume that in our experimental model CO₂, whose main source are metabolic pathways, can be detected in the absence of L-glutamate (in the absence of H₂O₂ generating agents) and simultaneously in the absence of 2-ketobutyrate.




Figure 4. CO₂ assessment in the cell-free complete culture medium. The cell-free culture medium was incubated for 24 h in the presence or absence of 5 mM sodium L-glutamate. After treatment, sodium 2-ketobutyrate was added (final concentrations: 0.5–15 mM, respectively). The CO₂ level was determined using *cis*-[Cr(C₂O₄)₂(pm)(OH)₂]⁺ and a stopped-flow technique. Data are presented as mean ± SD of three independent experiments. *p < 0.05, **p < 0.01 statistically significant differences between samples incubated in the presence and samples incubated in the absence of 5 mM L-glutamate.

Under these conditions CO₂ does not originate from the reaction of 2-ketobutyrate with H₂O₂ as well as is not affected by L-glutamate, hence its main source can probably be metabolic pathways. Our results have shown that the CO₂ level in lysates of cells incubated in the absence of 5 mM L-glutamate, measured without the addition of 2-ketobutyrate, is 0.006 µmol mg⁻¹ protein (±0.002). It reflects the background CO₂ level for lysates of cells nontreated with L-glutamate. The CO₂ level measured in the absence of L-glutamate, but after the addition of 2-ketobutyrate includes CO₂ originating from metabolic pathways. Another source can be endogenous H₂O₂ related to the function of some cell organelles, e.g. peroxisomes, mitochondria. Noteworthy, the CO₂ level in lysates of cells incubated in the presence of 5 mM L-glutamate, measured without the addition of 2-ketobutyrate, is 0.025 µmol mg⁻¹ protein (±0.006). It may be considered as the background CO₂ level for lysates of cells treated with L-glutamate. It appears slightly higher than in the absence of 5 mM L-glutamate suggesting that 5 mM L-glutamate may affect the CO₂ level. The background signal may be a disadvantage of the proposed method. However, a similar problem or disadvantage related to the background signal exists in many other experimental methods. Cellular autofluorescence, for example, resulting from natural fluorescence properties of some structural components of cells, affects the sensitivity of flow cytometry assays^{26–28}. Autofluorescence of cells or tissues is also a problem in another widely used technique such as fluorescence microscopy^{29,30}. The background fluorescence interferes with detection of specific fluorescent signals. In our proposed method the problem related to the background CO₂ level in cell lysates can be solved. The possible solution could be to subtract the background signal from the analysis.

The results indicated that, the 10 mM–15 mM concentration range of 2-ketobutyrate is sufficient to assess the CO₂ level in supernatants separated from 5 mM L-glutamate-treated HT22 cells. The CO₂ level in the samples measured was then similar and stopped increasing. As mentioned previously, the main source of CO₂ in supernatants was H₂O₂ released from the cells where it was produced²³. However, the additional source of CO₂ can also be H₂O₂ generated during oxidation of chemical components of the culture medium^{24,25}. Therefore, we also determined the CO₂ level in supernatants separated from cells incubated in the absence of 5 mM L-glutamate. We noticed that it was much lower than in the presence of 5 mM L-glutamate. CO₂ detected in supernatants separated from cells nontreated with 5 mM L-glutamate, measured without the addition of 2-ketobutyrate, is 0.949 µM (±0.114) and corresponds to the background CO₂ level in supernatants separated from cells nontreated with 5 mM L-glutamate. It results mostly from the oxidation of chemical components of the culture medium. CO₂ detected in the supernatant separated from cells treated with 5 mM L-glutamate, measured without the addition of 2-ketobutyrate, is 1.315 µM (±0.220) and reflects the background CO₂ level in supernatants separated from cells treated with 5 mM L-glutamate. Its source is related not only to the oxidation of chemical components of the culture medium, but also to the effects of L-glutamate on culture medium components.

We found that 2-ketobutyrate at a concentration from 12 mM to 15 mM can be used to assess the CO₂ level in the cell-free culture medium incubated with 5 mM L-glutamate. The CO₂ concentration was then about 0.8 µM higher in the presence of L-glutamate than in its absence, suggesting that 5 mM L-glutamate may generate a small amount of H₂O₂ and/or CO₂ in the culture medium. The CO₂ level (corresponding to H₂O₂) in the presence of L-glutamate reflects the oxidation of components of the culture medium as well as the influence of L-glutamate on culture medium components. The CO₂ level measured without the addition of 2-ketobutyrate can be considered as the background CO₂ level in the cell-free culture medium. In the presence of 5 mM L-glutamate it is 1.153 µM (±0.206), whereas in the absence of 5 mM L-glutamate 0.998 µM (±0.068). The background CO₂ level may result mainly from the generation of CO₂ during the oxidation/decomposition of some its components. The CO₂ level in the presence of L-glutamate as well as the CO₂ level in the absence of L-glutamate, both measured after addition of 2-ketobutyrate, include the background CO₂ level. However, it is important to note that the CO₂ level measured in cell lysates should not be affected by the background CO₂ in culture medium/supernatants. Before analysis, cells are separated from the culture medium, washed and then lysed to obtain cell lysates.

It is important to note that cell lysates, supernatants and cell-free culture medium differ in chemical composition. Thus, they should be considered examples of different experimental models. The chemical composition of culture media is considered to resemble (at least in part) the natural cellular environment. It should imitate physiological conditions. However, further studies are needed to find out whether the method presented here can be applied to detect hydrogen peroxide in biological fluids, e.g. in the blood serum. Our results revealed that the concentration range of 2-ketobutyrate needed to assess the CO₂ level in different models may not be the same. This highlights the importance of preliminary studies required to establish experimental conditions in every case. However, it is not a big disadvantage, because in many experiments it is a common procedure, e.g. a standard curve in protein measurements.

Conclusions

The new method we propose can be useful in comparative studies, e.g. unknown samples can be compared to control samples. An advantage of this method is its high sensitivity. The lower limit of detection is equal to 10⁻⁷ M²². Another advantage of the method is that the results can be presented in concentration units. Majority of methods allow to use arbitrary instead of more precise concentration units, e.g. flow cytometry assays using 2',7'-dichlorofluorescein diacetate or hydroethidine to measure intracellular production of reactive oxygen species^{31,32}. In conclusion, the presented method based on both *cis*-[Cr(C₂O₄)(pm)(OH₂)₂]⁺ and 2-ketobutyrate seems to be a very promising tool to study the hydrogen peroxide generation.

Methods

Chemicals. Reagents: sodium 2-ketobutyrate and L-glutamic acid monosodium salt monohydrate were purchased from Sigma (USA). L-glutamic acid monosodium salt solutions were prepared in the sterile physiological saline solution before use. Each time sodium 2-ketobutyrate solutions before use were prepared in sterile water. *Cis*-[Cr(C₂O₄)(pm)(OH₂)₂]⁺ ion was synthesized according to standard literature procedures. The final product - *cis*-[Cr(C₂O₄)(L-L)(O₂CO)]⁻ (L-L denotes bidentate ligand - pyridoxamine (pm)) was prepared by the known method described in the literature¹⁷.

Cell culture. The mouse hippocampal neuronal HT22 cell line was kindly provided by Professor T. Grune (Institute of Biological Chemistry and Nutrition, University Hohenheim, Stuttgart). HT22 cells were maintained in CO₂ incubator, at 37 °C in a humidified atmosphere containing 5% CO₂. Cells were cultured in Dulbecco's Modified Eagle's Medium without sodium pyruvate (DMEM, Sigma-Aldrich, USA), supplemented with 10% heat-inactivated fetal bovine serum (Sigma-Aldrich, USA), 100 IU mL⁻¹ penicillin (Sigma-Aldrich, USA) and 100 µg/mL streptomycin (Sigma-Aldrich, USA).

Cell treatment and CO₂ measurement. HT22 cells were seeded onto 6-well plates (2 × 10⁵ cells per well) and allowed to attach for 24 h. Cells were then incubated for 24 h in the presence or absence of 5 mM sodium L-glutamate. Additionally, the cell-free culture medium (DMEM) was incubated for 24 h with or without 5 mM sodium L-glutamate. After incubation, cells as well as cell culture medium in which the cells were incubated were collected and separated by centrifugation (300 × g, 5 min, room temperature). Next, HT22 cells were washed with PBS and suspended in a lysis buffer (0.15 M NaCl, 0.005 M EDTA, 1% Triton X-100, 0.01 M Tris-HCl). The cell-free complete culture medium was also collected. Sodium 2-ketobutyrate solution was then added (final concentration: 0.5 mM, 1 mM, 2 mM, 3 mM, 4 mM, 5 mM, 6 mM, 7 mM, 8 mM, 9 mM, 10 mM, 11 mM, 12 mM, 13 mM, 14 mM, 15 mM) to the samples (cell lysates, supernatants, cell-free culture medium), prior to CO₂ measurement. The protein level in cell lysates was measured using Pierce BCATM Protein Assay Kit (Thermo Scientific, USA) according to the manufacturer's instruction. The CO₂ concentration was assessed using a spectrophotometric stopped-flow technique and a coordinate *cis*-[Cr(C₂O₄)(pm)(OH₂)₂]⁺ ion as a molecular biosensor of CO₂.

Statistical analysis. Statistical analysis was performed using the Statistica 9 software (StatSoft, Poland). Data are expressed as mean ± SD. Statistical differences were evaluated using the Mann-Whitney U test. Differences were considered significant at p < 0.05, p < 0.01, p < 0.001, respectively.

References

1. Grivennikova, V. G. & Vinogradov, A. D. Generation of superoxide by the mitochondrial Complex I. *Biochim Biophys Acta*. **1757**, 553–561 (2006).
2. McCord, J. M. & Fridovich, I. Superoxide dismutase an enzymic function for erythrocuprein (hemocuprein). *J Biol Chem*. **244**, 6049–6055 (1969).
3. Boveris, A. & Chance, B. The mitochondrial generation of hydrogen peroxide General properties and effect of hyperbaric oxygen. *Biochem J*. **134**, 707–716 (1973).
4. Fenton, H. J. H. Oxidation of tartaric acid in presence of iron. *J Chem Soc*. **65**, 899–910 (1894).
5. Pranczk, J., Jacewicz, D., Wyrzykowski, D. & Chmurzynski, L. Analytical methods for determination of reactive oxygen species. *Curr. Pharm. Anal.* **10**, 293–304 (2014).
6. Wyrzykowski, D., Inkielewicz-Stepniak, I., Pranczk, J., Żamojć, K., Zięba, P., Tesmar, A., Jacewicz D., Ossowski, T. & Chmurzyński, L. Physicochemical properties of ternary oxovanadium(IV) complexes with oxydiacetate and 1, 10-phenanthroline or 2, 2'-bipyridine. Cytoprotective activity in hippocampal neuronal HT22 cells. *BioMetals* **28**, 307–320 (2015).
7. Friedman, J., Peleg, E., Kagan, T., Shnizer, S. & Rosenthal, T. Oxidative stress in hypertensive, diabetic, and diabetic hypertensive rats. *Am. J. Hypertens.* **16**, 1049–1052 (2003).
8. Reynolds, A., Laurie, C., Mosley, R. L. & Gendelman, H. E. Oxidative stress and the pathogenesis of neurodegenerative disorders. *Int. Rev. Neurobiol.* **82**, 297–325 (2007).
9. Schleicher, E. & Friess, U. Oxidative stress, AGE, and atherosclerosis. *Kidney Int.* **106**, 17–26 (2007).
10. Reuter, S., Gupta, S. C., Chaturvedi, M. M. & Aggarwal, B. B. Oxidative stress, inflammation, and cancer: how are they linked. *Free Radical Bio. Med.* **49**, 1603–1616 (2010).

11. Pranczk, J., Jacewicz, D., Wyrzykowski, D. & Chmurzynski, L. Platinum(II) and palladium(II) complex compounds as anti-cancer drugs. Methods of cytotoxicity determination. *Curr. Pharm. Anal.* **10**, 2–9 (2014).
12. Jia, J., Ma, L., Wu, M., Zhang, L., Zhang, X., Zhai, Q., Jiang, T., Wang, Q. & Xiong, L. Anandamide protects HT22 cells exposed to hydrogen peroxide by inhibiting CB1 receptor-mediated type 2 NADPH oxidase. *Oxid. Med. Cell Longev.*, doi: 10.1155/2014/893516 (2014).
13. Rao, W., Peng, C., Zhang, L., Su, N., Wang, K., Hui, H. & Fei, Z. Homer1a attenuates glutamate-induced oxidative injury in HT-22 cells through regulation of store-operated calcium entry. *Sci. Rep.* **6**, 33975 (2016).
14. Rhee, S. G., Chang, T. S., Jeong, W. & Kang, D. Methods for detection and measurement of hydrogen peroxide inside and outside of cells. *Mol. cells* **29**, 539–549 (2010).
15. Żamojć, K., Zdrowowicz, M., Jacewicz, D., Wyrzykowski, D. & Chmurzyński, L. Fluorescent probes used for detection of hydrogen peroxide under biological conditions. *Cr. Rev. Anal. Chem.* **46**, 171–200 (2016).
16. Holleman, M. A. F. Notice sur l'action de l'eau oxygénée sur les acides (-cetoniques et sur les dicétones 1.2. *Recl. Trav. Chim. Pays-bas Belg.* **23**, 169–172 (1904).
17. Jacewicz, D., Dąbrowska, A., Banecki, B., Kolisz, I., Woźniak, M. & Chmurzyński, L. A stopped-flow study on the kinetics and mechanism of CO₂ uptake by chromium(III) complexes with histamine and pyridoxamine. *Transit. Metal Chem.* **30**, 209–216 (2005).
18. Jacewicz, D., Szkatuła, M., Chylewska, A., Dąbrowska, A. & Chmurzyński, L. Coordinate *cis*-[Cr(C₂O₄)(pm)(OH)₂]⁺ cation as molecular biosensor of pyruvates protective activity against hydrogen peroxide mediated cytotoxicity. *Sensors* **8**, 4487–4504 (2008).
19. Coyle, J. T. & Puttfarcken, P. Oxidative stress, glutamate, and neurodegenerative disorders. *Sci.* **262**, 689–695 (1993).
20. Yoon, S. W., Kang, S., Ryu, S. E. & Poo, H. Identification of tyrosine-nitrated proteins in HT22 hippocampal cells during glutamate-induced oxidative stress. *Cell Prolif.* **43**, 584–593 (2010).
21. Kita, E. Model quasi-enzyme compounds of chromium(III) with vitamins B6 and histamine. *Pol J Chem.* **61**, 361–367 (1987).
22. Jacewicz, D., Żamojć, K., Wyrzykowski, D. & Chmurzyński, L. Analytical methods for determination of CO and CO₂ and their applicability in biological studies. *Curr. Pharm. Anal.* **9**, 226–235 (2013).
23. Halliwell, B. Antioxidants in human health and disease. *Ann. Rev. Nutr.* **16**, 33–50 (1996).
24. Hua Long, L. & Halliwell, B. Oxidation and generation of hydrogen peroxide by thiol compounds in commonly used cell culture media. *Biochem. Biophys. Res. Commun.* **286**, 991–994 (2001).
25. Grzelak, A., Rychlik, B. & Bartosz, G. Light-dependent generation of reactive oxygen species in cell culture media. *Free Radical Bio. Med.* **30**, 1418–1425 (2001).
26. Aubin, J. E. Autofluorescence of viable cultured mammalian cells. *J. Histochem. Cytochem.* **27**, 36–43 (1979).
27. Alberti, S., Parks, D. R. & Herzenberg, L. A. A single laser method for subtraction of cell auto fluorescence in flow cytometry. *Cytometry* **8**, 114–119 (1987).
28. Mosiman, V. L., Patterson, B. K., Canterero, L. & Goolsby, C. L. Reducing cellular autofluorescence in flow cytometry: an *in situ* method. *Cytometry* **30**, 151–156 (1997).
29. Steinkamp, J. A. & Stewart, C. C. Dual-laser, differential fluorescence correction method for reducing cellular background autofluorescence. *Cytometry* **7**, 566–574 (1986).
30. Neumann, M. & Gabel, D. Simple method for reduction of autofluorescence in fluorescence microscopy. *J. Histochem. Cytochem.* **50**, 437–439 (2002).
31. Ubezio, P. & Civoli, F. Flow cytometric detection of hydrogen peroxide production induced by doxorubicin in cancer cells. *Free Radical Bio. Med.* **16**, 509–516 (1994).
32. Rothe, G. & Valet, G. Flow cytometric analysis of respiratory burst activity in phagocytes with hydroethidine and 2',7'-dichlorofluorescein. *J. Leukocyte Biol.* **47**, 440–448 (1990).

Acknowledgements

This work was supported by National Science Centre, Poland grant numbers 2015/19/N/ST5/00276

Author Contributions

D.J. and K.S.-K. designed the study and wrote the paper. D.J., K.S.-K., J.D., A.P., D.W., A.T. and K.Ż. conducted the experiments and analysed the data. L.C. supervised the work.

Additional Information

Competing Interests: The authors declare no competing financial interests.

How to cite this article: Jacewicz, D. *et al.* Method for detection of hydrogen peroxide in HT22 cells. *Sci. Rep.* **7**, 45673; doi: 10.1038/srep45673 (2017).

Publisher's note: Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>

© The Author(s) 2017