



CORRECTION

Correction: Associations of measures of systemic blood flow used in a randomized trial of delayed cord clamping in preterm infants

Himanshu Popat^{1,2}, Kristy P. Robledo^{2,3}, Adrienne Kirby^{2,3}, Lucille Sebastian^{2,3}, Nicholas Evans^{2,4}, Andrew Gill⁵, Martin Kluckow^{2,6}, Sanjay Sinhal⁷, Koert de Waal⁸, William Tarnow-Mordi^{2,3} and David Osborn^{2,4}

Pediatric Research (2019) 86:138; <https://doi.org/10.1038/s41390-019-0382-z>

Correction to: *Pediatric Research* <https://doi.org/10.1038/s41390-019-0348-1>, published online 21 February 2019

In the original version of this article, five of the row labels (Treated ROP, Chronic lung disease, PDA—Treatment,

PDA—Surgery, NEC—Surgery) in Table 3 were inadvertently interchanged. This has now been corrected in both the PDF and HTML versions of the article. The original, incorrect table is displayed here for reference.

Table 3. Low systemic blood flow expressed as SVC flow ≤ 55 ml/kg/min and RVO < 150 ml/kg/min and clinical outcomes at 36 weeks postmenstrual age (n/N (%) or mean (SD). Relative risk (95% CI) unadjusted

	SVC ≤ 55 ml/kg/min	SVC > 55 ml/kg/min	Relative risk (95% CI)	P value	RVO ≤ 150 ml/kg/min	RVO > 150 ml/kg/min	Relative risk (95% CI)	P value
Death by 36 weeks	6/86 (7.0%)	8/177 (4.5%)	1.27 (0.30, 5.37)	0.71	9/81 (11.1%)	5/183 (2.7%)	3.83 (1.39, 10.56)	0.04
IVH grade 3 or 4	5/86 (5.8%)	6/175 (3.4%)	1.14 (0.04, 31.14)	0.90	6/79 (7.6%)	5/183 (2.7%)	2.41 (0.83, 6.99)	0.21
Any IVH	16/86 (18.6%)	25/175 (14.3%)	1.17 (0.67, 2.05)	0.59	18/79 (22.8%)	23/183 (12.6%)	1.82 (1.04, 3.20)	0.06
Late IVH	9/85 (10.6%)	18/169 (10.7%)	0.92 (0.43, 1.96)	0.83	13/78 (16.7%)	14/177 (7.9%)	2.16 (1.05, 4.45)	0.06
PVL	1/82 (1.2%)	1/172 (0.6%)	2.10 (0.13, 32.96)	0.64	0/75 (0.0%)	2/180 (1.1%)	NA	NA
NEC	5/86 (5.8%)	8/175 (4.6%)	1.32 (0.46, 3.77)	0.63	4/79 (5.1%)	9/183 (4.9%)	1.10 (0.35, 3.44)	0.87
NEC—surgery	5/82 (6.1%)	8/170 (4.7%)	1.29 (0.43, 3.83)	0.66	5/74 (6.8%)	8/179 (4.5%)	1.53 (0.53, 4.48)	0.47
Treated ROP	47/80 (58.8%)	76/168 (45.2%)	1.26 (0.98, 1.63)	0.09	43/71 (60.6%)	81/178 (45.5%)	1.27 (0.99, 1.62)	0.08
Chronic lung disease	46/86 (53.5%)	50/175 (28.6%)	1.68 (1.25, 2.27)	0.002	34/79 (43.0%)	62/183 (33.9%)	1.31 (0.97, 1.76)	0.10
PDA—treatment	2/74 (2.7%)	3/132 (2.3%)	1.21 (0.21, 7.03)	0.84	1/66 (1.5%)	4/141 (2.8%)	0.53 (0.06, 4.69)	0.51
PDA—surgery	0/71 (0.0%)	3/139 (2.2%)	NA	NA	2/64 (3.1%)	1/147 (0.7%)	4.69 (0.43, 51.16)	0.28
Blood transfusion	57/86 (66.3%)	82/175 (46.9%)	1.38 (1.13, 1.69)	0.004	47/79 (59.5%)	92/183 (50.3%)	1.12 (0.89, 1.43)	0.34
Peak HCT in first 7 days, %	49.7 (7.9)	50.0 (8.4)	−0.14 (−2.25, 1.98)	0.90	51.7 (8.8)	49.1 (7.8)	2.48 (0.22, 4.74)	0.03
Peak Bilirubin in first 7 days, $\mu\text{mol/L}$	148.5 (37.1)	151.2 (39.8)	−2.52 (−11.2, 6.17)	0.57	147.7 (38.4)	151.7 (39.2)	−3.37 (−13.1, 6.41)	0.50

IVH intraventricular hemorrhage, PVL periventricular leukomalacia, NEC necrotizing enterocolitis, ROP retinopathy of prematurity, PDA patent ductus arteriosus, HCT hematocrit, NA not applicable

¹Children's Hospital at Westmead, Westmead, NSW, Australia; ²University of Sydney, Camperdown, Australia; ³NHMRC Clinical Trials Centre, Camperdown, NSW, Australia; ⁴Royal Prince Alfred Hospital, Camperdown, NSW, Australia; ⁵Centre for Neonatal Education and Research, University of Western Australia, Perth, WA, Australia; ⁶Royal North Shore Hospital, St Leonards, NSW, Australia; ⁷Flinders Medical Centre, Bedford Park, SA, Australia and ⁸John Hunter Children's Hospital, New Lambton, NSW, Australia
Correspondence: Himanshu Popat (himanshu.popat@health.nsw.gov.au)