

ARTICLE

Open Access

Auxin regulates anthocyanin biosynthesis through the Aux/IAA–ARF signaling pathway in apple

Yi-cheng Wang^{1,2}, Nan Wang^{1,2}, Hai-feng Xu^{1,2}, Sheng-hui Jiang^{1,2}, Hong-cheng Fang^{1,2}, Meng-yu Su^{1,2}, Zong-ying Zhang^{1,2}, Tian-liang Zhang^{1,2} and Xue-sen Chen^{1,2}

Abstract

Auxin signaling, which is crucial for normal plant growth and development, mainly depends on ARF–Aux/IAA interactions. However, little is known regarding the regulatory effects of auxin signaling on anthocyanin metabolism in apple (*Malus domestica*). We investigated the functions of MdARF13, which contains a repression domain and is localized to the nucleus. This protein was observed to interact with the Aux/IAA repressor, MdIAA121, through its C-terminal dimerization domain. Protein degradation experiments proved that MdIAA121 is an unstable protein that is degraded by the 26S proteasome. Additionally, MdIAA121 stability is affected by the application of exogenous auxin. Furthermore, the overexpression of *MdIAA121* and *MdARF13* in transgenic red-fleshed apple calli weakened the inhibitory effect of MdARF13 on anthocyanin biosynthesis. These results indicate that the degradation of MdIAA121 induced by auxin treatment can release MdARF13, which acts as a negative regulator of the anthocyanin metabolic pathway. Additionally, yeast two-hybrid, bimolecular fluorescence complementation, and pull-down assays confirmed that *MdMYB10* interacts with MdARF13. A subsequent electrophoretic mobility shift assay and yeast one-hybrid assay demonstrated that MdARF13 directly binds to the promoter of *MdDFR*, which is an anthocyanin pathway structural gene. Interestingly, chromatin immunoprecipitation–quantitative real-time PCR results indicated that the overexpression of *MdIAA121* clearly inhibits the recruitment of MdARF13 to the *MdDFR* promoter. Our findings further characterized the mechanism underlying the regulation of anthocyanin biosynthesis via Aux/IAA–ARF signaling.

Introduction

Color is an important fruit quality index that influences the purchasing behavior of consumers to some degree¹. As a type of secondary metabolite, anthocyanins constitute one of the main pigments responsible for fruit color^{2,3}. Additionally, their positive effects on human health include enhancing vascular elasticity, preventing cardiovascular disease, and protecting the liver from damage^{4–7}.

Three main protein families (MYB, bHLH, and WD40) regulate anthocyanin biosynthesis by forming the MYB–bHLH–WD40 complex^{8,9}. The MYB transcription factors (TFs) belong to one of the largest plant TF families, with some members associated with anthocyanin biosynthesis having been isolated and identified in many plant species, including *Arabidopsis thaliana*, maize, strawberry, and grape^{10–13}. In apple, the three genes encoding MYB activators (*MYB10*, *MYB1*, and *MYBA*) involved in anthocyanin biosynthesis are alleles¹⁴. Two MYB TF genes (*MdMYB1* and *MdMYBA*), which were first isolated from fruit skin, have been confirmed to be responsible for the accumulation of anthocyanin^{15,16}. Furthermore, *MdMYB10* consists of a characteristic R6 rearrangement in its promoter that is

Correspondence: X.-s. Chen (chanyetixi@163.com)

¹State Key Laboratory of Crop Biology, College of Horticulture Science and Engineering, Shandong Agricultural University, Tai-An, Shandong, China

²Collaborative Innovation Center of Fruit & Vegetable Quality and Efficient Production in Shandong, Tai-An, Shandong, China

associated with increased anthocyanin biosynthesis in red-fleshed apples¹⁷.

Anthocyanin biosynthesis is influenced by environmental stimuli (e.g., light and temperature) and plant hormones (e.g., auxin and jasmonate)^{18–21}. Moreover, auxins (naphthalene acetic acid (NAA) or 2,4-dichlorophenoxyacetic acid (2,4-D)) regulate secondary metabolic pathways, including those related to phenylpropanoid, flavonoid, and anthocyanin metabolism^{22–25}. An earlier study involving a carrot suspension culture revealed that 2,4-D strongly inhibits anthocyanin biosynthesis²⁰. Additionally, the application of exogenous NAA and 2,4-D can decrease the anthocyanin contents of transgenic tobacco calli²⁴. In apple, studies involving red-fleshed apple calli revealed that increasing auxin concentrations can significantly inhibit anthocyanin biosynthesis^{26,27}. However, the role of the auxin signaling pathway during the regulation of anthocyanin biosynthesis is unclear.

Auxin (indole-3-acetic acid (IAA)) regulates plant development by inducing rapid cellular responses and changes in gene expression. Auxin response factors (ARFs) together with auxin/IAA (Aux/IAA) proteins are TFs that function as key regulators of auxin-responsive transcription in plants^{28–30}. Current auxin signaling models indicate that auxin responses mainly depend on the interaction between the homologous C-terminal domains of Aux/IAA and ARF proteins^{31,32}. When cellular auxin concentrations are low, ARF activators at the promoters of auxin-responsive genes are thought to be inactive because of their association with Aux/IAA repressors^{29,31}. When auxin concentrations increase, the Aux/IAA repressors are recruited to auxin receptors and degraded via the ubiquitin–proteasome pathway^{33,34}. This degradation enables ARF activators to repress/activate the expression of auxin-responsive genes³³.

Most ARFs consist of an N-terminal DNA-binding domain (DBD), a middle region that functions as an activation domain (AD) or a repression domain (RD), and a C-terminal dimerization domain (CTD)³⁵. ARFs can specifically bind to the TGTCTC auxin-response element (AuxRE) in the promoters of primary/early auxin-response genes³⁵. In the last few years, several new details regarding the role of ARFs in plant growth and development have been revealed. For example, AtARF3 and AtARF4 are important for the development of reproductive and vegetative tissues^{36,37}. Additionally, AtARF5 is involved in embryonic and vascular tissue development^{38,39}, while OsARF8 mediates hypocotyl elongation and influences auxin homeostasis⁴⁰. Meanwhile, SIARF7 may negatively regulate fruit set, whereas the upregulated expression of SIARF10 suggests that the encoded protein positively affects fruit development^{41,42}. Thus, ARFs exhibit dynamic functions in plants and have a central role in the hub of transcriptional networks.

In the present study, we investigated the effects of different auxin concentrations on anthocyanin biosynthesis in red-fleshed apple calli. We also characterized an *ARF* gene, *MdARF13*, and confirmed that the *MdARF13*-mediated auxin signaling pathway helps regulate anthocyanin biosynthesis in apple, thereby revealing a novel metabolic function for ARF proteins.

Materials and methods

Plant materials and hormone treatments

Calli induced as previously described were cultured on Murashige and Skoog (MS) medium supplemented with 2 μmol/L 6-benzylaminopurine and 5, 10, 20, or 40 μmol/L NAA⁴³. The pH of all media was adjusted to 5.8 ± 0.1. The red-fleshed apple calli were incubated under a 16 h light/8 h dark photoperiod (light intensity: 1000–2000 lx) for 18 days before being harvested.

Anthocyanin extraction and absorbance measurements

Harvested calli (0.5 g each) were ground to a powder in liquid nitrogen and then treated with 15 mL 1% (v/v) HCl-methanol for 24 h at 4 °C in darkness. The solutions were then centrifuged for 10 min at 8000 × g. The absorbance of the supernatants was measured at 530 nm using a UV-2450 spectrophotometer (Shimadzu, Kyoto, Japan).

RNA isolation and quantitative real-time polymerase chain reaction

Total RNA was isolated using an RNAPrep Pure Plant Kit (Tiangen, Beijing, China). First-strand complementary DNA (cDNA) was synthesized using the RevertAid First-Strand cDNA Synthesis Kit (Fermentas, St. Leon-Roth, Germany). Three replicates were prepared for each sample. Quantitative real-time polymerase chain reaction (qPCR) analyses were conducted using 50 ng/μL cDNA as the template, SYBR Green PCR Master Mix (TransGen Biotech, Beijing, China), and the iCycler iQ5 system (Bio-Rad, Hercules, CA, USA). Details regarding the qPCR primers are provided in Supplementary Table S1. *MdActin* was used as the internal control.

Amino acid sequence analysis and phylogenetic tree construction

The ARF amino acid sequences were obtained from the NCBI (National Center for Biotechnology Information) database and aligned using the DNAMAN software (Lynnon Biosoft, USA). A phylogenetic tree was constructed according to the neighbor-joining method using the MEGA5.0 program.

Red-fleshed apple callus transformation

The intact *MdARF13* coding sequence (CDS) was inserted into the pRI101 vector containing the 35S promoter and a *green fluorescent protein* (*GFP*) tag sequence to generate




Fig. 1 Red-fleshed apple calli on medium supplemented with different NAA concentrations (5, 10, 20, and 40 $\mu\text{mol/L}$). **a** Phenotypes of calli treated with different NAA concentrations, as well as the extraction of anthocyanin. **b** Relative anthocyanin content at 18 days. The relative anthocyanin content was calculated as follows: absorbance (530 nm)/fresh weight (g). **c** Relative expression levels of anthocyanin biosynthesis-related genes in response to different NAA treatments at 18 days

the 35S::*MdARF13*-GFP construct. Additionally, the intact *MdIAA121* CDS was inserted into the pCAMBIA1301 vector containing the 35S promoter and a GFP tag sequence to prepare the 35S::*MdIAA121*-GFP construct. The resulting recombinant vectors were transferred to *Agrobacterium tumefaciens* LBA4404 cells. Two-week-old calli grown in liquid medium were cocultured with *A. tumefaciens* LBA4404 cells carrying the 35S::*MdARF13*-GFP construct on MS medium containing 2 $\mu\text{mol/L}$ NAA and 4 $\mu\text{mol/L}$ 6-benzylaminopurine at 25 °C for 2 days in darkness. The calli were then transferred to fresh MS medium supplemented with 662 $\mu\text{mol/L}$ carbenicillin and 74 $\mu\text{mol/L}$ kanamycin to screen for transgene-carrying calli. To obtain calli cotransfected with *MdARF13*+*MdIAA121*,

the same method was used, but calli were transferred to MS medium supplemented with 662 $\mu\text{mol/L}$ carbenicillin, 74 $\mu\text{mol/L}$ kanamycin, and 40 $\mu\text{mol/L}$ hygromycin. GFP-overexpressing calli were used as controls.

Yeast two-hybrid assay

Yeast (*Saccharomyces cerevisiae*) two-hybrid (Y2H) assays were conducted according to the manufacturer's instructions (Clontech, Mountain View, CA, USA). The *MdARF13* CDS was inserted into the pGKBT7 vector (i.e., bait), while the *MdIAA7*, *MdIAA121*, *MdMYB9*, and *MdMYB10* CDSs were inserted into the pGADT7 vector (i.e., prey). All recombinant vectors were cotransformed into yeast strain Y2H Gold cells using the lithium acetate method. The cells




Fig. 2 Bioinformatic analyses and *MdARF13* subcellular localization. **a** Phylogenetic analysis of ARF transcription factors in different species. **b** Alignment of *MdARF13*, *AtARF4*, *CiARF4*, *EgrARF4*, and *SiARF4* proteins. **c** Subcellular localization of the *MdARF13*-GFP fusion protein in transformed apple callus protoplasts. The protoplasts producing GFP alone were used as the control. Bars = 50 μ m

were cultured on synthetic defined (SD) medium lacking leucine and tryptophan (SD/-Leu/-Trp). Putative transformants were transferred to SD medium lacking adenine, histidine, leucine, and tryptophane (SD/-Ade/-His/-Leu/-Trp; Clontech) with or without X- α -gal.

Bimolecular fluorescence complementation assay

For a bimolecular fluorescence complementation (BiFC) assay, the *MdARF13* CDS was inserted into the pSPYNE-35S vector (with an *NYFP* tag sequence) to generate the *MdARF13*-*NYFP* recombinant vector. The *MdIAA121* and *MdMYB10* CDSs were inserted into the pSPYCE-35S vector (with a *CYFP* tag sequence) to generate the *MdIAA121*-*CYFP* and *MdMYB10*-*CYFP* recombinant vectors. *A. tumefaciens* LBA4404 cells were transformed with the recombinant vectors and then cultured in liquid medium until the optical density (600 nm) reached 0.6. Onion epidermal cells were treated with equal volumes of different combinations of *A. tumefaciens* strains. Onion cells were cultured at 23 °C for 48 h. The yellow fluorescence protein (YFP) signal was detected using a Zeiss CLSM-5 confocal laser scanning microscope (excitation wavelength of 488 nm).

Pull-down assay

The *MdIAA121* and *MdMYB10* open reading frames were cloned into the pET32a vector containing a poly-histidine


(HIS) tag sequence. The intact *MdARF13* CDS was ligated into the pGEX-4T-1 vector containing a glutathione *S*-transferase (GST) tag sequence. The recombinant vectors were inserted into *Escherichia coli* BL21 (DE3) cells (TransGen) to induce the production of fusion proteins. The resulting proteins were mixed together and then purified in columns using the HIS tag. The purified mixed proteins were subsequently analyzed by western blotting with anti-HIS or anti-GST antibodies (Abmart, Shanghai, China).

Yeast one-hybrid analysis

Yeast one-hybrid (Y1H) assays were conducted using yeast strain Y187 cells (Clontech) according to the manufacturer's instructions. *MdARF13* was cloned into the pGADT7 vector, while the *MdDFR* promoter was inserted into the pHIS2 vector. Different combinations of recombinant and empty vectors were cotransformed into yeast Y187 cells, and the interactions were examined on SD medium lacking Trp, Leu, and His (SD/-Trp/-Leu/-His) with an optimal concentration of 3-amino-1,2,4-triazole.

Electrophoretic mobility shift assays

Electrophoretic mobility shift assay (EMSA) experiments were completed using an EMSA kit (Pierce, Rockford, IL, USA) and biotin-labeled probes. Briefly, biotin-labeled probes were incubated in 1× binding buffer



(2.5% glycerol, 10 mM EDTA, 5 mM MgCl₂, and 50 mM KCl) with or without proteins at 24 °C for 25 min. An unlabeled probe was added to the reactions.

Chromatin immunoprecipitation–quantitative real-time PCR analysis

The chromatin immunoprecipitation (ChIP) experiment was completed using a modified version of a published method⁴⁴. The ChIP Assay Kit (Upstate Biotechnology, Lake Placid, NY, USA) and anti-GFP antibody (Abmart) were used for cross-linking, the removal of cross-linkers, immunoprecipitation, and elution. The ChIP signal was quantified by qPCR as the percentage of the total input DNA. The experiment was repeated three times.

Subcellular localization analysis

Protoplasts isolated from apple callus cells were prepared and transformed as described by Hu et al.⁴⁵. Transformed apple protoplasts were stained with the DNA-specific dye 4',6-diamidino-2-phenylindole, which highlights the location of the nucleus. The protoplasts from transgenic calli were observed by fluorescence microscopy.


In vitro protein degradation assay

Protein degradation assays were used to detect the dynamic changes to MdIAA13 in vitro. The assay buffer contained 25 mM Tris-HCl, pH 7.5, 10 mM NaCl, 10 mM MgCl₂, 5 mM dithiothreitol, 10 mM ATP, and 4 mM phenylmethylsulfonyl fluoride. Wild-type red-fleshed calli were treated with buffer, and the extract was incubated with the MdIAA13-His fusion protein for specific periods. The relative MdIAA121 abundance was detected by western blot using anti-His monoclonal antibodies.

Results

Anthocyanin content and related gene expression under different auxin concentrations

The anthocyanin contents of red-fleshed apple calli gradually decreased with increasing NAA concentrations (5–40 μmol/L), which caused the callus color to fade (Fig. 1a). Moreover, the anthocyanin content of calli cultured in 5 μmol/L NAA was 1.39-, 2.42-, and 5.98-fold higher than that of calli grown in 10, 20, and 40 μmol/L NAA, respectively (Fig. 1b). These results are consistent with the findings of Ji et al.^{26,27} and implied that low auxin



concentrations were conducive to anthocyanin accumulation in red-fleshed apple calli.

The changes in the anthocyanin content of calli grown under different auxin conditions were also reflected at the transcript level. As the auxin concentration increased, the expression levels of anthocyanin biosynthesis-related genes decreased by varying degrees (Fig. 1c). However, the transcript levels for the *Aux/IAA* and *ARF* genes increased with increasing NAA concentrations (Fig. 1c). In particular, the *MdARF13* expression levels in calli treated with 40 μmol/L NAA were 2–5-fold higher than in calli exposed to the other NAA concentrations. This finding suggested that *MdARF13* may repress anthocyanin biosynthesis.

Bioinformatics analysis of the nuclear protein *MdARF13*

Extensive bioinformatics and biochemical analyses of plants have clarified the number of *ARF* genes and their potential functions. In *A. thaliana*, experiments have revealed that AtARF4 contains an active RD and functions as a repressor in plant protoplast transfection assays³⁵. A phylogenetic analysis to elucidate the relationship between *MdARF13* and *ARF* proteins from other plant species indicated that *MdARF13* and AtARF4 belong to the same clade (Fig. 2a). Additionally, the aligned protein sequences revealed that *MdARF13* and *ARF* proteins from other plant species contain a conserved B3-like DBD, an RD, and a domain homologous to the *Aux/IAA* proteins (CTD) (Fig. 2b).

The subcellular localization of *MdARF13* was studied by introducing a 35S::*MdARF13-GFP* construct into the protoplasts of red-fleshed apple calli. The transgenic calli carrying the empty vector (35S::GFP) were used as the control. In protoplasts expressing 35S::*MdARF13-GFP*, the GFP signal was observed only in the nucleus, while the GFP signal was detected throughout the control protoplasts (Fig. 2c), implying that *MdARF13* is localized to the nucleus.

MdARF13 interacts with *MdIAA121*


The *ARFs* contain a dimerization domain responsible for the formation of heterodimers with *Aux/IAA* proteins³¹. The Y2H assays, which were used to test whether *MdARF13* interacts with *Aux/IAA* proteins, revealed that *MdARF13* interacts only with *MdIAA121* (Fig. 3a). The BiFC assay involving onion epidermal cells cotransformed with *MdARF13-NYFP* and *MdIAA121-CYFP* constructs confirmed that *MdARF13* interacts with *MdIAA121* (Fig. 3b). The nuclei of these onion epidermal cells produced a strong YFP signal according to laser scanning confocal microscopy. In contrast, the YFP signal was not detected in cells in which *MdIAA121-CYFP* was replaced with CYFP alone. Meanwhile, our pull-down assays revealed that the recombinant *MdIAA121-HIS* fusion protein could be purified with *MdARF13-GST*, but not with GST alone (Fig. 3c). These results were consistent with those of the Y2H and BiFC assays. Our data suggested that *MdARF13* can interact with *MdIAA121* under *in vivo* and *in vitro* conditions.

Auxin affects *MdIAA121* stability

Ubiquitination, which represents an important post-translational modification, is a key regulator of various processes, such as hormone signaling, responses to light and sugars, plant development, and plant immunity. In the *Aux/IAA-ARF* signaling pathway, auxin treatments promote the degradation of *Aux/IAA* by the 26S proteasome to release *ARF* proteins⁴⁶. We conducted protein degradation assays to study the posttranslational regulation of *MdIAA121* *in vitro*. The *MdIAA121-His* proteins were rapidly degraded after NAA was added (Fig. 4a). Furthermore, samples were treated with MG132 and dimethyl sulfoxide (DMSO). The MG132 treatment noticeably weakened the effect of auxin on *MdIAA121* stability (Fig. 4b). These findings indicated that auxin induces the degradation of *MdIAA121* via the 26S proteasome.

Overexpression of *MdARF13* repressed anthocyanin accumulation in the apple callus

To examine whether *MdARF13* affects anthocyanin accumulation, *MdARF13* was overexpressed in red-fleshed apple calli. The color of the resulting transgenic calli changed from dark red to yellow, while the transcript levels of anthocyanin biosynthesis-related genes and the accumulation of anthocyanin decreased (Fig. 5). These




observations implied that *MdARF13* represses anthocyanin biosynthesis. To assess whether *MdIAA121* inhibits *MdARF13* activities, we generated calli cotransfected with *MdARF13+MdIAA121* by *A. tumefaciens*-mediated transformation. The expression levels of anthocyanin structural genes and anthocyanin content were obviously higher in cotransfected calli than in calli overexpressing

MdARF13 (Fig. 5). These results demonstrated that *MdARF13* together with *MdIAA121* mediate auxin signaling to regulate anthocyanin biosynthesis.

MdARF13* interacts with *MdMYB10

The MYB TFs are crucial for anthocyanin biosynthesis, and the expression of the corresponding genes is regulated



by the external environment. In this study, we hypothesized that *MdARF13* is responsive to auxin and affects *MYB* expression. To verify this hypothesis, we completed Y2H assays to investigate whether *MdARF13* interacts with *MdMYB9* and *MdMYB10*. We observed that *MdARF13* interacts only with *MdMYB10* (Fig. 6a). The physical interaction between *MdARF13* and *MdMYB10* was further confirmed in pull-down and BiFC assays (Fig. 6b, c). Thus, we speculated that interactions between *MdARF13* and *MdMYB10* affect anthocyanin biosynthesis.


MdARF13 binds to the *MdDFR* promoter

ARF TFs regulate their downstream target genes by binding to the TGTCTC AuxRE in the promoter region and subsequently promoting or repressing expression²⁹. We analyzed the promoter elements of the anthocyanin biosynthesis structural genes (*MdCHS*, *MdCHI*, *MdDFR*, *MdLDOX*, *MdFLS*, and *MdUFGT*) using the PlantCARE *cis*-acting regulatory element database (<http://bioinformatics.psb.ugent.be/webtools/plantcare/html/>) and determined that only *MdDFR* contains a putative AuxRE in its promoter region (Fig. 7b). We conducted an EMSA to clarify whether *MdARF13* is recruited to the *MdDFR* promoter region. The assay results indicated that *MdARF13* could interact with DNA probes containing the TGTCTC AuxRE (Fig. 7b). Moreover, Y1H assays confirmed that *MdARF13* could interact with the *MdDFR* promoter (Fig. 7a), suggesting that *MdARF13* suppresses the expression of *MdDFR* to decrease

anthocyanin accumulation. Additionally, the ChIP-qPCR assay results suggested that the *MdARF13*-GFP fusion protein could bind to the *DFR* promoter, while the over-expression of *MdIAA121* in *MdARF13*-overexpressing calli clearly inhibited the recruitment of *MdARF13* to the *MdDFR* promoter (Fig. 7c).


Discussion

Auxin has long been recognized for its essential role in plant growth and development. Previous studies concluded that auxin signaling regulates anthocyanin biosynthesis in plant calli, which is associated with changes in the external auxin concentration^{20,26}. In separate studies, Ji et al.²⁶ and Liu et al.⁴⁷ observed that increasing auxin concentrations within a certain range can inhibit anthocyanin biosynthesis in apple and *A. thaliana*. In this study, we observed that increasing NAA concentrations increased the inhibition of anthocyanin accumulation. Deikman and Hammer⁴⁸ reported that the application of exogenous auxins suppressed the expression of anthocyanin regulatory and structural genes to varying degrees. Similarly, we observed that the expression levels of most structural and regulatory genes decreased in calli treated with NAA. Furthermore, the expression of genes encoding two types of TFs (Aux/IAA and ARF) associated with auxin signaling also increased to varying degrees, especially *MdARF13*. Thus, we speculate that these auxin signaling factors help regulate the expression of genes related to the anthocyanin biosynthesis pathway.



The Aux/IAA and ARF TFs are an integral part of the complex and elaborate auxin signaling pathway. Genome-wide analyses of ARF families have been completed in many plant species, including rice, tomato, and maize^{49–51}. In *A. thaliana*, in addition to ARF3, 13, 17, and 23, the ARF family mainly comprises proteins with the following three parts: an N-terminal DBD, a middle region that functions as an AD or RD, and a CTD²⁹. In this study, we determined that MdARF13 is homologous to AtARF4 and has a similar structure. ARFs exist in the nucleus for a relatively short period²⁹. Consistent with this previous report, we observed that MdARF13 is a nuclear

protein. Most Aux/IAAs function as repressors and are believed to dimerize with ARF activators via their CTDs. However, the Aux/IAA repressors are rapidly degraded as auxin concentrations increase, thereby impairing their inhibitory effect on early auxin-responsive genes³⁵. Our Y2H, BiFC, and pull-down assays revealed that MdARF13 can interact with MdIAA121. Additionally, exogenous auxin treatment induces the degradation of MdIAA121 by the 26S proteasome, suggesting that MdIAA121 might serve as a labile repressor of MdARF13. However, future studies will need to identify the specific ubiquitination mechanisms associated with MdIAA121 degradation.



To further investigate the functions of MdARF13 during anthocyanin biosynthesis, *MdARF13* was overexpressed in red-fleshed apple calli. A previous study involving transfected protoplasts indicated that AtARF4 functions as a transcriptional repressor³⁵. In the current study, qPCR data revealed that the transcript levels of auxin-regulated anthocyanin biosynthesis-related genes were down-regulated more in the calli with the 35S::*MdARF13-GFP* construct than in the wild-type control. Thus, MdARF13 is likely an inhibitor of anthocyanin biosynthesis in apple. Additionally, Aux/IAA can function as a repressor that dimerizes with the target ARF⁵². We observed that overexpressing *MdIAA121* in calli overexpressing *MdARF13* weakened the inhibitory effect of MdARF13 on anthocyanin biosynthesis, suggesting that MdARF13 may be targeted by MdIAA121. The interaction between these two proteins may attenuate the inhibitory effect of MdARF13 on anthocyanin biosynthesis.

The MYB TFs are crucial for regulating anthocyanin biosynthesis in apples. External environmental factors affect anthocyanin accumulation because the associated signaling elements regulate the expression of *MYB* genes. An earlier study indicated that the regulation of anthocyanin biosynthesis by *MdMYB1* is responsive to light, with COP1 functioning as a negative regulator that degrades *MdMYB1* in darkness⁵³. The HY5 TF targets the *MYBD* promoter in cytokinin-induced signaling pathways, leading to the accumulation of anthocyanin⁵⁴. In this study, Y2H, BiFC, and pull-down assays verified the *in vitro* and *in vivo* interaction between MdARF13 and *MdMYB10*, suggesting that MdARF13 might repress anthocyanin accumulation by down-regulating *MdMYB10* expression. Moreover, ARF TFs contain DNA-binding motifs and play key roles in the

regulation of downstream genes through TGTCTC sequence-specific interactions with promoters³⁰. We detected an MdARF13-binding site upstream of the *MdDFR* gene (Fig. 7a). EMSA, ChIP, and Y1H assays were conducted to confirm the specific binding of MdARF13 to the *MdDFR* promoter. We also proved that MdIAA121 functions as a repressor of the auxin signal and interacts with MdARF13 to inhibit the recruitment of MdARF13 to the *MdDFR* promoter. Therefore, in apple, exogenous auxins repress anthocyanin biosynthesis most likely through the MdIAA121–MdARF13 signal transduction pathway.

ARF TFs are important for auxin signaling, and are involved in many processes related to plant growth and development. In this study, we observed that the inhibition of anthocyanin biosynthesis by high auxin concentrations involves MdARF13 (Fig. 8). Clarifying the role of auxin in the regulation of anthocyanin biosynthesis by ARFs may provide new insights into the regulation of anthocyanin metabolism by other hormones. During the cultivation of fruit trees, the rational application of hormones based on different environmental conditions can improve the appearance and nutritional value of fruit crops. Herein, we discussed the involvement of ARFs in the anthocyanin biosynthesis pathway, which may have implications for the development of new cultivation techniques aimed at improving fruit coloration under diverse environmental conditions.

Acknowledgements

This work was supported by grants from the National Key Research Project (2016YFC0501505) and the National Natural Science Foundation of China (31572091 and 31730080).

Conflict of interest

The authors declare that they have no conflict of interest.

Publisher's note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Supplementary Information accompanies this paper at (<https://doi.org/10.1038/s41438-018-0068-4>).

Received: 17 March 2018 Revised: 7 June 2018 Accepted: 11 June 2018
Published online: 01 December 2018

References

- King, M. C. & Cliff, M. A. Development of a model for prediction of consumer liking from visual attributes of new and established apple cultivars. *J. Am. Pom. Soc.* **56**, 223–229 (2002).
- Horbowicz, M., Kosson, R., Grzesiuk, A. & Debski, H. Anthocyanins of fruits and vegetables—their occurrence, analysis and role in human nutrition. *Veg. Crops Res. Bull.* **68**, 5–22 (2008).
- Mol, J., Grotewold, E. & Koes, R. How genes paint flowers and seeds. *Trends Plant Sci.* **3**, 212–217 (1998).
- Knekt, P. et al. Dietary flavonoids and the risk of lung cancer and other malignant neoplasms. *Am. J. Epidemiol.* **146**, 223–230 (1997).

5. Middleton, J. E., Kandaswami, C. & Theoharides, T. C. The effects of plant flavonoids on mammalian cells: implications for inflammation, heart disease, and cancer. *Pharmacol. Rev.* **52**, 673–751 (2000).
6. Hollman, P. & Katan, M. Absorption, metabolism and health effects of dietary flavonoids in man. *Biomed. Pharmacother.* **51**, 305–310 (1997).
7. Zafra-Stone, S. et al. Berry anthocyanins as novel antioxidants in human health and disease prevention. *Mol. Nutr. Food Res.* **51**, 675–683 (2007).
8. Baudry, A. et al. TT2, TT8, and TTG1 synergistically specify the expression of BANYULS and proanthocyanidin biosynthesis in *Arabidopsis thaliana*. *Plant J.* **39**, 366–380 (2004).
9. Quattrochio, F. et al. PH4 of petunia is an R2R3 MYB protein that activates vacuolar acidification through interactions with basic-helix-loop-helix transcription factors of the anthocyanin pathway. *Plant Cell* **18**, 1274–1291 (2006).
10. Borevitz, J. O., Xia, Y., Blount, J., Dixon, R. A. & Lamb, C. Activation tagging identifies a conserved MYB regulator of phenylpropanoid biosynthesis. *Plant Cell* **12**, 2383–2394 (2000).
11. Cone, K. C., Coccilone, S. M., Burr, F. A. & Burr, B. Maize anthocyanin regulatory gene *pl* is a duplicate of *c1* that functions in the plant. *Plant Cell* **5**, 1795–1805 (1993).
12. Aharoni, A. et al. The strawberry FaMYB1 transcription factor suppresses anthocyanin and flavonol accumulation in transgenic tobacco. *Plant J.* **28**, 319–332 (2001).
13. Deluc, L. et al. The transcription factor VvMYB5b contributes to the regulation of anthocyanin and proanthocyanidin biosynthesis in developing grape berries. *Plant Physiol.* **147**, 2041–2053 (2008).
14. Mcghee, T. K. et al. An R2R3 MYB transcription factor associated with regulation of the anthocyanin biosynthetic pathway in rosaceae. *BMC Plant Biol.* **10**, 50 (2010).
15. Takos, A. M. et al. Light-induced expression of a MYB gene regulates anthocyanin biosynthesis in red apples. *Plant Physiol.* **142**, 1216–1232 (2006).
16. Ban, Y. et al. Isolation and functional analysis of a MYB transcription factor gene that is a key regulator for the development of red coloration in apple skin. *Plant Cell Physiol.* **48**, 958–970 (2007).
17. Espley, R. V. et al. Red colouration in apple fruit is due to the activity of the MYB transcription factor, MdMYB10. *Plant J.* **49**, 414–427 (2007).
18. Cominelli, E. et al. Expression analysis of anthocyanin regulatory genes in response to different light qualities in *Arabidopsis thaliana*. *J. Plant Physiol.* **165**, 886–894 (2008).
19. Rowan, D. D. et al. Environmental regulation of leaf colour in red 35SPAP1 *Arabidopsis thaliana*. *New Phytol.* **182**, 102–115 (2009).
20. Ozeki, Y. & Komamine, A. Effects of growth regulators on the induction of anthocyanin synthesis in carrot suspension cultures. *Plant Cell Physiol.* **27**, 1361–1368 (1986).
21. Qi, T. et al. The Jasmonate-ZIM-Domain proteins interact with the WD-Repeat/bHLH/MYB complexes to regulate jasmonate-mediated anthocyanin accumulation and trichome initiation in *Arabidopsis thaliana*. *Plant Cell* **23**, 1795–1814 (2011).
22. Meyer, H. J. & Staden, J. The in vitro production of an anthocyanin from callus cultures of *Oxalis linearis*. *Plant Cell Tissue Organ Cult.* **40**, 55–58 (1995).
23. Ozeki, Y. & Komamine, A. Effects of inoculum density, zeatin and sucrose on anthocyanin accumulation in a carrot suspension culture. *Plant Cell Tissue Organ Cult.* **5**, 45–53 (1985).
24. Zhou, L. L., Zeng, H. N., Shi, M. Z. & Xie, D. Y. Development of tobacco callus cultures over expressing Arabidopsis PAP1/MYB75 transcription factor and characterization of anthocyanin biosynthesis. *Planta* **229**, 37–51 (2008).
25. Murthy, H. N., Lee, E. J. & Paek, K. Y. Production of secondary metabolites from cell and organ cultures: strategies and approaches for biomass improvement and metabolite accumulation. *Plant Cell Tissue Organ Cult.* **118**, 1–16 (2004).
26. Ji, X. H. et al. Effect of auxin, cytokinin and nitrogen on anthocyanin biosynthesis in callus cultures of red-fleshed apple (*Malus sieversii* f. *niedzwetzkyana*). *Plant Cell Tissue Organ Cult.* **120**, 325–337 (2015).
27. Ji, X. H., Zhang, R., Wang, N., Yang, L. & Chen, X. S. Transcriptome profiling reveals auxin suppressed anthocyanin biosynthesis in red-fleshed apple callus (*Malus sieversii* f. *niedzwetzkyana*). *Plant Cell Tissue Organ Cult.* **123**, 389–404 (2015).
28. Guilfoyle, T. J., Ulmasov, T. & Hagen, G. The ARF family of transcription factors and their role in plant hormone-responsive transcription. *Cell Mol. Life Sci.* **54**, 619–627 (1998).
29. Ulmasov, T., Hagen, G. & Guilfoyle, T. J. Activation and repression of transcription by auxin-response factors. *Proc. Natl. Acad. Sci. USA* **96**, 5844–5849 (1999).
30. Tiwari, S. B., Wang, X. J., Hagen, G. & Guilfoyle, T. J. AUX/IAA proteins are active repressors, and their stability and activity are modulated by auxin. *Plant Cell* **13**, 2809–2822 (2001).
31. Guilfoyle, T. J. & Hagen, G. Auxin response factors. *J. Plant Growth Regul.* **10**, 281–291 (2001).
32. Liscum, E. & Reed, J. W. Genetics of Aux/IAA and ARF action in plant growth and development. *Plant Mol. Biol.* **49**, 387–400 (2002).
33. Benfey, P. N. Auxin action: slogging out of the swamp. *Curr. Biol.* **12**, R389–R390 (2002).
34. Hellmann, H. & Estelle, M. Plant development: regulation by protein degradation. *Science* **297**, 793–797 (2002).
35. Tiwari, S. B., Hagen, G. & Guilfoyle, T. The roles of auxin response factor domains in auxin-responsive transcription. *Plant Cell* **15**, 533–543 (2003).
36. Sessions, A. et al. ETTIN patterns the *Arabidopsis* floral meristem and reproductive organs. *Development* **124**, 4481–4491 (1997).
37. Pekker, I., Alvarez, J. P. & Eshed, Y. Auxin response factors mediate *Arabidopsis* organ asymmetry via modulation of KANADI activity. *Plant Cell* **17**, 2899–2910 (2005).
38. Hardtke, C. S. et al. Overlapping and nonredundant functions of the *Arabidopsis* auxin response factors MONOPTEROS and NONPHOTOTROPIC HYPOCOTYL 4. *Development* **131**, 1089–1100 (2004).
39. Hardtke, C. S. & Berleth, T. The *Arabidopsis* gene MONOPTEROS encodes a transcription factor mediating embryo axis formation and vascular development. *EMBO J.* **17**, 1405–1411 (1998).
40. Tian, C. E. et al. Disruption and overexpression of auxin response factor 8 gene of *Arabidopsis* affect hypocotyl elongation and root growth habit, indicating its possible involvement in auxin homeostasis in light condition. *Plant J.* **40**, 333–343 (2004).
41. Vriezen, W. H., Feron, R., Maretto, F., Keijman, J. & Mariani, C. Changes in tomato ovary transcriptome demonstrate complex hormonal regulation of fruit set. *New Phytol.* **177**, 60–76 (2008).
42. De, J. M., Woltersarts, M., Feron, R., Mariani, C. & Vriezen, W. H. The *Solanum lycopersicum* auxin response factor 7 (SIARF7) regulates auxin signaling during tomato fruit set and development. *Plant J.* **57**, 160–170 (2009).
43. Sun, J. et al. Effects of methyl jasmonate and abscisic acid on anthocyanin biosynthesis in callus cultures of red-fleshed apple (*Malus sieversii*, f. *niedzwetzkyana*). *Plant Cell Tissue Organ Cult.* **130**, 227–237 (2017).
44. An, X. H. et al. MdMYB9 and MdMYB11 are involved in the regulation of the JA-induced biosynthesis of anthocyanin and proanthocyanidin in apples. *Plant Cell Physiol.* **56**, 650–662 (2015).
45. Hu, D. G. et al. MdMYB1 regulates anthocyanin and malate accumulation by directly facilitating their transport into vacuoles in apples. *Plant Physiol.* **170**, 1315–1330 (2016).
46. Smallle, J. & Vierstra, R. D. The ubiquitin 26S proteasome proteolytic pathway. *Annu. Rev. Plant Biol.* **55**, 555–590 (2004).
47. Liu, Z., Shi, M. Z. & Xie, D. Y. Regulation of anthocyanin biosynthesis in *Arabidopsis thaliana* red pap1-d cells metabolically programmed by auxins. *Planta* **239**, 765–781 (2014).
48. Deikman, J. & Hammer, P. E. Induction of anthocyanin accumulation by cytokinins in *Arabidopsis thaliana*. *Plant Physiol.* **108**, 47–57 (1995).
49. Wang, D. et al. Genomewide analysis of the auxin response factors (ARF) gene family in rice (*Oryza sativa*). *Gene* **394**, 13–24 (2007).
50. Kumar, R., Tyagi, A. K. & Sharma, A. K. Genome-wide analysis of auxin response factor (ARF) gene family from tomato and analysis of their role in flower and fruit development. *Mol. Genet. Genomics* **285**, 245–260 (2011).
51. Xing, H. et al. Genome-wide identification and expression profiling of auxin response factor (ARF) gene family in maize. *BMC Genomics* **12**, 178 (2011).
52. Tiwari, S. B., Hagen, G. & Guilfoyle, T. J. Aux/iaa proteins contain a potent transcriptional repression domain. *Plant Cell* **16**, 533–543 (2004).
53. Li, Y. Y. et al. MdCop1 ubiquitin e3 ligases interact with mdmyb1 to regulate light-induced anthocyanin biosynthesis and red fruit coloration in apple. *Plant Physiol.* **160**, 1011–1022 (2012).
54. Nguyen, N. H. et al. MYBD employed by HY5 increases anthocyanin accumulation via repression of MYBL2 in *Arabidopsis*. *Plant J.* **84**, 1192–1205 (2015).