

ARTICLE

Open Access

# ABA-mediated regulation of leaf and root hydraulic conductance in tomato grown at elevated CO<sub>2</sub> is associated with altered gene expression of aquaporins

Liang Fang <sup>1,5</sup>, Lamis Osama Anwar Abdelhakim<sup>1</sup>, Josefina Nymark Hegelund<sup>1</sup>, Shenglan Li<sup>1</sup>, Jie Liu<sup>1,2</sup>, Xiaoying Peng<sup>1,3</sup>, Xiangnan Li<sup>4</sup>, Zhenhua Wei<sup>1,2</sup> and Fulai Liu<sup>1,2</sup>

## Abstract

Elevated CO<sub>2</sub> concentration in the air ( $e[\text{CO}_2]$ ) decreases stomatal density (SD) and stomatal conductance ( $g_s$ ) where abscisic acid (ABA) may play a role, yet the underlying mechanism remains largely elusive. We investigated the effects of  $e[\text{CO}_2]$  (800 ppm) on leaf gas exchange and water relations of two tomato (*Solanum lycopersicum*) genotypes, Ailsa Craig (WT) and its ABA-deficient mutant (*flacca*). Compared to plants grown at ambient CO<sub>2</sub> (400 ppm),  $e[\text{CO}_2]$  stimulated photosynthetic rate in both genotypes, while depressed the  $g_s$  only in WT. SD showed a similar response to  $e[\text{CO}_2]$  as  $g_s$ , although the change was not significant.  $e[\text{CO}_2]$  increased leaf and xylem ABA concentrations and xylem sap pH, where the increases were larger in WT than in *flacca*. Although leaf water potential was unaffected by CO<sub>2</sub> growth environment,  $e[\text{CO}_2]$  lowered osmotic potential, hence tended to increase turgor pressure particularly for WT.  $e[\text{CO}_2]$  reduced hydraulic conductance of leaf and root in WT but not in *flacca*, which was associated with downregulation of gene expression of aquaporins. It is concluded that ABA-mediated regulation of  $g_s$ , SD, and gene expression of aquaporins coordinates the whole-plant hydraulics of tomato grown at different CO<sub>2</sub> environments.

## Introduction

Stomata controls the photosynthesis ( $A_n$ ) and transpiration rates. The ability of plants to regulate the stomatal conductance ( $g_s$ ), through either modulating the aperture of the stomatal pore in a short term or changing the stomatal density (SD) in a long term, is crucial for their survival in an ever-changing environment. Among other environmental factors, the rising CO<sub>2</sub> concentration ([CO<sub>2</sub>]) in the atmosphere will have profound impacts on plant physiological processes, particularly those related to

stomatal control of leaf gas exchange and plant water relations<sup>1</sup>.

The influences of CO<sub>2</sub> elevation ( $e[\text{CO}_2]$ ) on stomatal morphology and physiology have been well documented<sup>1–5</sup>. Accumulated evidence showed that  $e[\text{CO}_2]$  reduces SD<sup>6–9</sup>. It has been suggested that reduction in SD caused by  $e[\text{CO}_2]$  could be modulated by abscisic acid (ABA) levels<sup>10,11</sup>. Earlier studies have shown that SD correlates positively with plant ABA level<sup>12–14</sup>. However, whether such a relationship also exists for plants grown in different CO<sub>2</sub> environments remains unknown. The low SD of plants grown at  $e[\text{CO}_2]$  could curtail the maximal  $g_s$  in a long term, while an immediate reduction of  $g_s$  after exposure to  $e[\text{CO}_2]$  has often been observed<sup>4,15</sup>. Guard cells could sense the change of [CO<sub>2</sub>] growth environment through responding to intercellular [CO<sub>2</sub>] (C<sub>i</sub>) and not leaf

Correspondence: Fulai Liu (fl@plen.ku.dk)

<sup>1</sup>Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Højbakkegaard Alle 13, 2630 Taastrup, Denmark

<sup>2</sup>Key Laboratory of Agricultural Soil and Water Engineering in Arid and Semiarid Areas, Ministry of Education, Northwest A&F University, 712100 Yangling, Shaanxi, China

Full list of author information is available at the end of the article.

surface  $[CO_2]$ <sup>16</sup>.  $e[CO_2]$  has been found to affect several ion channel activities, which may cause depolarization of the guard cell membrane potential<sup>4</sup>. In addition, ABA could play an important role in inducing stomatal closure in plants grown under  $e[CO_2]$ <sup>10,17</sup>. An earlier study showed that ABA could enhance the response of stomata to changes of  $[CO_2]$ <sup>18</sup>. More recently, literature revealed that  $e[CO_2]$ -caused closure of stomata might be mediated by ABA<sup>11</sup>. On the other hand, a recent study reported that  $e[CO_2]$ -induced stomatal closure is ABA independent via modulating OST1/SnRK2 kinases<sup>19</sup>. Therefore, the role of ABA in mediating  $g_s$  response to  $e[CO_2]$  merits further investigations. Moreover, it is well recognized that the distribution of ABA in plants is affected by the apoplast pH<sup>20</sup>, which could be affected by the  $CO_2$  growth environment hence modulating the efficiency of the ABA-mediated stomatal response to  $e[CO_2]$ . However, until now this aspect has not been explored.

Many researchers have reported that plants grown at  $e[CO_2]$  could maintain higher (less negative) leaf water potential ( $\Psi_l$ ), which could be partially attributed to the lowered  $g_s$  and hence transpiration rate at  $e[CO_2]$ <sup>5,21</sup>. Nevertheless, higher  $\Psi_l$  of plants grown at  $e[CO_2]$  was not always the case, even though  $g_s$  and transpiration rate were found to be lower, but hydraulic conductance could also be reduced in plants grown at  $e[CO_2]$ <sup>22,23</sup>, which may offset the positive effect of lowered  $g_s$  and transpiration rate on  $\Psi_l$ . Moreover, the response of plant hydraulic conductance to  $e[CO_2]$  was variable as controversial results were reported<sup>21</sup>. The changes of hydraulic conductance may be associated with changes of the abundance or activity of aquaporins that control plasma membrane water permeability<sup>24–26</sup>. Yet, it remains largely unknown whether  $e[CO_2]$  affects the expression of genes encoding aquaporins in leaf and root and whether endogenous ABA is involved in this process.


This study aimed to investigate the responses of leaf gas exchange, water relation characteristics, and hydraulic conductance of tomato plants to  $e[CO_2]$ . To achieve this, two tomato genotypes (GEs) differing in the endogenous ABA level were tested. We hypothesized that ABA would exert an important role in mediating the responses of stomatal behavior and plant water status to  $e[CO_2]$  by modulating both stomatal aperture and SD as well as the expression of aquaporins and thereby the whole-plant hydraulics and water balance.

## Results


The ABA-deficient *flacca* tomato had significantly small leaf area and shoot biomass in relation to the wild-type (WT) plants; although  $e[CO_2]$  tended to increase the growth for both of the GEs, the increments were not statistically significant (Fig. S1).

## Leaf gas exchange

Compared to WT, *flacca* had significantly higher  $A_n$  and  $g_s$  under both  $CO_2$  growth conditions (Fig. 1a, b). The  $A_n$  of both WT and *flacca* were significantly higher in the  $e[CO_2]$  plants than in the  $a[CO_2]$  plants. In relation to plants grown at  $a[CO_2]$ , a reduction of  $g_s$  at  $e[CO_2]$  was only noticed in WT and not in *flacca*.



**Fig. 1 Leaf gas exchange and stomatal morphology response of the two tomato genotypes to different  $CO_2$  growth environments.** Net photosynthetic rate ( $A_n$ ) (a), stomatal conductance ( $g_s$ ) (b), and stomatal density (SD) (c) of wild-type tomato "Ailsa Craig" (WT) and its respective ABA-deficient mutant (*flacca*) grown at ambient (400 ppm,  $a[CO_2]$ ) and elevated (800 ppm,  $e[CO_2]$ ) atmospheric  $CO_2$  concentrations. The effects of  $CO_2$  growth environment ( $CO_2$ ) and genotype (GE) as well as their interactions  $CO_2 \times GE$  are presented (two-way ANOVA). The different letters on the columns indicate statistically significant difference between the treatments by Tukey's test at  $P < 0.05$ . Error bars indicate standard error of the means ( $n = 8$ )



**Fig. 2 Leaf and xylem sap ABA concentration and xylem sap pH of the two genotypes of tomato as affected by different CO<sub>2</sub> growth environments.** Leaf ABA concentration ([ABA]<sub>leaf</sub>) (a), xylem sap ABA concentration ([ABA]<sub>xylem</sub>) (b), and xylem pH (c) of wild-type tomato "Ailsa Craig" (WT) and its respective ABA-deficient mutant (*flacca*) grown under ambient (400 ppm, a[CO<sub>2</sub>]) and elevated (800 ppm, e[CO<sub>2</sub>]) CO<sub>2</sub> environments. The effects of CO<sub>2</sub> growth environment (CO<sub>2</sub>) and genotype (GE) as well as their interactions CO<sub>2</sub> × GE are presented (two-way ANOVA). The different letters on the columns indicate statistically significant difference between the treatments by Tukey's test at P < 0.05. Error bars indicate standard error of the means (SE) (n = 8)

### Stomatal density

Significantly higher SD in *flacca* than in WT was noticed across the two CO<sub>2</sub> growth environments. Compared to the a[CO<sub>2</sub>] plants, SD tended to be lower when grown at e[CO<sub>2</sub>] for WT (although not statistically significant), whereas for *flacca* a slight increase of SD was noticed in plants grown at e[CO<sub>2</sub>], resulting in a significant interaction between CO<sub>2</sub> and GE (Fig. 1c).

### Leaf and xylem sap ABA concentration

As expected, significantly higher leaf and xylem ABA concentrations were observed in WT compared to *flacca* (Fig. 2a, b). In relation to the a[CO<sub>2</sub>] plants, e[CO<sub>2</sub>] significantly increased [ABA]<sub>leaf</sub> and [ABA]<sub>xylem</sub>, while the magnitude of increase was greater in WT than in *flacca*, although no significant CO<sub>2</sub> × GE effect was found.

### Xylem sap pH

The e[CO<sub>2</sub>] plants had higher xylem pH than the a[CO<sub>2</sub>] plants; and in general *flacca* had higher xylem pH than WT irrespective to the CO<sub>2</sub> growth environments (Fig. 2c).

For WT, g<sub>s</sub> was negatively correlated with [ABA]<sub>leaf</sub> across the two CO<sub>2</sub> growth environments (P < 0.001); although a similar relationship was also noticed in *flacca*, the linear regression was not statistically significant (Fig. 3a). Likewise, negative linear relationships between [ABA]<sub>xylem</sub> and g<sub>s</sub> was observed across the two CO<sub>2</sub> growth environments for both GEs; the linear regressions, however, were not statistically significant (Fig. 3b). No obvious relationship between xylem pH and g<sub>s</sub> were evident (Fig. 3c).

### Plant water relations


*flacca* had lower (more negative)  $\Psi_l$  and  $\Psi_n$  and lower  $\Psi_p$  compared to WT (Fig. 4). CO<sub>2</sub> growth environment had no effect on  $\Psi_l$ , while e[CO<sub>2</sub>] decreased  $\Psi_n$  as compared to a[CO<sub>2</sub>] (Fig. 4b). e[CO<sub>2</sub>] increased the  $\Psi_p$  of WT but not of *flacca* (Fig. 4c).

### Hydraulic conductance

Compared to the a[CO<sub>2</sub>] plants, lower K<sub>l</sub> when grown at e[CO<sub>2</sub>] was observed (Fig. 5a); however, the reduction was less significant in *flacca* than in WT resulting in a significant interaction between CO<sub>2</sub> and GE. The K<sub>r</sub> of WT was significantly higher than that of *flacca* when grown at a[CO<sub>2</sub>], whereas they had a similar K<sub>r</sub> when grown at e[CO<sub>2</sub>] (Fig. 4b). e[CO<sub>2</sub>] decreased K<sub>r</sub> only in WT while it slightly increased K<sub>r</sub> in *flacca* in relation to the plants grown at a[CO<sub>2</sub>] (Fig. 5b).

### Expression of genes encoding aquaporins of the plasma membrane intrinsic protein (PIP) subgroup


In leaves of WT, transcripts of four PIPs (PIP1.5, PIP2.1, PIP2.8, and PIP2.9) responded to e[CO<sub>2</sub>] with a 2–5-fold downregulation of expression levels (Fig. 6a). PIP1.3 and PIP2.4 showed similar trends but were not significant or below the twofold change cut-off. In *flacca*, PIPs showed only minor fluctuations in transcript levels none of which were significant when comparing a[CO<sub>2</sub>] to e[CO<sub>2</sub>] growth conditions. When comparing leaf PIP expression between the two GEs grown at a[CO<sub>2</sub>], WT showed



**Fig. 3** Correlations between stomatal conductance to leaf and xylem ABA concentration and xylem sap pH of the two genotypes of tomato grown at different  $\text{CO}_2$  levels. Correlations of stomatal conductance ( $g_s$ ) to leaf ABA concentration ( $[\text{ABA}]_{\text{leaf}}$ ) (a) and xylem sap ABA concentration ( $[\text{ABA}]_{\text{xylem}}$ ) (b), and xylem pH (c) of wild-type tomato "Ailsa Craig" (WT) and its ABA-deficient mutant (flacca) grown under ambient (400 ppm, a[ $\text{CO}_2$ ]) and elevated (800 ppm, e[ $\text{CO}_2$ ])  $\text{CO}_2$  environments. Triple asterisks (\*\*\*\*) indicates that the regression line is statistically significant ( $P < 0.001$ ) and ns denotes no significance


significantly higher *PIP2.1*, *PIP2.4* and *PIP2.9* expression than *flacca*.

In roots of WT, PIP transcript responses to e[ $\text{CO}_2$ ] showed a similar response as in leaves. All 8 root PIPs



**Fig. 4** Leaf water relation characteristics of the two tomato genotypes as affected by different  $\text{CO}_2$  growth environments. Leaf water potential ( $\psi_l$ ) (a), osmotic potential ( $\psi_\pi$ ) (b), and turgor pressure ( $\psi_p$ ) (c) of wild-type tomato "Ailsa Craig" (WT) and its ABA-deficient mutant (flacca) grown under ambient (400 ppm, a[ $\text{CO}_2$ ]) and elevated (800 ppm, e[ $\text{CO}_2$ ])  $\text{CO}_2$  environments. The effects of  $\text{CO}_2$  growth environment ( $\text{CO}_2$ ) and genotype (GE) as well as their interactions  $\text{CO}_2 \times \text{GE}$  are presented (two-way ANOVA). The different letters on the columns indicate significant difference between the treatments by Tukey's test at  $P < 0.05$ . Error bars indicate standard error of the means ( $n = 8$ )

showed 2–4-fold downregulation in response to e[ $\text{CO}_2$ ]; however, just transcriptional changes in 5 PIPs were found to be significant (*PIP1.3*, *PIP2.1*, *PIP2.4*, *PIP2.5*, and *PIP2.8*) (Fig. 6b). As in leaves, PIPs of *flacca* did not follow the clear response observed in WT. Five *flacca* root PIPs did not respond in transcript abundance to the e[ $\text{CO}_2$ ] growth environment. However, three PIP transcripts responded with significant twofold upregulation (*PIP2.8* and *PIP2.9*) or downregulation (*PIP2.4*) in *flacca* grown at




**Fig. 5 Leaf and root hydraulic conductance of the two tomato genotypes as affected by different  $\text{CO}_2$  growth environments.**

Leaf hydraulic conductance ( $K_l$ ) (a) and root hydraulic conductance ( $K_r$ ) (b) of wild-type tomato "Ailsa Craig" (WT) and its representative ABA-deficient mutant (flacca) grown under ambient (400 ppm,  $a[\text{CO}_2]$ ) and elevated (800 ppm,  $e[\text{CO}_2]$ )  $\text{CO}_2$  environments. The effects of  $\text{CO}_2$  growth environment ( $\text{CO}_2$ ) and genotype (GE) as well as their interactions  $\text{CO}_2 \times \text{GE}$  are presented (two-way ANOVA). The different letters on the columns indicate significant difference between the treatments by Tukey's test at  $P < 0.05$ . Error bars indicate standard error of the means (SE) ( $n = 4$ )

$e[\text{CO}_2]$  in relation to that grown at  $a[\text{CO}_2]$ . Furthermore, at  $a[\text{CO}_2]$   $\text{PIP}1.3$ ,  $\text{PIP}2.4$ ,  $\text{PIP}2.8$ , and  $\text{PIP}2.9$  had significantly lower expression level while  $\text{PIP}1.5$  had significantly higher expression level, respectively, in flacca than in WT (Fig. 6b).

## Discussion

It is well known that  $e[\text{CO}_2]$  enhances  $A_n$  while reducing  $g_s$ , although the response may vary among species and different growth environments<sup>4,27</sup>. Consistent with this, here  $A_n$  was stimulated by  $e[\text{CO}_2]$  in both WT and flacca plants (Fig. 1a); however, reduction of  $g_s$  by  $e[\text{CO}_2]$  was only observed in the WT and not in flacca (Fig. 1b). As expected, at both  $\text{CO}_2$  growth environments, WT plants possessed significantly greater  $[\text{ABA}]_{\text{leaf}}$  and  $[\text{ABA}]_{\text{xylem}}$  than flacca (Fig. 2a, b); also,  $e[\text{CO}_2]$  increased  $[\text{ABA}]_{\text{leaf}}$  and  $[\text{ABA}]_{\text{xylem}}$  more pronounced in WT than in flacca. For WT,  $g_s$  was negatively correlated with  $[\text{ABA}]_{\text{leaf}}$  (Fig. 3a), revealing that  $g_s$  was most probably



**Fig. 6 Effect of different  $\text{CO}_2$  growth environments on the relative gene expression of aquaporins in leaf and root of the two tomato genotypes.** Relative expression of genes encoding the aquaporin subfamily of plasma membrane intrinsic proteins (PIPs) in leaf (a) and root (b) of wild-type tomato "Ailsa Craig" (WT) and its representative ABA-deficient mutant (flacca) grown under ambient (400 ppm,  $a[\text{CO}_2]$ ) and elevated (800 ppm,  $e[\text{CO}_2]$ )  $\text{CO}_2$  environments. Different letters on the top of the columns for each PIP gene indicate significant difference between the treatments by Tukey's test at  $P < 0.05$ . Error bars indicate standard error of the means (SE) ( $n = 4$ )

controlled by  $[\text{ABA}]_{\text{leaf}}$  across the two  $\text{CO}_2$  growth environments. Such relationship, however, was not evident for  $[\text{ABA}]_{\text{xylem}}$  and  $g_s$ , although earlier studies have frequently reported that  $g_s$  correlated better with  $[\text{ABA}]_{\text{xylem}}$  than with  $[\text{ABA}]_{\text{leaf}}$ <sup>28</sup>. Besides, for the two GEs the change of  $g_s$  in response to  $e[\text{CO}_2]$  was associated with a similar pattern of change in SD (Fig. 1c), suggesting that the endogenous ABA level exerted an important role in the  $e[\text{CO}_2]$ -induced modulation of SD and thus  $g_s$ . Therefore, it is reasonable to postulate that the endogenous ABA level had influenced the responsiveness of SD and  $g_s$  to  $e[\text{CO}_2]$  in tomato plants. In line with this, it has been reported that  $e[\text{CO}_2]$ -induced stomatal closure and reductions in SD was modulated by plant ABA levels<sup>10,11</sup>. However, a positive correlation between SD and plant ABA level previously reported in other studies is contradictory to results obtained here<sup>12–14</sup>. Moreover, in addition to ABA, cytokinins and other phytohormones could have also been involved in stomatal regulation in plants grown at  $e[\text{CO}_2]$ <sup>29</sup>.

An earlier study suggested that the higher  $[\text{ABA}]_{\text{leaf}}$  in the  $e[\text{CO}_2]$  plants might be caused by slight osmotic stress

due to the relative higher solutes' accumulation induced by rising  $A_n$  when plants grow at  $e[\text{CO}_2]$ <sup>30</sup>. This was seemingly true here as  $e[\text{CO}_2]$  led to more negative  $\Psi_n$  in all plants (Fig. 4b). However, ABA synthesis in leaf is believed to be linked with  $\Psi_p$ <sup>31</sup>, and an increased  $\Psi_p$  in plants grown at  $e[\text{CO}_2]$  (Fig. 3c) would result in a low  $[\text{ABA}]_{\text{leaf}}$  disagreeing with the results of the present study. Recently, evidence has indicated that ABA accumulation in drying leaves is due to a decrease in cell volume, not due to reduction of  $\Psi_p$ <sup>32</sup>. Moreover, the greater  $[\text{ABA}]_{\text{xylem}}$  of the  $e[\text{CO}_2]$  plants could be linked to their higher xylem sap pH in relation to the  $a[\text{CO}_2]$  plants (Fig. 2b, c)<sup>33</sup>. Besides, the lowered root hydraulic conductance at  $e[\text{CO}_2]$  could also contribute to the greater  $[\text{ABA}]_{\text{xylem}}$  in the  $e[\text{CO}_2]$  plants<sup>22</sup>, assumingly attributed to a reduced rate of sap flow during collection, which may cause a concentration effect on the xylem sap. Our results disagree with Li et al., who reported that  $e[\text{CO}_2]$  did not affect  $[\text{ABA}]_{\text{leaf}}$  in tomato plants<sup>34</sup>, and the reasons behind this disagreement are unknown, which merit further studies.

In literature, very little information is available about how  $e[\text{CO}_2]$  influences xylem sap pH. In this study, higher xylem sap pH was observed in the  $e[\text{CO}_2]$  plants compared to the  $a[\text{CO}_2]$  plants (Fig. 2c), suggesting that xylem sap pH was affected by  $[\text{CO}_2]$ . This is a novel finding, although the mechanisms behind remain speculative. One mechanism could be due to the bicarbonate ion ( $\text{HCO}_3^-$ ), which is produced when  $\text{CO}_2$  dissolve in xylem sap that modulates the pH. Another mechanism might be linked to a disturbed root ion (e.g., nitrate) uptake caused by  $e[\text{CO}_2]$ <sup>35</sup>; a reduced nitrate uptake under  $e[\text{CO}_2]$  would result in an increase of xylem pH as suggested by a previous study<sup>36</sup>. Interestingly, compared to WT plants, *flacca* had greater xylem sap pH (Fig. 2c); this contradicts the common consensus that a high xylem sap pH would enable more efficient stomatal closure<sup>33</sup>, yet the reasons behind this are unknown. As mentioned previously, an increased xylem sap pH could retain ABA in the apoplast thereby more efficiently inducing stomatal closure<sup>20,33,37</sup>. Here, in addition to the contribution of a slightly lowered SD, the  $e[\text{CO}_2]$ -induced reduction in  $g_s$  in the WT could be partially ascribed to the higher  $[\text{ABA}]_{\text{leaf}}$  and/or  $[\text{ABA}]_{\text{xylem}}$  as well as a greater xylem sap pH.

Accumulated evidence indicates that changes in  $g_s$  could lead to changes in  $\Psi_l$  by altering the transpiration rate in plants under well-watered conditions<sup>38</sup>. In the present study, the greater  $g_s$  of *flacca* could have resulted in lower  $\Psi_l$ , and vice versa for the WT plants (Fig. 3a), consistent with previous findings in the same GE<sup>39</sup>. Early studies have indicated that  $e[\text{CO}_2]$  could lead to a higher  $\Psi_l$  in plants<sup>5,23</sup>. In agreement with this, the  $\Psi_l$  of WT plants was slightly higher (less negative) under  $e[\text{CO}_2]$  than at  $a[\text{CO}_2]$ , though the overall  $e[\text{CO}_2]$  effect on  $\Psi_l$  was

not statistically significant (Fig. 4a). In addition,  $e[\text{CO}_2]$  decreased  $\Psi_n$  in all plants affirming our earlier findings in tomato<sup>5</sup>; while *flacca* had significantly lower  $\Psi_n$  than WT under both  $\text{CO}_2$  growth environments (Fig. 3b), which could be a result of enhanced solutes' accumulation caused by the greater photosynthetic rate ( $A_n$ ) in those plants (Fig. 1a). Also, a higher  $[\text{ABA}]_{\text{leaf}}$  might induce greater vacuolar invertase activity in the leaf, which could enhance hexose concentrations thereby contributing to a lowered  $\Psi_n$ <sup>40</sup>. The significantly greater  $\Psi_p$  in the  $e[\text{CO}_2]$  plants was most likely a consequence of the lowered  $\Psi_n$  as the  $\Psi_l$  was almost unaffected by  $\text{CO}_2$  growth conditions. Further, it was noticed that the  $\Psi_p$  of *flacca* was much lower than that of WT tomato at both  $\text{CO}_2$  growth conditions (Fig. 4c) and that could be attributed to the relatively greater dehydration of the leaf caused by the greater  $g_s$  in *flacca*.

Several early studies have demonstrated that plant hydraulic conductance was reduced when grown at  $e[\text{CO}_2]$ <sup>22,41</sup>. In line with this, here the  $e[\text{CO}_2]$  plants possessed significantly lower  $K_l$  and  $K_r$  in WT (Fig. 4a, b). The change of hydraulic conductance of WT plants grown at  $e[\text{CO}_2]$  was closely associated with the change of  $g_s$ , indicating that the reduction in hydraulic conductance could be due to a homeostatic adjustment by the plants in order to match hydraulic conductance with the lowered  $g_s$  at  $e[\text{CO}_2]$ <sup>42</sup>. However, this was not the case in *flacca*, where the  $K_l$  and  $K_r$  were almost identical at both  $\text{CO}_2$  growth environments (even a slight increase of  $K_r$  of the  $e[\text{CO}_2]$  plants as compared to the  $a[\text{CO}_2]$  plants) (Fig. 5a, b). Moreover, in the present study, *flacca* had lower  $K_l$  and  $K_r$  compared to WT under  $a[\text{CO}_2]$  (Fig. 4a, b). This was in agreement with earlier findings that a higher endogenous ABA level linked to a greater hydraulic conductance<sup>43–45</sup>. Recently, a study also reported that in barley the ABA-deficit mutant possessed significantly lower hydraulic conductance as compared with the WT<sup>46</sup>. These authors suggested that high ABA level and hence greater aquaporin abundance and higher hydraulic conductivity seem essential to sustain the  $\Psi_l$  in barley plants. Nonetheless, although the endogenous ABA level was greater in WT plants grown at  $e[\text{CO}_2]$  than at  $a[\text{CO}_2]$  (Fig. 2a, b), the hydraulic conductance was lower in those plants (Fig. 5a, b), indicating that, beside endogenous ABA, other factors might also be involved in the modulation of plant hydraulic conductance under  $e[\text{CO}_2]$ .

To explore the mechanisms underlying the  $e[\text{CO}_2]$ -induced changes in leaf and root hydraulic conductance, the expression of gene encoding major PIP aquaporins were investigated. To date, there is no information available about how  $e[\text{CO}_2]$  affects the gene expression of aquaporins in tomato plants. A study<sup>26</sup> suggested that the changes in aquaporins expression could be regulated by  $\text{CO}_2$ , which might contribute to the changes of hydraulic

conductance in soybean plants, but there was no direct evidence given in the paper. Here in WT plants, genes encoding five out of six and eight PIPs in leaf and root, respectively, were constantly and significantly downregulated by growing at  $e[\text{CO}_2]$  (Fig. 6). Consistent with this, a study in broccoli (*Brassica oleracea* L. var Italica) showed that  $e[\text{CO}_2]$  decreased the abundance of PIP1 and PIP2 protein in both leaf and root as compared to  $a[\text{CO}_2]$ <sup>47</sup>. Similarly, in tobacco (*Nicotiana tabacum*) leaves a downregulation of *NtPIP2;1* gene expression was noticed when grown at  $e[\text{CO}_2]$ <sup>48</sup>. However, this was not the case for *flacca* where most of the genes were unaffected or even upregulated by  $e[\text{CO}_2]$ , revealing that the endogenous ABA level exerts a crucial role in mediating the response of aquaporins to  $e[\text{CO}_2]$ . In line with this, several earlier studies demonstrated that ABA is involved in modulating gene expression of PIPs<sup>44,49</sup>. For instance, a study showed that PIPs were upregulated in response to elevated ABA level in *Arabidopsis thaliana*<sup>49</sup>; likewise, another study reported that PIPs were downregulated in response to low endogenous ABA level in transgenic maize plants with silenced ABA synthesis<sup>44</sup>. Most interestingly, the changes of aquaporin gene expression coincided well with the changes in  $K_l$  and  $K_r$ , indicating that modulation of the gene expression of aquaporins in the leaf and root contributed essentially to the changes of hydraulic conductance in the  $e[\text{CO}_2]$  plants. The mechanisms underlying such root and shoot coordination in controlling water balance via modulating PIP expression of plants grown at  $e[\text{CO}_2]$  remain unknown; the modified N nutrition could be involved as suggested by a recent study<sup>50</sup>. Nonetheless, this finding is of great significance for improving our understanding about the responses of tomato plants to  $e[\text{CO}_2]$  and the role of ABA in mediating these responses.

Taken together, the results of this study reveal that endogenous ABA is involved in modulating the physiological responses of tomato plants to  $e[\text{CO}_2]$ . ABA-mediated regulation of  $g_s$  and  $K_l$  and  $K_r$  coordinates the whole-plant hydraulics and water balance of tomato plants under different  $\text{CO}_2$  growth environments.

## Materials and methods

### Plant material and growth conditions

Seeds of isogenic WT (cv. Ailsa Craig) tomato and an ABA-deficient tomato mutant (*flacca*) (*Solanum lycopersicum*) were provided by the Lancaster Environment Centre (Lancaster University, UK). The *flacca* is impaired in the oxidation of ABA-aldehyde to ABA thus possessing significantly lower (ca. 20-folds less) endogenous ABA concentrations than WT<sup>51,52</sup>. All potted plants were grown in a climate-controlled greenhouse at Taastrup campus of University of Copenhagen, Denmark (55°67' N, 12°30' E). The seeds were sown in 4 L pots filled with

2,600 g of peat material (Plugg-och Såjord-Dry matter ca.110 kg m<sup>-3</sup>, organic matter >95%, pH 5.5–6.5 and EC 1.5–2.5 mS cm<sup>-1</sup>) on February 7, 2018. In total, 32 pots were established. Four weeks after sowing, fertilizers were added together with irrigation water in the form of  $\text{NH}_4\text{NO}_3$  (2.8 g) and  $\text{H}_2\text{KPO}_4$  (3.5 g) per pot to avoid any nutrient deficiency.

After sowing, the plants were grown in two separated greenhouse cells (cell 1 and cell 2) with different atmospheric  $\text{CO}_2$  concentrations: ambient (400 ppm,  $a[\text{CO}_2]$ ) and elevated (800 ppm,  $e[\text{CO}_2]$ ), respectively. In each cell, 16 plants (8 WT and 8 *flacca*) were randomly distributed on a growth table. The  $\text{CO}_2$  was enriched inside the cell by emission of pure  $\text{CO}_2$  at one point from a bottle tank and distributed through the ventilation system. The  $[\text{CO}_2]$  was monitored every 6 s by a  $\text{CO}_2$  Transmitter (Series GMT220, Vaisala Group, Helsinki, Finland).

The day/night air temperature in the both greenhouse cells were set at 20/18 ± 2 °C, relative humidity at 60 ± 2%, photoperiod at 16 h, and photosynthetic active radiation (PAR) at >250  $\mu\text{mol m}^{-2} \text{s}^{-1}$  supplied by sunlight plus LDE lamps. The vapor pressure deficit ranged from 0.8 to 1 kPa. The climate data were monitored every 5 min and recorded by a climate computer. The daily average  $[\text{CO}_2]$ , air temperature, and relative humidity in the greenhouse cells during the experiment period are shown in Fig. 7. All pots were well watered to 95% pot water holding capacity after seedling establishment.


### Measurements

#### Leaf gas exchange

Six weeks after sowing, net photosynthetic rate ( $A_n$ ), stomatal conductance ( $g_s$ ), and transpiration rate ( $T_r$ ) were measured on upper canopy fully expanded leaves (one leaflet per plant, eight WT and eight *flacca* plants per cell, respectively) between 9:00 to 12:00 using a portable photosynthetic system (LiCor-6400XT, LI-Cor, NE, USA). Measurements were done at 20 °C chamber temperature and 1200  $\text{mol m}^{-2} \text{s}^{-1}$  PAR, and 400 ppm in cuvette for  $a[\text{CO}_2]$  and 800 ppm in cuvette for  $e[\text{CO}_2]$  growth environment, respectively.

#### Stomatal density

SSD was measured using a digital microscope (Dino lite AM4113/AD4113 series with ver. 1.4.1, Vidy Precision Equipment Co. Ltd, Wuxi, China). For each plant, three images (calibrated image size: 654 × 490  $\mu\text{m}$ ) from both the adaxial and abaxial leaf surfaces were taken (one leaflet per plant, eight WT and eight *flacca* plants per cell, respectively). ImageJ software (Version 1.51k, Wayne Rasband, National Institutes of Health, USA, Java 1.6.0–24 (64 bit)) was used for counting the stomatal number.



**Fig. 7** The daily average of atmospheric CO<sub>2</sub> concentration [CO<sub>2</sub>], air temperature, and relative humidity in the two greenhouse cells during the experimental period. Error bars indicate standard error of the mean (SE = 96)

#### Plant water relations

Midday leaf water potential ( $\Psi_l$ ) was measured on young fully expanded leaf (one leaflet per plant, eight WT and eight *flacca* plants per cell, respectively) using a scholander-type pressure chamber (Soil Moisture Equipment Corp., Santa Barbara, CA, USA). After measuring  $\Psi_l$ , the leaf was immediately cut into two pieces and packed in aluminum foil separately and frozen in liquid nitrogen for later determination of leaf osmotic potential ( $\Psi_\pi$ ) and leaf ABA concentration ( $[\text{ABA}]_{\text{leaf}}$ ).  $\Psi_\pi$  was

measured using a psychrometer (C-52 sample chamber, Wescor Crop, Logan, UT, USA) connected to a microvoltmeter (HR-33T, Wescor, Logan, UT, USA) at 22 ± 1 °C. Turgor pressure ( $\Psi_p$ ) was calculated as  $\Psi_l - \Psi_\pi$ .

Leaf hydraulic conductance ( $K_l$ , mmol m<sup>-2</sup> s<sup>-1</sup> MPa<sup>-1</sup>) was calculated as:

$$K_l = \frac{T_r}{\Psi_l} \quad (1)$$

where  $T_r$  is the transpiration rate and  $\Psi_l$  is the leaf water potential.

Root water potential was measured on four WT and four *flacca* plants, respectively, in each greenhouse cell with a scholander-type pressure chamber (AGRSCI, KVL, Denmark). The whole pots were put into the chamber, then the chamber was sealed and only the above-soil part of the plants was left out. The stem was cut with a scalpel at approximate 10 cm above the soil surface. By pressuring the whole root system, the  $\Psi_r$  was determined when the xylem sap started to appear from the cutting surface. And the pressure was increased until it equaled  $\Psi_l$  of the plant to ensure a sap flow rate similar to the transpiration rate of the plant. Approximately 0.5–1 ml of xylem sap was collected to Eppendorf tubes using a pipette. Immediately after collection, the xylem sap was weighed and then frozen in liquid nitrogen and stored at -80 °C for ABA analysis. The time for collecting the sap was recorded and the stem cross-section area was measured. Then the hydraulic conductance of the whole root system ( $K_r$ , g cm<sup>-2</sup> min<sup>-1</sup> MPa<sup>-1</sup>) was calculated as:

$$K_r = \frac{\text{Xylem mass}}{T \times P \times S} \quad (2)$$

where xylem mass is the weight of the collected xylem sap (g);  $T$  is the collection time (s);  $P$  is the chamber pressure (MPa), which was maintained during collection; and  $S$  is the stem cross-section area (cm<sup>2</sup>).

Plant leaf area was determined by a leaf area meter (LICOR 3100, LI-COR Inc., Lincoln, NB) and the shoot biomass was determined after oven-drying at 70 °C for 48 h.

#### Xylem sap pH

After thawing for 30 min, the pH of the xylem sap was determined with a microelectrode (model PHR-146, Lazar Research Laboratories, Inc., CA, USA) interfaced with a pH meter (Model 60, Jenco Instruments Inc., CA, USA).

#### Leaf and xylem sap ABA concentration

Enzyme-linked immunosorbent assay was used to determine ABA concentration in the leaf and xylem sap samples following the protocol of Asch<sup>53</sup>. For the leaf ABA assay, we used the same leaf samples for determining  $\Psi_l$ , which could have caused dehydration of the leaf thus

affecting leaf ABA concentration. To clarify this, an extra test was done where ABA concentration of leaf samples from the same plants with and without  $\gamma_1$  measurements was compared, and no differences in ABA concentration were found between the two groups of leaves. Therefore, our method is valid for evaluating the leaf ABA concentration under the different treatments.

#### DNA/RNA extractions, cDNA synthesis, and PCR reactions

DNA and RNA extractions were done from 80 to 100 mg grinded leaf or root material using the DNeasy Plant Mini Kit or the RNeasy Plant Mini Kit, respectively, as recommended by the supplier (Qiagen, Germany). DNA or RNA yield and purity were estimated using Nanodrop<sup>TM</sup> 1000 spectrophotometer (Thermo Fisher Scientific Inc., USA). RNA integrity was verified on agarose gels. Purified RNA was stored at  $-80^{\circ}\text{C}$ . For expression analyses, 1  $\mu\text{g}$  of RNA was treated with DNase I Amplification Grade (Sigma-Aldrich, USA) and cDNA were synthesized using the iScript cDNA Synthesis Kit (Bio-Rad, USA) as recommended. cDNA was diluted fivefold in RNase/DNase free Tris-EDTA pH 7.4 (Sigma-Aldrich) for initial tests of PIPs in reverse transcriptase PCR. To target plasma membrane-localized aquaporins likely to transport water, the PIP subfamily were selected. Subsequently, tomato-specific PIP primers developed previously<sup>54</sup> were used to pinpoint which PIPs were expressed in the tissues of this study. All initial PCR reactions using gDNA or cDNA were done using Ex taq polymerase (Takara Bio Inc, Japan) as recommended with 2% (v/v) dimethyl sulfoxide in final reactions. PCR conditions were  $94^{\circ}\text{C}$  4 min, 35 cycles of [30 s  $94^{\circ}\text{C}$ , 1 min  $60^{\circ}\text{C}$ , 45 s  $72^{\circ}\text{C}$ ], and 7 min  $72^{\circ}\text{C}$ . Among the 12 PIPs tested (*PIP1.1*–*PIP1.3*, *PIP1.5*, *PIP1.7*, *PIP2.1*, *PIP2.4*–*PIP2.6*, *PIP2.8*, *PIP2.9*, and *PIP2.12*), 4 were not suitable for the subsequent quantitative PCR (qPCR) analyses. *PIP1.2*, *PIP2.6*, and *PIP2.12* were detected in very low abundances or were not expressed. *PIP1.7* was found to be highly unstable and were excluded from the analyses.

#### Quantitative real-time PCR analyses (RT-qPCR)

Reactions of RT-qPCR were performed using SsoAdvanced<sup>TM</sup> Universal SYBR<sup>®</sup> Green Supermix as recommended (Bio-Rad) with a CFX Connect<sup>TM</sup> Real-Time PCR Detection System (Bio-Rad). Analyses of primer temperature optimization, melting curves, standard curves for primer pair efficiencies, Cq values, and normalized expression (Cq) were conducted in CFX Maestro Software supplied by Bio-Rad. In addition to PIP primer pairs, tomato-specific *TIP4.1*, *SAND*, *CAC*, and *Expressed* reference gene candidates developed elsewhere were included in the analyses<sup>55</sup>. *CAC* was selected as reference gene in RefFinder<sup>56</sup>. Primer-specific temperature settings and efficiencies are available in Supporting Information

Table S1. Each treatment type were analyzed with three technical and four biological replicates. Changes to fold change less than twofold up or down were considered minor. The full RT-qPCR assay were conducted twice from the level of RNA extractions.

#### Statistics

Data were statistically analyzed using Microsoft Excel, SPSS 22.0 software (IBM SPSS Software, New York, USA), and CFX Maestro Software (Bio-Rad). The effects of  $\text{CO}_2$  growth environment and GE and their interaction on variables were analyzed using two-way analysis of variance (ANOVA). In addition, in order to discriminate the means between the four treatments, one-way ANOVA (Tukey's test) was conducted to determine the significant differences. Differences between treatments were considered significant when  $P < 0.05$ .

#### Acknowledgements

We thank Professor Ian C. Dodd at the Lancaster Environment Centre (Lancaster University, UK) for sharing ABA-deficient tomato mutant (*flacca*) and wild-type tomato (cv. Ailsa Craig). Technical assistance by Rene Hvidberg Petersen and Lene Korsholm Jørgensen is gratefully acknowledged.

#### Author details

<sup>1</sup>Department of Plant and Environmental Sciences, Faculty of Science, University of Copenhagen, Højbakkegaard Alle 13, 2630 Taastrup, Denmark.

<sup>2</sup>Key Laboratory of Agricultural Soil and Water Engineering in Arid and Semiarid Areas, Ministry of Education, Northwest A&F University, 712100 Yangling, Shaanxi, China.

<sup>3</sup>College of Bioscience and Biotechnology, Hunan Agricultural University, 410128 Changsha, Hunan, China.

<sup>4</sup>Key Laboratory of Mollisols Agroecology, Northeast Institute of Geography and Agroecology, Chinese Academy of Sciences, 130102 Changchun, China.

<sup>5</sup>Present address: Centre for Crop Systems Analysis, Department of Plant Sciences, Wageningen University & Research, PO Box 430, 6700 AK Wageningen, The Netherlands

#### Conflict of interest

The authors declare that they have no conflict of interest.

**Supplementary Information** accompanies this paper at (<https://doi.org/10.1038/s41438-019-0187-6>).

Received: 12 April 2019 Revised: 16 July 2019 Accepted: 3 August 2019

Published online: 11 September 2019

#### References

1. Xu, Z., Jiang, Y., Jia, B. & Zhou, G. Elevated- $\text{CO}_2$  response of stomata and its dependence on environmental factors. *Front. Plant Sci.* **7**, 657 (2016).
2. Royer, D. L. Stomatal density and stomatal index as indicators of paleoatmospheric  $\text{CO}_2$  concentration. *Rev. Palaeobot. Palynol.* **114**, 1–28 (2001).
3. Hetherington, A. M. & Woodward, F. I. The role of stomata in sensing and driving environmental change. *Nature* **424**, 901 (2003).
4. Ainsworth, E. A. & Rogers, A. The response of photosynthesis and stomatal conductance to rising  $[\text{CO}_2]$ : mechanisms and environmental interactions. *Plant Cell Environ.* **30**, 258–270 (2007).
5. Yan, F., Li, X. & Liu, F. ABA signaling and stomatal control in tomato plants exposure to progressive soil drying under ambient and elevated atmospheric  $\text{CO}_2$  concentration. *Environ. Exp. Bot.* **139**, 99–104 (2017).
6. Beerling, D. J. & Woodward, F. I. Changes in land plant function over the Phanerozoic: reconstructions based on the fossil record. *Bot. J. Linn. Soc.* **124**, 137–153 (1997).
7. Woodward, F. I. & Kelly, C. K. The influence of  $\text{CO}_2$  concentration on stomatal density. *New Phytol.* **131**, 311–327 (1995).

8. Woodward, F. I. Do plants really need stomata? *J. Exp. Bot.* **49**, 471–480 (1998).
9. Woodward, F. I., Lake, J. A. & Quick, W. P. Stomatal development and CO<sub>2</sub>: ecological consequences. *New Phytol.* **153**, 477–484 (2002).
10. Chater, C. et al. Elevated CO<sub>2</sub>-induced responses in stomata require ABA and ABA signalling. *Curr. Biol.* **25**, 2709–2716 (2015).
11. Tazoe, Y. & Santrucek, J. Superimposed behaviour of gm under ABA-induced stomata closing and low CO<sub>2</sub>. *Plant Cell Environ.* **38**, 385–387 (2015).
12. Franks, P. J. & Farquhar, G. D. The effect of exogenous abscisic acid on stomatal development, stomatal mechanics, and leaf gas exchange in *Tradescantia virginiana*. *Plant Physiol.* **125**, 935–942 (2001).
13. Lake, J. A. & Woodward, F. I. Response of stomatal numbers to CO<sub>2</sub> and humidity: control by transpiration rate and abscisic acid. *New Phytol.* **179**, 397–404 (2008).
14. Yan, F., Sun, Y., Song, F. & Liu, F. Differential responses of stomatal morphology to partial root-zone drying and deficit irrigation in potato leaves under varied nitrogen rates. *Sci. Hortic.* **145**, 76–83 (2012).
15. Urban, O. Physiological impacts of elevated CO<sub>2</sub> concentration ranging from molecular to whole plant responses. *Photosynthetica* **41**, 9–20 (2003).
16. Mott, K. A. & Buckley, T. N. Stomatal heterogeneity. *J. Exp. Bot.* **49**, 407–417 (1998).
17. Engineer, C. B. et al. CO<sub>2</sub> sensing and CO<sub>2</sub> regulation of stomatal conductance: advances and open questions. *Trends Plant Sci.* **21**, 16–30 (2016).
18. Raschke, K. Stomatal action. *Annu. Rev. Plant Physiol.* **26**, 309–340 (1975).
19. Hsu, P. K. et al. Abscisic acid-independent stomatal CO<sub>2</sub> signal transduction pathway and convergence of CO<sub>2</sub> and ABA signalling downstream of OST1 kinase. *Proc. Natl. Acad. Sci. USA* **115**, E9971–E9980 (2018).
20. Kaiser, W. M. & Hartung, W. Uptake and release of abscisic acid by isolated photoautotrophic mesophyll cells, depending on pH gradients. *Plant Physiol.* **68**, 202–206 (1981).
21. Wullschleger, S. D., Tschaplinski, T. J. & Norby, R. J. Plant water relations at elevated CO<sub>2</sub>—implications for water-limited environments. *Plant Cell Environ.* **25**, 319–331 (2002).
22. Bunce, J. A. Growth at elevated carbon dioxide concentration reduces hydraulic conductance in alfalfa and soybean. *Glob. Change Biol.* **2**, 155–158 (1996).
23. Bunce, J. A. & Ziska, L. H. Decreased hydraulic conductance in plants at elevated carbon dioxide. *Plant Cell Environ.* **21**, 121–126 (1998).
24. Clarkson, D. T. et al. Root hydraulic conductance: diurnal aquaporin expression and the effects of nutrient stress. *J. Exp. Bot.* **51**, 61–70 (2000).
25. Kaldenhoff, R. & Fischer, M. Functional aquaporin diversity in plants. *Biochim. Biophys. Acta* **1758**, 1134–1141 (2006).
26. Locke, A. M. & Ort, D. R. Diurnal depression in leaf hydraulic conductance at ambient and elevated [CO<sub>2</sub>] reveals anisohydric water management in field-grown soybean and possible involvement of aquaporins. *Environ. Exp. Bot.* **116**, 39–46 (2015).
27. Wand, S. J., Midgley, G. F., Jones, M. H. & Curtis, P. S. Responses of wild C<sub>4</sub> and C<sub>3</sub> grass (Poaceae) species to elevated atmospheric CO<sub>2</sub> concentration: a meta-analytic test of current theories and perceptions. *Glob. Change Biol.* **5**, 723–741 (1999).
28. Davies, W. J. & Zhang, J. Root signals and the regulation of growth and development of plants in drying soil. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **42**, 55–76 (1991).
29. Yong, J. W. H., Wong, S. C., Letham, D. S., Hocart, C. H. & Farquhar, G. D. Effect of elevated [CO<sub>2</sub>] and nitrogen nutrition on cytolinins in the xylem sap and leaves of cotton. *Plant Physiol.* **124**, 767–779 (2000).
30. Mamatha, H., Srinivasa Rao, N. K. & Vijayalakshmi, T. Physiological responses of tomato (*Lycopersicon esculentum* Mill) cv. Arka Ashish to elevated atmospheric CO<sub>2</sub> under water limiting conditions. *Indian J. Agric. Res.* **49**, 299–307 (2015).
31. McAdam, S. A. M. & Brodrribb, T. J. Linking turgor with ABA biosynthesis: implications for stomatal responses to vapor pressure deficit across land plants. *Plant Physiol.* **171**, 2008–2016 (2016).
32. Sack, L., John, G. P. & Buckley, T. N. ABA accumulation in dehydrating leaves is associated with decline in cell volume, not turgor pressure. *Plant Physiol.* **176**, 489–493 (2018).
33. Liu, F., Jensen, C. R. & Andersen, M. N. A review of drought adaptation in crop plants: changes in vegetative and reproductive physiology induced by ABA-based chemical signals. *Aust. J. Agric. Res.* **56**, 1245–1252 (2005).
34. Li, X. et al. Carbon dioxide enrichment alleviates heat stress by improving cellular redox homeostasis through an ABA-independent process in tomato plants. *Plant Biol.* **17**, 81–89 (2015).
35. Li, X., Jiang, D. & Liu, F. Soil warming enhances the hidden shift of elemental stoichiometry by elevated CO<sub>2</sub> in wheat. *Sci. Rep.* **6**, 23313 (2016). p.
36. Bahrin, A., Jensen, C. R., Asch, F. & Mogensen, V. O. Drought-induced changes in xylem pH, ionic composition, and ABA concentration act as early signals in field-grown maize (*Zea mays* L.). *J. Exp. Bot.* **53**, 251–263 (2002).
37. Seo, M. & Koshiba, T. Transport of ABA from the site of biosynthesis to the site of action. *J. Plant Res.* **124**, 501–507 (2011).
38. Farquhar, G. D. & Sharkey, T. D. Stomatal conductance and photosynthesis. *Annu. Rev. Plant Physiol.* **33**, 317–345 (1982).
39. Dodd, I. C., Theobald, J. C., Richer, S. K. & Davies, W. J. Partial phenotypic reversion of ABA-deficient *flacca* tomato (*Solanum lycopersicum*) scions by a wild-type rootstock: normalizing shoot ethylene relations promotes leaf area but does not diminish whole plant transpiration rate. *J. Exp. Bot.* **60**, 4029–4039 (2009).
40. Trouverie, J., Thévenot, C., Rocher, J.-P., Sotta, B. & Prioul, J.-L. The role of abscisic acid in the response of a specific vacuolar invertase to water stress in the adult maize leaf. *J. Exp. Bot.* **54**, 2177–2186 (2003).
41. Bunce, J. A. Effects of humidity on short-term responses of stomatal conductance to an increase in carbon dioxide concentration. *Plant Cell Environ.* **21**, 115–120 (1998).
42. Robredo, A. et al. Elevated CO<sub>2</sub> alleviates the impact of drought on barley improving water status by lowering stomatal conductance and delaying its effects on photosynthesis. *Environ. Exp. Bot.* **59**, 252–263 (2007).
43. Hose, E., Steudle, E. & Hartung, W. Abscisic acid and hydraulic conductivity of maize roots: a study using cell-and root-pressure probes. *Planta* **211**, 874–882 (2000).
44. Parent, B. et al. Drought and abscisic acid effects on aquaporin content translate into changes in hydraulic conductivity and leaf growth rate: a trans-scale approach. *Plant Physiol.* **149**, 2000–2012 (2009).
45. Pantin, F. et al. The dual effect of abscisic acid on stomata. *New Phytol.* **197**, 65–72 (2013).
46. Veselov, D. S. et al. Rapid changes in root HvPIP2; 2 aquaporins abundance and ABA concentration are required to enhance root hydraulic conductivity and maintain leaf water potential in response to increased evaporative demand. *Funct. Plant Biol.* **45**, 143–149 (2018).
47. Zaghdoud, C. et al. Elevated CO<sub>2</sub> alleviates negative effects of salinity on broccoli (*Brassica oleracea* L. var *Italica*) plants by modulating water balance through aquaporins abundance. *Environ. Exp. Bot.* **95**, 15–24 (2013).
48. Secchi, F., Schubert, A. & Lovisolo, C. Changes in air CO<sub>2</sub> concentration differentially alter transcript levels of *NtAQP1* and *NtPIP2;1* aquaporin genes in tobacco leaves. *Int. J. Mol. Sci.* **17**, 567 (2016).
49. Jang, J. Y., Kim, D. G., Kim, Y. O., Kim, J. S. & Kang, H. An expression analysis of a gene family encoding plasma membrane aquaporins in response to abiotic stress in *Arabidopsis thaliana*. *Plant Mol. Biol.* **54**, 713–725 (2004).
50. Ding, L. et al. The enhanced drought tolerance of rice plants under ammonium is related to aquaporin (AQP). *Plant Sci.* **234**, 14–21 (2015).
51. Taylor, I. B., Linforth, R. S. T., Al-Naib, R. J., Bowman, W. R. & Marples, B. A. The wilty tomato mutants *flacca* and *sitiens* are impaired in the oxidation of ABA-aldehyde to ABA. *Plant Cell Environ.* **11**, 739–745 (1988).
52. Netting, A. G., Theobald, J. C. & Dodd, I. C. Xylem sap collection and extraction methodologies to determine *in vivo* concentrations of ABA and its bound forms by gas chromatography-mass spectrometry (GC-MS). *Plant Methods* **8**, 11 (2012).
53. Asch, F. *Determination of Abscisic Acid by Indirect Enzyme Linked Immuno Sorbent Assay (ELISA)*. Technical Report, (Laboratory for Agro-hydrology and Bioclimatology, Department of Agricultural Sciences, 2000).
54. Reuscher, S. et al. Genome-wide identification and expression analysis of aquaporins in tomato. *PLoS ONE* **8**, e79052 (2013).
55. Expósito-Rodríguez, M., Borges, A. A., Borges-Pérez, A. & Pérez, J. A. Selection of internal control genes for quantitative real-time RT-PCR studies during tomato development process. *BMC Plant Biol.* **22**, 131 (2008).
56. Xie, F., Xiao, P., Chen, D., Xu, L. & Zhang, B. miRDeepFinder: a miRNA analysis tool for deep sequencing of plant small RNAs. *Plant Mol. Biol.* **80**, 75–84 (2012).