

SCIENTIFIC REPORTS



OPEN

Publisher Correction: Upregulation of prefrontal metabotropic glutamate receptor 5 mediates neuropathic pain and negative mood symptoms after spinal nerve injury in rats

Geehoon Chung  ^{1,2}, Chae Young Kim ^{1,3}, Yeong-Chan Yun ¹, Sang Ho Yoon ^{1,3}, Myoung-Hwan Kim ^{1,3}, Yu Kyeong Kim ⁴ & Sang Jeong Kim  ^{1,2,3,5}

Correction to: *Scientific Reports* <https://doi.org/10.1038/s41598-017-09991-8>, published online 29 August 2017

In this Article, the two parts of Table 1 are merged together. The correct Table 1 appears below.

¹Department of Physiology, Seoul National University College of Medicine, Seoul, Korea. ²Department of Brain and Cognitive Sciences, Seoul National University College of Natural Sciences, Seoul, Korea. ³Department of Biomedical Sciences, Seoul National University College of Medicine, Seoul, Korea. ⁴Department of Nuclear Medicine, Seoul National University College of Medicine, Seoul, Korea. ⁵Neuroscience Research Institute, Seoul National University College of Medicine, Seoul, Korea. Geehoon Chung and Chae Young Kim contributed equally to this work. Correspondence and requests for materials should be addressed to S.J.K. (email: sangjikim@snu.ac.kr)

Brain region	K _e	T	Z	Peak p	Location (mm)		
					ML	AP	DV
Sham < SNL							
Medial prefrontal cortex(Prelimbic and infralimbic cortices)	97	5.50	4.06	2.54E-06	0.2	2.6	-4.8
Primary somatosensory cortex(Dysgranular zone)	24	5.88	4.23	1.16E-05	-4.6	-0.2	-3.4
Primary, secondary somatosensory cortices and caudate putamen	37	4.42	3.52	0.0002	4.8	-0.4	-5.2
Retrosplenial cortex	25	6.70	4.56	2.45E-05	0.2	-4.6	-2.4
Medial septum	30	3.51	2.98	0.0014	0.4	-0.8	-4.8
Sham > SNL							
Insular cortex and piriform cortex	102	7.64	4.89	4.98E-07	-3.2	3.6	-7.0
		4.73	3.69	0.0001	-3.4	3.0	-7.6
		4.60	3.62	0.0001	-3.6	2.6	-5.8
Insular cortex and primary somatosensory cortex	31	4.12	3.35	0.0004	-4.8	3.4	-4.8
Endopiriform nucleus	20	4.06	3.32	0.0005	3.8	1.4	-7.2
Nucleus accumbens	30	3.74	3.12	0.0009	1.6	1.8	-6.4
Piriform cortex	61	3.72	3.11	0.0009	-6.2	-4.8	-8.6
Corpus callosum and caudate putamen	36	3.71	3.11	0.0009	1.6	-0.4	-3.4
Corpus callosum and caudate putamen	76	3.65	3.07	0.0011	3.6	-2.0	-3.8

Table 1. Brain regions with significant differences between sham and SNL rats (two-sample t-test, $p < 0.005$, $K_e > 20$). The peak voxel location is represented by the distance from the bregma (mm).



Open Access This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons license and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>.

© The Author(s) 2018