

# SCIENTIFIC REPORTS



OPEN

## Plant tolerance to excess light energy and photooxidative damage relies on plastoquinone biosynthesis

Received: 08 October 2014

Accepted: 07 May 2015

Published: 03 June 2015


Brigitte Ksas<sup>1,2,3,\*</sup>, Noëlle Becuwe<sup>1,2,3,\*</sup>, Anne Chevalier<sup>1,2,3</sup> & Michel Havaux<sup>1,2,3</sup>

Plastoquinone-9 is known as a photosynthetic electron carrier to which has also been attributed a role in the regulation of gene expression and enzyme activities via its redox state. Here, we show that it acts also as an antioxidant in plant leaves, playing a central photoprotective role. When *Arabidopsis* plants were suddenly exposed to excess light energy, a rapid consumption of plastoquinone-9 occurred, followed by a progressive increase in concentration during the acclimation phase. By overexpressing the plastoquinone-9 biosynthesis gene *SPS1* (*SOLANESYL DIPHOSPHATE SYNTHASE 1*) in *Arabidopsis*, we succeeded in generating plants that specifically accumulate plastoquinone-9 and its derivative plastoehromanol-8. The *SPS1*-overexpressing lines were much more resistant to photooxidative stress than the wild type, showing marked decreases in leaf bleaching, lipid peroxidation and PSII photoinhibition under excess light. Comparison of the *SPS1* overexpressors with other prenyl quinone mutants indicated that the enhanced phototolerance of the former plants is directly related to their increased capacities for plastoquinone-9 biosynthesis.

Photosynthesis inevitably produces toxic molecules derived from oxygen. Indeed, molecular oxygen can interact with the photosynthetic electron transport chain, leading to the formation of reduced forms of oxygen, such as superoxide, or with excited chlorophyll molecules, generating singlet oxygen ( ${}^1\text{O}_2$ )<sup>1–3</sup>. Superoxide can spontaneously or enzymatically dismutate into hydrogen peroxide which can subsequently lead to the hydroxyl radical in the presence of metals. Light-induced production of reactive oxygen species (ROS) is amplified under environmental stress conditions when the photosynthetic processes are inhibited and the absorption of light energy becomes excessive relative to the photosynthetic activity. One way to dissipate this excess energy is to transfer electrons and/or excitation to oxygen. However, to cope with the resulting production of harmful ROS, chloroplasts contain a variety of antioxidant mechanisms including soluble and lipophilic low molecular weight antioxidants<sup>4–6</sup>, detoxification enzymes and repair mechanisms<sup>7–9</sup>.

${}^1\text{O}_2$  is produced within the photosystems (PS) from excited chlorophyll molecules in the triplet state<sup>10,11</sup>.  ${}^1\text{O}_2$  is thought to be the major ROS produced in plant cells at high light intensities<sup>12</sup> and to be instrumental in the execution of ROS-induced cell death in leaves<sup>13</sup>. This ROS has a short lifetime (*ca.* 100 ns in biological tissues), suggesting a small diffusion path in cells<sup>11</sup>. Consequently,  ${}^1\text{O}_2$  reacts primarily in the close vicinity of its production site, and efficient  ${}^1\text{O}_2$  detoxification mechanisms must function close to the sites of  ${}^1\text{O}_2$  production. Accordingly, thylakoid membranes contain various lipid-soluble compounds that can quench  ${}^1\text{O}_2$  within the photosystems and around. Carotenoids are considered to be the first line of defense against  ${}^1\text{O}_2$  toxicity because of their high efficiency of  ${}^1\text{O}_2$  quenching and their

<sup>1</sup>CEA, IBEB, Laboratoire d'Ecophysiologie Moléculaire des Plantes, F-13108 Saint-Paul-lez-Durance, France. <sup>2</sup>CNRS, UMR 7265 Biologie Végétale et Microbiologie Environnementales, F-13108 Saint-Paul-lez-Durance, France. <sup>3</sup>Aix-Marseille Université, F-13284 Marseille, France. \*These authors contributed equally to the article. Correspondence and requests for materials should be addressed to M.H. (email: michel.havaux@cea.fr)




**Figure 1. Responses of WT *Arabidopsis* plants to excess light energy.** Plants (Col 0) aged 4 weeks were grown at a PFD of  $170 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Plants were exposed to high light (PFD,  $1300 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) and low air temperature ( $5^\circ\text{C}$ ). **A**) Maximal quantum yield of PSII photochemistry (Fv/Fm chlorophyll fluorescence ratio). Data are mean values of 10 measurements  $\pm$  SD. **B**) Changes in prenyl lipid concentrations in relative values:  $\alpha$ -tocopherol ( $\alpha$ -Toc), plastoquinol-8 (PC) and total plastoquinone-9 (PQ). For each compound, values were normalized to the control value at time 0. Data are mean values of 4 measurements  $\pm$  SD. A value of 1 corresponds to  $1.84 \text{ nmol cm}^{-2}$  for plastoquinone-9,  $19.3 \text{ pmol cm}^{-2}$  for plastoquinol-8 and  $270 \text{ ng cm}^{-2}$  for  $\alpha$ -tocopherol.

localization in close proximity with the chlorophyll molecules in the light-harvesting complexes and the reaction centers of the photosystems<sup>4,14</sup>. However, prenyl lipids of the tocopherol family have also been shown to participate in the protection against  ${}^1\text{O}_2$ . Tocopherols can quench  ${}^1\text{O}_2$ , thus protecting PSII from photoinhibition, and can terminate lipid peroxidation chain-reactions, thus protecting the thylakoid membranes<sup>6,15–18</sup>. However, chloroplasts contain other prenyl lipids, such as plastoquinone-9, which could provide additional protection against photooxidative stress. Plastoquinones are viewed essentially as mobile electron carriers involved in electron transfer between PSII and PSI<sup>19</sup>. Through their redox state, there are also recognized as regulators of gene expression and enzyme activities<sup>20,21</sup>. However, it has been shown *in vitro* that plastoquinone-9 also has protective and antioxidant properties, being able to dissipate energy in the chlorophyll antennae<sup>22</sup>, to quench  ${}^1\text{O}_2$  and to inhibit oxidation of lipid membranes<sup>23–25</sup>. If this function does occur *in vivo*, it could be of great physiological importance because plastoquinones are diffusible molecules present in relatively high amounts in the thylakoids (estimated to be  $\sim 7$  molecules per PSII reaction center, ref. 26). Moreover, both the head group and the isoprenoid side chain of plastoquinol are able to quench  ${}^1\text{O}_2$ , likely making this molecule a better antioxidant than tocopherols. The possible role of plastoquinone-9 *in planta* as an antioxidant and photoprotector is analyzed here in leaves of the model plant *Arabidopsis thaliana*. To this end, the plastoquinone biosynthesis pathway was manipulated to generate plants that contain noticeably more plastoquinone-9 than the wild type (WT), and the behavior of those plastoquinone-accumulating plants under high light stress conditions was compared to that of wild-type (WT) plants.

## Results

**The plastoquinone-9 content of *Arabidopsis* leaves is highly sensitive to excess light energy.** High excitation pressure on the photosystems can be achieved by increasing light irradiance and/or decreasing temperature<sup>27</sup>. WT *Arabidopsis* plants aged 4 weeks were suddenly exposed to high photon flux density (PFD,  $1300 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ ) and low air temperature ( $5^\circ\text{C}$ ). The decreased air temperature led to leaf temperature (around  $18^\circ\text{C}$ ) in the temperature range of control leaves grown at  $170 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ , thus avoiding leaf heating usually associated with high PFDs. Under those conditions, PSII underwent rapid photoinhibition, as shown by the fall of the Fv/Fm chlorophyll fluorescence ratio from 0.8 to 0.2 after 7 h in high light (Fig. 1A). Then, PSII slowly recovered, with Fv/Fm progressively increasing and finally reaching, after 5 d in high light, values close to the initial values recorded before stress. Plastoquinone-9 (reduced + oxidized) appeared to follow the same trends: it drastically decreased by 80% after 7 h in high light (Fig. 1B). Subsequently, the total plastoquinone-9 concentration increased, finally leading to a strong accumulation corresponding to almost 4 times the




**Figure 2. Responses of the *Arabidopsis ch1* mutant to excess light energy.** Plants grown for 5 weeks at PFD 170  $\mu\text{mol}$  photons  $\text{m}^{-2} \text{s}^{-1}$  were exposed to high light (PFD, 1000  $\mu\text{mol}$  photons  $\text{m}^{-2} \text{s}^{-1}$ ) and low temperature (10°C). **A**) Maximal quantum yield of PSII photochemistry (Fv/Fm chlorophyll fluorescence ratio). Data are mean values of 10 measurements  $\pm$  SD. **B**) Changes in prenyl lipid concentrations in relative values:  $\alpha$ -tocopherol ( $\alpha$ -Toc), plastoChroman-8 (PC) and total plastoquinone-9 (PQ). For each compound, values were normalized to the control value at time 0. Data are mean values of 4 measurements  $\pm$  SD. A value of 1 corresponds to 1.9  $\text{nmol cm}^{-2}$  for plastoquinone-9, 16.8  $\text{pmol cm}^{-2}$  for plastoChroman-8 and 539  $\text{ng cm}^{-2}$  for  $\alpha$ -tocopherol.

plastoquinone-9 levels measured before stress. Those changes in plastoquinone-9 concentration were associated with limited changes in the plastoquinone reduction state: the plastoquinone-9 reduction levels decreased from ca. 80% reduction to ca. 70% during the high light treatment and increased back to the initial values after 5 d in high light (Supplementary Fig. 1). The plastoChroman-8 content also changed with the light conditions, but compared to plastoquinone-9, the changes were very attenuated and occurred much more slowly (Fig. 1B). In contrast with plastoChroman-8 and plastoquinone-9, no loss of  $\alpha$ -tocopherol was observed during high light stress: the  $\alpha$ -tocopherol concentration remained stable during the first two days of light stress and then it increased noticeably.

We also analyzed a photosensitive *Arabidopsis* mutant, the *ch1* mutant, which has been shown to release more singlet oxygen ( $^1\text{O}_2$ ) from the PSII reaction centers compared to WT<sup>28</sup>. Because of their high photosensitivity, *ch1* mutant plants aged 5 weeks were exposed to less severe stress conditions than those used for WT in Fig. 1: 1000  $\mu\text{mol}$  photons  $\text{m}^{-2} \text{s}^{-1}$  at 10°C. Nevertheless, this milder treatment caused a drastic inhibition of PSII photochemistry which did not reverse (Fig. 2A). Again, plastoquinone-9 underwent drastic changes in concentration during the high light treatment which mirrored the inhibition of PSII photochemistry (Fig. 2B). There was a decrease in the percentage reduction state of plastoquinone-9 from ca. 80% at time 0 to ca. 60% at the end of the experiments after 50 h of light stress (Supplementary Fig. 1). PlastoChroman-8 behaved similarly to plastoquinone-9, exhibiting a noticeable decrease, but this effect was delayed relative to the plastoquinone-9 changes. In comparison,  $\alpha$ -tocopherol exhibited very limited changes and remained close to the control levels before stress. To sum up, among the prenyl lipids of *Arabidopsis* leaves, plastoquinone-9 appeared to be by far the most sensitive to the light conditions, displaying strong reductions during photooxidative stress and marked accumulations during stress acclimation. In addition, we observed a correlation between the changes in PSII activity and the changes in plastoquinone-9 concentration.

**High light-induced changes in the expression of genes involved in the plastoquinone-9 biosynthesis pathway.** A scheme of the biosynthetic pathways of plastoquinone-9, plastoChroman-8 and  $\alpha$ -tocopherol is presented in Fig. 3A. The first committed step in plastoquinone-9 biosynthesis is the condensation of the aromatic compound homogentisate (HGA) with solanesyl diphosphate by homogentisate solanesyl diphosphate transferase (HST), leading to the formation of MSBQ (methyl-solanesyl-benzoquinone). HGA is shared in common with the tocopherol biosynthetic pathway, with prenylation of HGA with phytol diphosphate producing MPBQ (methyl-phytol-benzoquinone), a precursor of  $\alpha$ -tocopherol. MPBQ is converted to  $\alpha$ -tocopherol by the action of VTE3, a



**Figure 3. Light regulation of the plastoquinone-9 biosynthesis pathway. A)** Scheme of the biosynthesis pathway. **B)** qRT-PCR measurements of the expression of several key genes involved in the biosynthesis of plastoquinone-9 in leaves of WT *Arabidopsis* plants exposed to high light (1300  $\mu\text{mol}$  photons  $\text{m}^{-2}$   $\text{s}^{-1}$  at air temperature of 7°C). Data are normalized to the values at time 0. Data are mean values of 3 separate experiments  $\pm$  SD. The expression level of VTE3 at time 147 h (87  $\pm$  2) is out of scale in the graph. Metabolites: HPP, hydroxyphenylpyruvate; HGA, homogentisate; DP, diphosphate; IPP, isopentenyl phosphate; GGDP, geranylgeranyl diphosphate; MPBQ, methyl-phytyl-benzoquinone; MSBQ, methyl-solanesyl-benzoquinone, DMPBQ, dimethyl-phytyl-benzoquinone. Enzymes: HPPD, HPP dioxygenase; HST, homogentisate solanesyl diphosphate transferase; SPS, solanesyl diphosphate synthase; VTE1, tocopherol cyclase; VTE2, homogentisate phytol transferase; VTE3, MPBQ/MSBQ methyl transferase; VTE4,  $\gamma$ -tocopherol methyl transferase.

methyl transferase<sup>29</sup> that is also involved in plastoquinone-9 biosynthesis by converting MSBQ (methyl-solanesyl-benzoquinone) to plastoquinone.

HGA synthesis from hydroxyphenylpyruvate (HPP) is catalyzed by HPP dioxygenase (HPPD) while solanesyl diphosphate is synthesized from geranylgeranyl diphosphate (GGDP) and isopentenyl phosphate (IPP) by a reaction catalyzed by solanesyl diphosphate synthase (SPS). Three SPS enzymes are present in *Arabidopsis*, SPS1, SPS2 and SPS3<sup>30,31</sup>. While SPS3 is a ubiqinone biosynthetic enzyme localized in mitochondria<sup>31</sup>, SPS1 and SPS2 have been recently demonstrated to be targeted to plastids and to be involved in plastoquinone biosynthesis<sup>32</sup>. Previously, SPS1 was supposed to be involved in the synthesis of the side chain of ubiquinone while plastoquinone biosynthesis was believed to be dependent on SPS2 only<sup>33</sup>. However, this view has been challenged by recent observations revealing that a mitochondria-targeted gene (SPS3) different from SPS1 and SPS2 is the main, if not sole, contributor of solanesyl diphosphate synthase activity required for ubiquinone biosynthesis<sup>31</sup>. Moreover, the *Arabidopsis* mutants *AtSPS1* and *AtSPS2* were found to be affected in plastoquinone-9 and plastochromanol-8 biosynthesis, not in ubiquinone-9 synthesis. Thus, the current view is that both SPS1 and SPS2 catalyze the elongation of the prenyl side chain of plastoquinone<sup>32</sup>. Plastochromanol-8 has been demonstrated to originate from reduced plastoquinone-9 through the action of VTE1<sup>34,35</sup>, a tocopherol cyclase enzyme also involved in the biosynthesis of  $\alpha$ -tocopherol from its direct precursor,  $\gamma$ -tocopherol (Fig. 3A).

In Fig. 3B, we examined the effect of excess light energy on the expression of several genes of the plastoquinone-9 and plastochromanol-8 biosynthesis pathway. One can see that the expression of both SPS1 and SPS2 genes was rapidly induced after transfer of plants aged 4 weeks from low light to high


light, with the accumulation of *SPS1* transcripts being noticeably more pronounced than that of *SPS2*. The expression pattern of *HPPD* was close to that of *SPS2*, with an induction in high light. In striking contrast, *HST* expression was not affected by light. The *VTE1* and *VTE3* genes were also activated by high light but this effect was more progressive and continuous than the up-regulation of *SPS1*, *SPS2* and *HPPD*. So, the plastoquinone-9 biosynthesis pathway is globally up-regulated by high light, with a marked effect on the *SPS1* gene in less than 3 h after the transfer from low light to high light. Light induction of the plastoquinone pathway is consistent with early data on the incorporation of radiolabelled tyrosine into prenyl lipids<sup>36</sup>. Upon illumination of *Xanthium* leaves, incorporation of radioactivity into plastoquinone was observed to be much more pronounced and to occur more rapidly than incorporation into tocopherols.

**Synthesis of plastoquinone-9 and plastoeranol-8 is boosted in *SPS1*-overexpressing plants and correlates with tolerance to excess light energy.** A previous work has shown that constitutive overexpression of *HST* in *Arabidopsis* has little effect on the plastoquinone-9 concentration in leaves<sup>37</sup>, suggesting that *HST* activity is not the limiting step for plastoquinone-9 biosynthesis. Our observation that *HST* gene expression is not responsive to a condition associated with plastoquinone-9 accumulation could be seen as a fact in line with this suggestion. Considering the strong and rapid expression of *SPS1* under conditions that induced plastoquinone-9 accumulation in leaves (Figs. 1,3), we decided to overexpress this gene in *Arabidopsis*. *Arabidopsis SPS1* cDNA was inserted under the control of the 35S promoter in a plant binary vector. This vector was used to generate transgenic *Arabidopsis* plants, and a number of stable lines derived from independent transformation events were obtained. Figure 4A shows a selection of homozygous lines (SPS1oex) exhibiting a strong accumulation of *SPS1* transcripts. This transformation had marked effects on plastoquinone-9 and its derivative plastoeranol-8 which accumulated in all lines (Fig. 4B). The effect was particularly marked for plastoeranol-8 with an accumulation factor of 2 to 3. The plastoquinone-9 accumulation was less marked, but nevertheless reached approximately 150–170% of the WT level. The reduction state of plastoquinone-9 increased very slightly in the SPS1oex leaves (Supplementary Fig. 2). In contrast, the  $\alpha$ -tocopherol concentration did not change significantly in any of the transformed lines. Similarly, the levels of minor forms of tocopherol ( $\delta$ - and  $\gamma$ -tocopherol) were not modified by the *SPS1* overexpression (data not shown). Thus, constitutive overexpression of *SPS1* selectively boosted the plastoquinone/plastoeranol pathway, as expected. Importantly, plastoquinone-9 and plastoeranol-8 accumulation in SPS1oex plants had no effect on growth (Supplementary Fig. 3A), did not modify the quantum yield of photosynthetic electron transport (Supplementary Fig. 3B) and did not affect the chlorophyll levels (Supplementary Fig. 3C). This could suggest that the extra plastoquinone-9 molecules that accumulate in the SPS1oex lines are not stored in the thylakoid membranes and are not connected to the photosynthetic electron transport chain. The pool of photoactive plastoquinone-9 (connected to the electron transport chain) was estimated in Fig. 5. The size of this pool did not differ significantly between SPS1oex leaves and WT leaves. In contrast, the pool of non-active plastoquinones was noticeably enlarged in the *SPS1*-overexpressing leaves, suggesting plastoquinone-9 storage in a compartment different from the thylakoid membranes.

The plastoquinone/plastoeranol-accumulating SPS1oex plants were exposed to photooxidative stress. We chose rather severe stress conditions in order to induce photooxidative damage in WT plants:  $1300\text{ }\mu\text{mol photons m}^{-2}\text{ s}^{-1}$  at  $5^\circ\text{C}$  with a photoperiod of 13 h. Moreover, for this experiment, plants were grown at the PFD of  $110\text{ }\mu\text{mol m}^{-2}\text{ s}^{-1}$  to increase the sensitivity of WT plants to high light stress. As shown by the autoluminescence image of Fig. 6A and the quantification of the autoluminescence signal in Fig. 6B, WT accumulated lipid peroxides after 28 h of such stress treatment. The increased oxidation of lipids was confirmed by the amount of hydroxy octadecatrienoic acid (HOTE, the hydroxy fatty acid derived from the oxidation of linolenic acid, the most abundant fatty acid in *Arabidopsis* leaves), which drastically increased in WT leaves (Fig. 6C). In striking contrast, the SPS1oex lines were much less sensitive to the light stress, with a noticeable reduction of lipid peroxidation (Fig. 6A–C). Extensive leaf bleaching was observed in WT plants, but not in the SPS1oex lines (data not shown). The increased phototolerance of SPS1oex lines is confirmed by the maintenance of higher PSII photochemical efficiency ( $F_v/F_m$ ) compared to WT (Fig. 6D).


Figures 6E and 6F show the time course of the changes in total plastoquinone-9 and in plastoeranol-8 in one of the SPS1oex lines (#3) in high light. Plastoquinone-9 initially decreased in high light-stressed WT and SPS1oex plants. Subsequently, the plastoquinone concentration increased, as previously shown in Fig. 1, but this effect was much more rapid and more pronounced in the SPS1oex line compared to WT. Thus, compared to WT, the plastoquinone-9 levels were much higher in the *SPS1* overexpressors throughout the stress treatment, and the difference was particularly marked during the acclimation phase after several days in high light. After 6 d in high light, the plastoquinone-9 levels were almost doubled in the SPS1oex line relative to WT. The plastoquinone reduction state was slightly higher in the SPS1oex #3 line relative to WT, but it did not change much with the light stress conditions (Supplementary Fig. 2). In contrast with plastoquinone-9, plastoeranol-8 was observed to decrease in both types of plants during the stress experiment (Fig. 6F).

**The phototolerance of tocopherol-deficient *Arabidopsis vte* mutants.** The *vte1* and *vte2* mutants are deficient in tocopherol cyclase and homogentisate phytol transferase, respectively, leading



**Figure 4. Overexpression of the SPS1 gene in *Arabidopsis*.** A) RT-PCR analysis of SPS1 transcripts in WT (col 0) and SPS1oex transformed lines. ACTIN2 mRNAs were used as loading controls. The SPS1 and ACTIN2 images are from the same experiment and are therefore directly comparable. B) Plastoquinone-9 (PQ), plastoehromanol-8 (PC) and  $\alpha$ -tocopherol ( $\alpha$ -Toc) levels in leaves. Values (measured on a leaf area basis) are normalized to the WT values. Data are mean values of 4 separate experiments  $\pm$  SD. The leaf specific weight (in mg dry weight  $\text{cm}^{-2}$ ) did not differ significantly between the genotypes:  $1.1460 \pm 0.1850$  for Col 0,  $1.2203 \pm 0.0808$  for SPS1oex #3,  $1.1670 \pm 0.0813$  for SPS1oex #12 and  $1.1354 \pm 0.1485$  for SPS1oex #14. C) Growth phenotype of the plants (WT and 3 SPS1oex lines). Plants were grown for 4 weeks at PFD  $170 \mu\text{mol m}^{-2} \text{s}^{-1}$ .

to a complete deficiency in tocopherols<sup>38,39</sup>, as confirmed in Fig. 7A. As previously reported<sup>34,35</sup>, *vte1* is also deficient in plastoehromanol-8 while the plastoehromanol-8 content of *vte2* leaves is normal (Fig. 7C). Interestingly, the combined lack of tocopherols and plastoehromanol-8 in *vte1* leaves was associated in a substantial decrease in plastoquinone-9 levels (Fig. 7B). Under normal growth conditions, the plastoquinone-9 pool was decreased to less than 75% of the WT level. As previously shown<sup>40,41</sup>, neither chlorophylls nor carotenoids are affected in the *vte1* and *vte2* mutants. Under high light stress, the plastoquinone-9 concentration fell to less than 50% of the WT level. In contrast, loss of tocopherol with normal levels of plastoehromanol-8 in the *vte2* mutant had no impact on the plastoquinone-9 concentration. When exposed to high light stress ( $1300 \mu\text{mol m}^{-2} \text{s}^{-1}$  and  $8^\circ\text{C}$ ), pronounced photooxidative




**Figure 5. Pool size of photoactive and non-photoactive plastoquinone-9.** Measurements were performed as detailed in the Methods section on leaves of WT Arabidopsis and two SPS1oex lines (#3 and #12). Data are mean values of 3 separate experiments  $\pm$  SD. F.W.= fresh weight. The stars indicate significant difference between WT and the SPS1oex lines at  $P < 0.05$  (Student's t test).

stress occurred in *vte1* leaves, as illustrated by the accumulation of lipid peroxides (Fig. 7D), the increase in hydroxy fatty acids (HOTEs) (Fig. 7E), the extensive leaf bleaching (data not shown), and the loss of PSII activity (Fv/Fm) (Fig. 7F). These effects were attenuated in WT and *vte2* plants which behaved similarly in response to high light stress. One can thus conclude that loss of tocopherol has rather limited effects on the photosensitivity of Arabidopsis, unless plastoeranol-8 is also missing. This suggests a functional overlap between tocopherols and plastoeranol-8 under high light stress. Our results also show that the protective action of those compounds is required to preserve the plastoquinone pool size.


**The phototolerance of VTE1-overexpressing plants.** Overexpression of the *VTE1* gene in Arabidopsis has two effects: it brings about a strong accumulation of plastoeranol-8<sup>35</sup> and it enhances the production of  $\gamma$ -tocopherol<sup>40</sup>, as expected from the functions of this gene in the biosynthesis pathway (Fig. 3A) and as confirmed in Fig. 8A-C. The plastoquinone-9 levels were not significantly affected by the genetic modification (Fig. 8B). Also, the concentration of  $\alpha$ -tocopherol was not modified in the VTE1 overexpressor (Fig. 8A), so that the increase in the total amount of tocopherols ( $\alpha + \gamma$ ) was rather moderate (ca. 120 vs 100 ng  $\text{cm}^{-2}$ ).  $\gamma$ -Tocopherol is a potent antioxidant *in vivo*, reducing oxidative stress-induced lipid peroxidation even better than  $\alpha$ -tocopherol<sup>42</sup>. In some plant species such as runner bean, young leaves contain very high levels of  $\gamma$ -tocopherol, supplementing other lipophilic antioxidants and contributing to the overall protection against oxidative stress<sup>43</sup>. So, the phototolerance of the VTE1 overexpressors should not be negatively affected by the changes in tocopherol composition. Neither the growth rate, nor the chlorophyll content of the VTE1 overexpressors was different from WT values (total chlorophyll in  $\mu\text{g cm}^{-2}$ : 23.51  $\pm$  1.85 for WT, 23.98  $\pm$  1.06 for VTE1oex #1 and 24.12  $\pm$  0.64 for VTE1oex#40). Both WT and the VTE1 overexpressors were exposed to a high light stress condition that induced photodamage and lipid peroxidation in WT leaves. To this end, we decreased the growth PFD (to 110  $\mu\text{mol m}^{-2} \text{s}^{-1}$ ) and increased the severity of the stress treatment (as in Fig. 6). Under those conditions, WT plants exhibited high autoluminescence emission, high HOTE levels and low Fv/Fm ratios. The VTE1oex lines behaved like WT: pronounced lipid peroxidation (Fig. 8D-E) and extensive PSII photoinhibition (Fig. 8F) were observed in all plants. Thus, accumulation of plastoeranol-8 itself is not sufficient to provide significant protection against photostress. This observation is in line with previous data obtained by Zbierzak *et al.*<sup>35</sup> who did not find any change in photosynthesis in the VTE1oex lines after several days in high light.

## Discussion


Plants are known to accumulate tocopherols in response to high PFDs (e.g.<sup>15,41,44,45</sup>). In comparison, the light dependence of plastoquinone-9 and plastoeranol-8 levels is poorly documented. A few reports indicated that plastoquinone-9 accumulates in leaves exposed to high PFDs<sup>34,46-48</sup>. In contrast, the changes in plastoeranol-8 content of Arabidopsis leaves induced by high light stress were observed to be small compared to plastoquinone-9<sup>32,34,35,49</sup>. The present study shows that the plastoquinone-9 concentration in Arabidopsis leaves is extremely sensitive to the light conditions. Exposure to excess light energy by transferring low light-grown Arabidopsis plants to high light induced a dramatic reduction of the total plastoquinone-9 content while longer-term acclimation to high light was associated with a marked rise in the plastoquinone-9 levels (Fig. 1B). Moreover, those changes in plastoquinone-9 content appeared to follow the changes in PSII photochemical efficiency, with PSII photoinhibition being accompanied by a loss of plastoquinone-9 and PSII recovery being paralleled by increasing concentrations



**Figure 6. Tolerance of SPS1-overexpressing *Arabidopsis* plants to high light stress.** Plants grown for 5 weeks at a PFD of  $110 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  were exposed to high light at low temperature ( $1300 \mu\text{mol photons m}^{-2} \text{s}^{-1}$  and  $5^\circ\text{C}$ ) for 28 h. **A**) Lipid peroxidation monitored by autoluminescence imaging in WT (col 0) and SPS1oex #14 *Arabidopsis* plants after the light stress. **B**) Average autoluminescence intensity in WT and several SPS1oex lines before and after high light stress (white and black bars, respectively). Data are mean values of 5 measurements  $\pm$  SD. **C**) HOTE concentrations before and after high light stress (white and black bars, respectively). Data are mean values of 3 measurements  $\pm$  SD. **D**) Maximal quantum yield of PSII photochemistry (Fv/Fm) after high light stress before and after high light stress (white and black bars, respectively). Data are mean values of 10 measurements  $\pm$  SD. **E-F**) Changes in plastоquinone-9 (panel E) and plastochromanol-8 (panel F) in WT and the SPS1oex #3 line during high light stress. Data are mean values of 4 measurements  $\pm$  SD. In panels B, C and D, the stars indicate significant difference between WT and the SPS1oex lines at  $P < 0.05$  (Student's t-test).



**Figure 7. Comparison of the *vte1* and *vte2* mutants exposed to excess light energy.** WT (Col 2) and mutant plants were grown for 4 weeks at a PFD of  $170\text{-}\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Plants were then exposed to high light ( $1300\mu\text{mol m}^{-2} \text{ s}^{-1}$  at  $8^\circ\text{C}$ ). A-C) Prenyl lipid levels ( $\alpha$ -tocopherol [ $\alpha$ -Toc], total plastoquinone-9 [PQ], plastoehromanol-8 [PC]) before and after exposure for 3 or 7 d to high light at low temperature (white, grey and black bars, respectively). Data are mean values of 4 experiments  $\pm$  SD. Leaf specific weight (in  $\text{mg dry weight cm}^{-2}$ ) did not differ significantly between the genotypes so that expression of the data on a leaf area basis or on a weight basis does not affect the conclusions:  $1.467 \pm 0.075$  for WT,  $1.470 \pm 0.044$  for *vte1* and  $1.420 \pm 0.081$  for *vte2*. D) Lipid peroxidation monitored by autoluminescence imaging after 28 h in high light. E) HOTE concentrations before and after high light stress (white and black bars, respectively). Data are mean values of 3 measurements  $\pm$  SD. F) PSII photochemical efficiency ( $F_v/F_m$  chlorophyll fluorescence ratio) measured before and after high light stress (white and black bars, respectively). Data are mean values of 12 measurements  $\pm$  SD. Panels B, E, F: the stars indicate significant difference between WT and the mutants at  $P < 0.05$  (Student's t-test).



**Figure 8. Responses of VTE1-overexpressing *Arabidopsis* plants to excess light energy.** WT (Col 2) and two VTE1oex lines (#1 and 40) were grown for 5 weeks at a PFD of  $110 \mu\text{mol photons m}^{-2} \text{ s}^{-1}$ . Plants were then exposed to high light ( $1300 \mu\text{mol m}^{-2} \text{ s}^{-1}$  at  $5^\circ\text{C}$ ). A-C) Prenyl lipid levels under control conditions (before light stress): A)  $\gamma$ -tocopherol ( $\gamma$ -Toc, white bars) and  $\alpha$ -tocopherol ( $\alpha$ -Toc, grey bars); B) plastoquinone-9 (PQ); C) plastoehromanol-8 (PC). Data are mean values of 4 measurements  $\pm$  SD. The leaf specific weight (mg dry weight/cm<sup>2</sup>) did not differ significantly between the genotypes:  $1.1601 \pm 0.0980$  for Col 2,  $1.1673 \pm 0.1319$  for VTE1oex #40 and  $1.1673 \pm 0.1491$  for VTE1oex #1. D) Lipid peroxidation monitored by autoluminescence imaging after 28 h in high light. E) HOTE concentrations before and after high light stress (white and black bars, respectively). Data are mean values of 3 measurements  $\pm$  SD. F) PSII photochemical efficiency (Fv/Fm chlorophyll fluorescence ratio) before and after high light stress (white and black bars, respectively). Data are mean values of 12 measurements  $\pm$  SD.

of plastoquinone-9. This correlation with PSII activity was not found for plasto chromanol-8 and  $\alpha$ -tocopherol. Although the changes in plastoquinone-9 and plasto chromanol-8 concentrations induced by high light followed the same trends, the plasto chromanol changes were delayed and occurred with much lower amplitude relative to the plastoquinone changes (Fig. 1B). After transfer to high PFDs, the tocopherol content remained constant while PSII photochemistry was strongly inhibited. These changes were amplified in the  ${}^1\text{O}_2$ -overproducing *ch1* *Arabidopsis* mutant (Fig. 2), confirming the close correlation between PSII photochemical efficiency and plastoquinone-9 content under high light stress as well as the temporal disconnection between PSII photoinhibition and the variations in plasto chromanol-8 and tocopherol contents.

PSII photoinhibition is related to  ${}^1\text{O}_2$  formation resulting from the interaction between molecular oxygen and the triplet excited state of the reaction center chlorophyll molecule P680<sup>10</sup> and is linked to the degradation of the D1 protein triggered by the formed  ${}^1\text{O}_2$ <sup>50,51</sup>. The involvement of  ${}^1\text{O}_2$  in the damage of the PSII reaction centers can also be indirect by inhibiting synthesis of D1 and impairing the repair processes<sup>52</sup>. Elimination of  ${}^1\text{O}_2$  produced in the PSII centers is believed to be fulfilled by the  $\beta$ -carotene molecules located in the PSII reaction center<sup>53</sup> as well as by  $\alpha$ -tocopherol<sup>16,54,55</sup>. However, plastoquinone-9 has been demonstrated to be another potent antioxidant, which is able to quench  ${}^1\text{O}_2$  *in vitro*<sup>23,25,56</sup> and to inhibit lipid peroxidation in model systems<sup>24</sup>. Addition of plastoquinone homologues to *Chlamydomonas* cultures grown in the presence of plastoquinone biosynthesis inhibitors prevented degradation of D1<sup>55</sup>. Moreover, plastoquinone-9 and plasto chromanol-8 incorporated into liposomes were found to be more active than  $\alpha$ -tocopherol in the inhibition of  ${}^1\text{O}_2$ -induced lipid peroxidation<sup>24</sup>, and plastoquinone-9 was observed to be a better quencher of  ${}^1\text{O}_2$  than  $\alpha$ -tocopherol in solvents<sup>23</sup>. Thus, the dramatic loss of plastoquinone-9 observed in our study in *Arabidopsis* leaves suddenly exposed to excess light energy (Fig. 1B) is likely to be related to the  ${}^1\text{O}_2$ -scavenging activity of this compound which involves oxidation and consumption of the prenyl-lipidic molecule. Accordingly, increased production of  ${}^1\text{O}_2$  from the PSII centers in the *ch1* mutant was associated with an accelerated loss of plastoquinone-9. It is clear that exhaustion of the available pool of plastoquinones under excess light energy can have important consequences for the PSII repair cycle by precluding PSII reassembly. This phenomenon can exacerbate the inhibition of PSII photochemical activity. However, as shown here, exposure of *Arabidopsis* plants to high PFDs triggered up-regulation of the plastoquinone biosynthesis pathway, thus enhancing the capacity for plastoquinone-9 synthesis during plant acclimation to high light and compensating for the initial loss of plastoquinone-9 (Fig. 3B). This phenomenon led *in fine* to a strong accumulation of plastoquinone-9 in photoacclimated *Arabidopsis* plants.

Importantly, the present study shows that constitutive enhancement of the plastoquinone-9 biosynthesis capacity of *Arabidopsis* by overexpressing the *SPS1* gene strongly reinforces the tolerance of *Arabidopsis* plants to photooxidative stress (Fig. 6). The *SPS1oex* transgenic plants contain more plastoquinone-9 than WT plants both in low light and in high light conditions. Particularly, the high light stress-induced increase in plastoquinone-9 concentration following the initial reduction was much more rapid and reached much higher levels in *SPS1oex* plants compared to WT plants (Fig. 6E). Also, the plastoquinone-9 depletion in leaves immediately after transfer to high light was less pronounced in the *SPS1oex* plants relative to WT, leading to plastoquinone-9 contents close to the control values measured in WT plants before stress. The enhanced plastoquinone-9 levels in the *SPS1oex* lines throughout the high light stress treatments compared to WT were accompanied by a marked decrease in photooxidative damage and lipid peroxidation and in the preservation of the PSII photochemical activity. These findings show that active and efficient synthesis of plastoquinone-9 plays a role in the photoprotection mechanisms of plants. Incidentally, our results also suggest that solanesyl diphosphate synthesis is a limiting step in the plastoquinone biosynthesis pathway.

It should be stressed that, in addition to plastoquinone-9 accumulation, plasto chromanol-8 levels were also noticeably increased in the *SPS1oex* lines. Plasto chromanol-8 is synthesized from reduced plastoquinone-9 by *VTE1*<sup>34,35</sup>. Thus, the strong accumulation of plasto chromanol-8 in the *SPS1oex* lines suggests that chloroplasts can accommodate a finite amount of plastoquinone molecules, with the excess molecules being converted to plasto chromanol-8. The structure of plasto chromanol-8 is similar to that of tocotrienols and it has been shown *in vitro* that, like tocotrienols, it has antioxidant properties<sup>23,57</sup>. Plasto chromanol-8 has also been reported to play a crucial role in lipid protection in *Arabidopsis* seeds in combination with tocopherols<sup>58</sup>. Although the protective role of this compound in plant leaves is not yet established, plasto chromanol-8 accumulation could participate in the phototolerance of the *SPS1*-overexpressing plants. However, this idea is not consistent with the phenotype of *VTE1oex* transgenic plants which accumulated large amounts of plasto chromanol-8 with normal plastoquinone-9 levels and close to normal tocopherol levels: the latter plants exhibited the same tolerance towards photooxidative stress as WT (Fig. 8). Thus, high plasto chromanol-8 levels do not provide obvious advantages to leaves, at least under the light stress conditions used in our study, indicating that accumulation of plasto chromanols cannot explain by itself the increased phototolerance of the *SPS1oex* plants.

Interestingly, concomitant suppression of plasto chromanol-8 and tocopherols in the *vte1* mutant had a strong impact on the plastoquinone-9 levels which were substantially lowered compared to the WT levels both in low light and in high light (Fig. 7). Such an effect was not observed in the tocopherol-deficient *vte2* mutant that contains normal levels of plasto chromanol-8. Moreover, the decreased pool of plastoquinone-9 in the *vte1* mutant was accompanied by a much higher sensitivity to high light stress

compared to *vte2*. This difference between the *vte1* and *vte2* mutants can be understood if we consider that prenyl lipids can have overlapping antioxidant functions. In the absence of both tocopherols and plastoerol-8, photoprotection provided by the latter compounds relies now on plastoquinone-9. Due to the enhanced solicitation of plastoquinone-9 as an antioxidant, its concentration drops as it is consumed during  ${}^1\text{O}_2$  scavenging. Cooperation between antioxidants in protection against photosensitized oxidation is known for carotenoids and tocopherols<sup>59</sup>; a similar phenomenon could occur between prenyl lipids.

A substantial fraction of the prenyl lipids plastoquinone-9, plastoerol-8 and tocopherols is stored in the plastoglobules that are attached to the thylakoid membranes<sup>35,47,60</sup>. We found that, under the growth conditions used in this study, about 50% of plastoquinones are present in the plastoglobules of WT *Arabidopsis* leaves (Fig. 5). The plastoquinone molecules contained in the plastoglobules are not directly connected to the photosynthetic electron transport chain and therefore they represent the photochemically non-active fraction of plastoquinone. This non-photoactive pool was shown to represent the majority of plastoquinones in high light-grown *Arabidopsis* plants<sup>34</sup>. In several other plant species, increased concentrations of plastoquinone-9 (e.g. with aging or high PFDs) correlate with appearance of plastoglobules<sup>47</sup>. The same phenomenon appeared to occur in the SPS1oex plants in which the extra plastoquinones were mostly non-photoactive. Plastoglobules are permanently coupled to thylakoid membranes, indicating that prenyl lipids have the possibility to diffuse from plastoglobules to thylakoids<sup>61</sup>. Actually, the rapid consumption of about 80% of the plastoquinones observed in the present study in *Arabidopsis* plants suddenly exposed to excess light energy (Fig. 1B) supports the idea of plastoquinone exchange between plastoglobules and thylakoids. Since the majority of plastoquinone-9 is located in the plastoglobules, high light-induced degradation of such a high percentage of the plastoquinone content implies that plastoquinone molecules moved from the plastoglobules to the thylakoid-based photosystems where ROS are produced and can interact with them. In this context, it is also interesting to note that knockdown of *FIBBRILIN4* gene expression in apple disrupted the partitioning of plastoquinone between the plastoglobules and the rest of the chloroplast at the expenses of the plastoglobules<sup>62</sup> and increased concomitantly the sensitivity to biotic and abiotic stresses including high light<sup>63</sup>. Plastoglobular accumulation of plastoquinone-9 and rapid channeling of the stored plastoquinone-9 molecules to the thylakoids could thus function as a photoprotective mechanism by maintaining efficient  ${}^1\text{O}_2$ -scavenging activity under high light illumination and by providing new plastoquinone molecules for the PSII repair cycle. Stimulation of plastoquinone synthesis in SPS1oex plants is likely to enhance this mechanism, thus leading to more phototolerant plants as observed in this study.

In conclusion, it is becoming clear that the function of plastoquinones goes beyond their traditional roles as photosynthetic electron carriers between PSII and PSI and redox signals regulating cellular activities<sup>20,21</sup>. In fact, the partitioning of plastoquinone-9 between thylakoid membranes and their attached plastoglobules underlies the involvement of this compound in different physiological mechanisms. It has been recently shown that the plastoquinones contained in the plastoglobules are involved in electron-transfer reactions important for the chloroplast metabolism<sup>36</sup>. Other possible functions have recently emerged from a number of *in vitro* studies that have revealed that plastoquinone-9 possesses antioxidant properties<sup>23,24</sup>. Our finding that SPS1 overexpression in *Arabidopsis* concomitantly boosts the plastoquinone biosynthesis pathway and the plant tolerance to photooxidative stress demonstrates that this class of molecules does fulfill a protective role *in planta*.

## Methods

**Plant material.** WT *Arabidopsis* plants (*Arabidopsis thaliana*, ecotype Col 0 or Col 2) were grown at an air temperature of 20 °C/18 °C (day/night), a PFD of ~170  $\mu\text{mol}$  photons  $\text{m}^{-2} \text{s}^{-1}$  with a photoperiod of 8 h (unless specified otherwise) and a relative air humidity of 70%. The following mutants were used in this study: the tocopherol cyclase mutant *vte1*<sup>38</sup>, the homogentisate phytol transferase mutant *vte2*<sup>39</sup>, the  ${}^1\text{O}_2$ -producing mutant *ch1*<sup>28</sup>, and two VTE1 overexpressing lines, one in the *vte1* mutant background (VTE1oex #1) and the other in the WT background (VTE1oex #40)<sup>40</sup>. The corresponding WT (Col 0 or Col 2) of the different mutants are indicated in the figure legends.

Experiments were done on plants aged 4 or 5 weeks, as specified in the figure legends. Biochemical/biophysical analyses were performed on mature, well-developed leaves. High light stress was imposed by transferring plants to a high PFD (1300  $\mu\text{mol}$  photons  $\text{m}^{-2} \text{s}^{-1}$ ) at a low air temperature of 5 or 8 °C. Under those stress conditions, leaf temperature was ca. 17 or 19 °C respectively, in the temperature range of unstressed leaves. Because of its high photosensitivity<sup>28</sup>, the *ch1* mutant plants were submitted to less severe stress conditions (1000  $\mu\text{mol}$  photons  $\text{m}^{-2} \text{s}^{-1}$  at 10 °C).

**Prenyl lipids.** Leaf discs were grinded in ethyl acetate. After centrifugation, the supernatant was filtered, evaporated on ice under a stream of  $\text{N}_2$ , recovered in methanol/hexane (17/1) and filtered before analysis by HPLC, as described in<sup>64</sup> and<sup>34</sup>. The column was a Phenomenex Kinetex 2.6  $\mu\text{m}$ , 100  $\times$  4.6 mm, 100 Å. Separation of tocopherols, plastoquinone-9 and plastoerol-8 was done in the isocratic mode with methanol/hexane (17/1) as solvent system and a flow rate of 0.8 ml/min. All prenyl lipids, except oxidized plastoquinone-9, were detected by their fluorescence at 330 nm with an excitation at 290 nm. Plastoquinone-9 in the oxidized state was measured by its absorbance at 255 nm. Plastoerol-8

and plastoquinone-9 standards were a kind gift from Dr. J. Kruk. Tocopherol standards were purchased from Sigma.

Leaf samples were frozen in liquid nitrogen as quickly as possible to avoid changes in the plastoquinone redox state during sample preparation. Determination of the photoactive and non-active fractions of plastoquinone-9 was done following the protocol described in<sup>64</sup>. Leaves were cut in half. To determine the pool of photoactive plastoquinone-9, we first measured the amount of reduced plastoquinone-9 (plastoquinol-9) in one half of the leaves after a dark-adaptation period of 2 h. The dark-adapted samples were frozen in liquid nitrogen after 3-s illumination with far-red light to ensure full oxidation of the photoactive pool of plastoquinone. Then, plastoquinol-9 was re-quantified in the other half of the leaves that had been exposed to a high PFD of white light ( $2000\mu\text{mol photons m}^{-2}\text{ s}^{-1}$ ) for 15 s. Illumination was done in a mortar that was filled with liquid nitrogen after the 15-s illumination in order to freeze the samples in the light. The size of the photoactive pool was calculated as the difference between the plastoquinol-9 content in the light and the plastoquinol-9 content in the dark. The size of the non-photoactive pool of plastoquinone-9 was the sum of the amount of plastoquinol-9 in the dark (not reoxidizable in the dark) and the amount of oxidized plastoquinone-9 in the light (not reducible by high light). This pool corresponds mainly to the plastoquinone molecules present in the plastoglobules although a small fraction can also be present in the chloroplast envelope<sup>47</sup>.

**Chlorophyll fluorescence.** Chlorophyll fluorescence was measured in attached leaves with a PAM-2000 chlorophyll fluorometer (Walz), as described in<sup>15</sup>. The maximal quantum yield of PSII photochemistry was measured after a period of dark adaptation as  $\text{Fv}/\text{Fm} = (\text{Fm}-\text{Fo})/\text{Fm}$  where Fo is the initial fluorescence level (measured with a weak, non-actinic red light) and Fm is the maximal fluorescence level (measured with a 800-ms pulse of saturating white light). The actual quantum yield of PSII photochemistry was measured in the light as  $\Delta\text{F}/\text{Fm}' = (\text{Fm}'-\text{Fs})/\text{Fm}$  where Fs is the steady-state fluorescence level and Fm' is the maximal fluorescence level.  $\Delta\text{F}/\text{Fm}'$  was measured in leaves adapted to different PFDs of white light.

**RNA Isolation and Quantitative RT-PCR.** The NucleoSpin RNA plant kit (Macherey-Nagel) was used to extract total RNA from 100 mg of leaves. RNA concentrations were measured on a NanoDrop2000 (Thermo Scientific, USA). Quantitative RT-PCR (qRT-PCR) was performed with cDNA synthesized with the Prime Script<sup>TM</sup> Reverse Transcriptase (Takara) from 500 ng of total RNA. Design of the specific primers was done for each gene (see supplemental Table 1) with the Primer3plus software ([www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi](http://www.bioinformatics.nl/cgi-bin/primer3plus/primer3plus.cgi)). qRT-PCR was carried out using the LightCycler 480 SYBR Green I Master (Roche) in the quantitative PCR thermal cycler (LightCycler 480 real-time PCR system; Roche). Reaction preparation (total volume of 5  $\mu\text{l}$ ): 2  $\mu\text{l}$  of cDNA diluted 10-fold, 2  $\mu\text{l}$  of SYBR Green I Master, and 1  $\mu\text{M}$  forward and reverse primers. The amplification profile was 95 °C for 10 min and 45 cycles (95 °C/15 s, 60 °C or 56 °C (depending on the gene)/15 s, and 72 °C/15 s). Reactions were performed in triplicates. Normalization of gene expressions was done using ACTIN2 as housekeeping gene. At least three biological replicates were performed for each gene tested.

**SPS1 overexpressing transgenic *Arabidopsis* plants.** To generate plants over-expressing *SPS1* (At1g78510), full-length cDNA was amplified using the primers allowing the addition of attB recombination sites. The cDNA was cloned into a pDONR201 vector (Life Technologies) and transferred into the binary GATEWAY destination vector pB2GW7,0 (Plant Systems Biology, VIB-Ghent University, Belgium)<sup>65</sup>. The pB2GW7,0 vector allows expression of a sense cDNA under the control of the cauliflower mosaic virus 35S promoter and includes a glufosinate-resistance gene. Transformation using *Agrobacterium tumefaciens* C58 strain was performed as described by Davis *et al.*<sup>66</sup>. Homozygous transgenic lines (T3) were produced and selected from resistance segregation assays.

Transformed lines were verified using PCR analysis on genomic DNA with the Phire Plant Direct PCR kit (Finnzymes) and gene-specific primers. ACTIN2 was used as a positive control for each PCR. The PCR program was 98 °C for 5 min, 30 cycles (denaturation at 95 °C for 5 s, annealing at 55 °C for 5 s, extension at 72 °C for 20 s) (GeneAmp, PCRSystem 2700, Applied Biosystems). For RT-PCR analysis, RNA extraction from leaves was done using NucleoSpin RNA Plant Kit (Macherey Nagel) according to the manufacturer's instructions. Reverse transcription was performed using total RNA (500 ng), a Superscript III Reverse Transcriptase (Life Technologies) and an oligo(dT)20 primer. RT-PCR assays were performed using the same gene-specific primers used for genotyping. Primers: *SPS1*-F, tggacgcctgccttactct; *SPS1*-R, gcaagcttatacacgacg; *ACTIN2*-F, aaaaatggccgatggtgaggatat; *ACTIN2*-R, caataccggtgtacgaccact.

**Lipid peroxidation analysis and imaging.** 0.5 g plant material (fresh weight) was grinded with Ultraturax T25 (Ika-Werk) in 2.5 ml  $\text{CHCl}_3$ -methanol (50-50) containing 5 mM triphenyl phosphine, 1 mM butylated hydroxytoluene and 1 ml citric acid (1M), with 15-hydroxy-11,13(*Z,E*)-eicosadienoic acid as internal standard. After evaporation under a stream of  $\text{N}_2$ , the organic phase was re-solubilized in 1.25 ml ethanol and 1.25 ml 3.5 M NaOH. The sample was hydrolyzed at 80 °C for 30 min. pH was adjusted between 4 and 5 by addition of 1 M citric acid, and hydroxy fatty acids were then extracted with hexane/ether (50/50). HOTE isomers (hydroxy octadecatrienoic acid, produced by the oxidation of linolenic acid) were separated and quantified by straight phase HPLC-UV analysis, as described in a previous

work<sup>67</sup>. Estimation of ROS-induced HOTEs and lipoxygenase-induced HOTEs was done according to the method described in<sup>67</sup>.

Lipid peroxidation was imaged by measuring autoluminescence emission. The latter signal is attributed to the spontaneous decomposition of lipid peroxides<sup>68</sup>. Spontaneous photon emission from whole plants was measured after a 2-h dark adaptation period using a cooled charge-coupled device (CCD) camera, as explained in<sup>68</sup>. Integration time was 20 min and pixel binning was 2 × 2. The images were analyzed with the ImageJ software.

## References

- Apel, K. & Hirt, H. Reactive oxygen species: metabolism, oxidative stress, and signal transduction. *Annu. Rev. Plant Biol.* **55**, 373–399 (2004).
- Asada, K. Production and scavenging of reactive oxygen species in chloroplasts and their functions. *Plant Physiol.* **141**, 391–396 (2006).
- Li, Z., Wakao, S., Fischer, B. B. & Niyogi, K. K. Sensing and responding to excess light. *Annu. Rev. Plant Biol.* **60**, 239–260 (2009).
- Edge, R., McGarvey, D. J. & Truscott, T. G. The carotenoids as anti-oxidants – a review. *J. Photochem. Photobiol. B: Biol.* **41**, 189–200 (1997).
- Noctor, G. & Foyer, C. H. Ascorbate and glutathione: keeping active oxygen species under control. *Annu. Rev. Plant Physiol. Plant Mol. Biol.* **49**, 249–279 (1998).
- Falk, J. & Munné-Bosch, S. Tocochromanol functions in plants: antioxidation and beyond. *J. Exp. Bot.* **61**, 1549–1566 (2010).
- Tarrago, L., Laugier, E. & Rey, P. Protein-repairing methionine sulfoxide reductases in photosynthetic organisms: gene organization, reduction mechanisms, and physiological roles. *Mol. Plant* **2**, 202–217 (2009).
- Tripathi, B. N., Bhatt, I. & Dietz, K. J. Peroxiredoxins: a less studied component of hydrogen peroxide detoxification in photosynthetic organisms. *Protoplasma* **235**, 2–15 (2009).
- Pospisil, P. The role of metals in production and scavenging of reactive oxygen species in photosystem II. *Plant Cell Physiol.* **55**, 1224–1232 (2014).
- Krieger-Liszka, A. Singlet oxygen production in photosynthesis. *J. Exp. Bot.* **56**, 337–346 (2005).
- Triantaphylidès, C. & Havaux, M. Singlet oxygen in plants: Production, detoxification and signalling. *Trends Plant Sci.* **14**, 219–228 (2009).
- Gonzalez-Perez, S. *et al.* Early transcriptional defense responses in *Arabidopsis* cell suspension culture under high-light conditions. *Plant Physiol.* **156**, 1439–1456 (2011).
- Triantaphylidès, C. *et al.* Singlet oxygen is the major reactive oxygen species involved in photooxidative damage to plants. *Plant Physiol.* **148**, 960–968 (2008).
- Frank, H. A. & Cogdell, R. J. Carotenoids in photosynthesis. *Photochem. Photobiol.* **63**, 257–264 (1996).
- Havaux, M., Eymery, F., Porfiriova, S., Rey, P. & Dörrmann, P. Vitamin E protects against photoinhibition and photooxidative stress in *Arabidopsis thaliana*. *Plant Cell* **17**, 3451–3469 (2005).
- Krieger-Liszka, A. & Trebst, A. Tocopherol is the scavenger of singlet oxygen produced by the triplet states of chlorophyll in the PSII reaction centre. *J. Exp. Bot.* **57**, 1677–1684 (2006).
- Dörrmann, P. Functional diversity of tocochromanols in plants. *Planta* **225**, 269–276 (2007).
- Mène-Saffrané, L. & Dellapenna, D. Biosynthesis, regulation and functions of tocochromanols in plants. *Plant Physiol. Biochem.* **48**, 301–309 (2010).
- Amesz, J. The function of plastoquinone in photosynthetic electron transport. *Biochim. Biophys. Acta* **301**, 35–51 (1973).
- Pfannschmidt, T. *et al.* Potential regulation of gene expression in photosynthetic cells by redox and energy state: approaches towards better understanding. *Ann. Bot.* **103**, 599–607 (2009).
- Rochaix, J. D. Redox regulation of thylakoid protein kinases and photosynthetic gene expression. *Antioxid. Redox Signal.* **18**, 2184–2201 (2014).
- Samson, G. & Bruce, D. Origins of the low yield of chlorophyll a fluorescence induced by single turnover flash in spinach thylakoid. *Biochim. Biophys. Acta* **1276**, 147–153 (1996).
- Gruszká, J., Pawłak, A. & Kruk, J. Tocochromanols, plastoquinol, and other biological prenyllipids as singlet oxygen quenchers – determination of singlet oxygen rate constants and oxidation products. *Free Rad. Biol. Med.* **45**, 920–928 (2008).
- Nowicka, B., Gruszká, J. & Kruk, J. Function of plastochemical and other biological prenyllipids in the inhibition of lipid peroxidation – A comparative study in model systems. *Biochim. Biophys. Acta* **1828**, 233–240 (2013).
- Yadav, D. K., Kruk, J., Sinha, R. K. & Pospisil, P. Singlet oxygen scavenging activity of plastoquinol in photosystem II of high plants: Electron paramagnetic resonance spin-trapping study. *Biochim. Biophys. Acta* **1797**, 1807–1811 (2010).
- Melis, A. & Brown, J. S. Stoichiometry of system I and system II reaction centers and of plastoquinone in different photosynthetic membranes. *Proc. Natl. Acad. Sci. USA* **77**, 4712–4716 (1980).
- Savitch, L. V., Maxwell, D. P. & Huner, P. A. Photosystem II excitation pressure and photosynthetic carbon metabolism in *Chlorella vulgaris*. *Plant Physiol.* **111**, 127–136 (1996).
- Ramel, F. *et al.* Light-induced acclimation of the *Arabidopsis chlorina1* mutant to singlet oxygen. *Plant Cell* **25**, 1445–1462 (2013).
- Cheng, Z. *et al.* Highly divergent methyltransferases catalyze a conserved reaction in tocopherol and plastoquinone synthesis in cyanobacteria and photosynthetic eukaryotes. *Plant Cell* **15**, 2343–2356 (2003).
- Jun, L., Saiki, R., Tatsumi, K., Nakagawa, T. & Kawamukai, M. Identification and subcellular localization of two solanesyl diphosphate synthases from *Arabidopsis thaliana*. *Plant Cell Physiol.* **45**, 1882–1888 (2004).
- Ducluzeau, A.-L. *et al.* Gene networking reconstruction identifies the authentic trans-prenyl diphosphate synthase that makes the solanesyl moiety of ubiquinone-9 in *Arabidopsis*. *Plant J.* **69**, 366–375 (2012).
- Block, A. *et al.* Functional modeling identifies paralogous solanesyl-diphosphate synthases that assemble the side chain of plastoquinone-9 in plastids. *J. Biol. Chem.* **288**, 27594–27606 (2013).
- Hirooka K *et al.* Functional analysis of two solanesyl diphosphate synthases from *Arabidopsis thaliana*. *Biosci. Biotechnol. Biochem.* **69**, 592–601 (2005).
- Szymanska, R. & Kruk, J. Plastoquinol is the main prenyl lipid synthesized during acclimation to high light conditions in *Arabidopsis* and is converted to plastochemical by tocopherol cyclase. *Plant Cell Physiol.* **51**, 537–545 (2010).
- Zbierzak, A. M. *et al.* Intersection of the tocopherol and plastoquinone metabolic pathways at the plastoglobule. *Biochem. J.* **425**, 389–399 (2009).
- Molina Torres, J., Laidman, D. L. & Gaunt, J. K. The influence of photoperiod on incorporation of precursors into tocopherols and plastoquinone in *Xanthium strumarium* L. *New Phytol.* **111**, 397–401 (1989).
- Sadre, R., Grulber, J. & Frentzen, M. Characterization of homogentisate prenyltransferases involved in plastoquinone-9 and tocochromanol biosynthesis. *FEBS Lett.* **580**, 5357–5362 (2006).

38. Porfirova, S., Bergmuller, E., Tropf, S., Lemke, R. & Dörmann, P. Isolation of an *Arabidopsis* mutant lacking vitamin E and identification of a cyclase essential for all tocopherol biosynthesis. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 12495–12500 (2002).
39. Sattler, S. E., Gilliland, L. U., Magallanes-Lundback, M., Pollard, M. & DellaPenna, D. Vitamin E is essential for seed longevity and for preventing lipid peroxidation during germination. *Plant Cell* **16**, 1419–1432 (2004).
40. Kanwischer, M., Porfirova, S., Bergmüller, E. & Dörmann, P. Alterations in tocopherol cyclase activity in transgenic and mutant plants of *Arabidopsis* affect tocopherol content, tocopherol composition, and oxidative stress. *Plant Physiol.* **137**, 713–723 (2005).
41. Maeda H., Song, W., Sage, T. L. & DellaPenna, D. Tocopherols play a crucial role in low-temperature adaptation and phloem loading in *Arabidopsis*. *Plant Cell* **18**, 2710–2731 (2006).
42. Abbasi, A.-R., Hajirezaei, M., Hofius, D., Sonnewald, U. & Voll, L. M. Specific roles of  $\alpha$ - and  $\gamma$ -tocopherol in abiotic stress responses of transgenic tobacco. *Plant Physiol.* **143**, 1720–1738 (2007).
43. Szymanska, R. & Kruk, J.  $\gamma$ -Tocopherol dominates in young leaves of runner bean (*Phaseolus coccineus*) under a variety of growing conditions: the possible functions of gamma-tocopherol. *Phytochemistry* **69**, 2142–2148 (2008).
44. Logan, B. A., Demmig-Adams, B. & Adams, W. W. III & Grace, S. C. Antioxidants and xanthophyll cycle-dependent energy dissipation in *Cucurbita pepo* L. and *Vinca major* L. acclimated to four growth PPFs in the field. *J. Exp. Bot.* **49**, 1869–1879 (1998).
45. Golan, T., Muller-Moulé, P. & Niyogi, K. K. Photoprotection mutants of *Arabidopsis thaliana* acclimate to high light by increasing photosynthesis and specific antioxidants. *Plant Cell Environ.* **29**, 879–887 (2008).
46. Lichtenthaler, H. K. in *Lipids and Lipid Polymers in Higher Plants* (eds Tevini, M. & Lichtenthaler, H. K.) 231–258 (Springer, 1977).
47. Lichtenthaler, H. K. Biosynthesis, accumulation and emission of carotenoids,  $\alpha$ -tocopherol, plastoquinone, and isoprene in leaves under high photosynthetic irradiance. *Photosynth. Res.* **92**, 163–179 (2007).
48. Kruk, J. & Trebst, A. Plastoquinol as a singlet oxygen scavenger in photosystem II. *Biochim. Biophys. Acta* **1777**, 154–162 (2008).
49. Rastogi, A. *et al.* Singlet oxygen scavenging activity of tocopherol and plastochemical in *Arabidopsis thaliana*: relevance to photooxidative stress. *Plant Cell Environ.* **37**, 392–401 (2014).
50. Miyao, M. Involvement of active oxygen species in degradation of the D1 protein under strong illumination in isolated supercomplexes of photosystem II. *Biochemistry* **33**, 9722–9730 (1994).
51. Adir, N., Zer, H., Shochat, S. & Ohad, I. Photoinhibition - an historical perspective. *Photosynth. Res.* **76**, 343–370 (2003).
52. Nishiyama, Y., Allakverdiev, S. I. & Murata, N. A new paradigm for the action of reactive oxygen species in the photoinhibition of photosystem II. *Biochim. Biophys. Acta* **1757**, 742–749 (2006).
53. Telfer, A. What is beta-carotene doing in the photosystem II reaction centre? *Philos. Trans. R. Soc. Lond. B. Biol. Sci.* **357**, 1431–1439 (2002).
54. Trebst, A., Depka, B. & Hollander-Czytko, H. A specific role for tocopherol and of chemical singlet oxygen quenchers in the maintenance of photosystem II structure and function in *Chlamydomonas reinhardtii*. *FEBS Lett.* **516**, 156–160 (2002).
55. Kruk, J., Hollander-Czytko, H., Oettmeier, W. & Trebst, A. Tocopherol as singlet oxygen scavenger in photosystem II. *J. Plant Physiol.* **162**, 749–757 (2005).
56. Hundal, T., Forsmark-Andree, P., Ernster, L. & Andersson, B. Antioxidant activity of reduced plastoquinone in chloroplast thylakoid membrane. *Arch. Biochem. Biophys.* **324**, 117–122 (1995).
57. Olejnik, D., Gogolewski, M. & Nogala-Kaucka, M. Isolation and some properties of plastochemical-8. *Nahrung* **41**, 101–104 (1997).
58. Mène-Saffrané, L., Jones, A. D. & DellaPenna, D. Plastochemical-8 and tocopherols are essential lipid-soluble antioxidants during seed desiccation and quiescence in *Arabidopsis*. *Proc. Natl. Acad. Sci. USA* **107**, 17815–17820 (2010).
59. Wrona, M., Kortykowski, W., Rozanowska, M., Sarna, T. & Truscott, T. G. Cooperation of antioxidants in protection against photosensitized oxidation. *Free Rad. Biol. Med.* **35**, 1319–1329 (2003).
60. Steinmüller, D. & Tevini, M. Composition and function of plastoglobuli. *Planta* **163**, 201–207 (1985).
61. Austin, J. R. 2nd, Frost, E., Vidi, P. A., Kessler, F. & Staehelin, P. A. Plastoglobules are lipoprotein subcompartments of the chloroplast that are permanently coupled to thylakoid membranes and contain biosynthetic enzymes. *Plant Cell* **18**, 1693–1703 (2006).
62. Singh, D. K., Laremore, T. N., Smith, P. B., Maximova, S. N. & McNellis, T. W. Knockdown of FIBBRILIN4 gene expression in apple decreases plastoglobule plastoquinone content. *PLOS One* **7**, e47547 (2012).
63. Singh, D. K. *et al.* FIBBRILIN4 is required for plastoglobule development and stress resistance in apple and *Arabidopsis*. *Plant Physiol.* **154**, 1281–1293 (2010).
64. Kruk, J. & Karpinski, S. An HPLC-based method of estimation of the total redox state of plastoquinone in chloroplasts, the size of the photochemically active plastoquinone-pool and its redox state in thylakoids of *Arabidopsis*. *Biochim. Biophys. Acta* **1757**, 1669–1675 (2006).
65. Karimi, M., Inze, D. & Depicker, A. GATEWAY vectors for *Agrobacterium*-mediated plant transformation. *Trends Plant Sci.* **7**, 193–195 (2002).
66. Davis, A. M., Hall, A., Millar, A. J., Darrah, C. & Davis, S. J. Protocol: Streamlined sub-protocols for floral-dip transformation and selection of transformants in *Arabidopsis thaliana*. *Plant Methods* **5**, 3 (2009).
67. Montillet, J. L. *et al.* The upstream oxylipin profile of *Arabidopsis thaliana*: A tool to scan for oxidative stresses. *Plant J.* **40**, 439–451 (2004).
68. Birtic, S. *et al.* Using spontaneous photon emission to image lipid oxidation patterns in plant tissues. *Plant J.* **67**, 1103–1115 (2011).

## Acknowledgments

VTE10ex *Arabidopsis* seeds were a kind gift from Peter Doermann (University of Bonn, Germany) and the plastoquinone-9 and plastochemical-8 standards were obtained from Jerzy Kruk (Jagiellonian University, Krakow, Poland). They are gratefully thanked for their help. We would also like to thank the members of the GRAP platform (CEA/Cadarache) for their help in growing and stressing the plants.

## Author Contributions

B.K., N.B. and M.H. conceived and designed the experiments. B.K., N.B. and A.C. performed the experiments. B.K. and M.H. analyzed the data. M.H. wrote the main manuscript text.

## Additional Information

Supplementary information accompanies this paper at <http://www.nature.com/srep>

**Competing financial interests:** The authors declare no competing financial interests.

**How to cite this article:** Ksas, B. *et al.* Plant tolerance to excess light energy and photooxidative damage relies on plastoquinone biosynthesis. *Sci. Rep.* **5**, 10919; doi: 10.1038/srep10919 (2015).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>