

SCIENTIFIC REPORTS



OPEN

Quantum coherence of steered states

Xueyuan Hu¹, Antony Milne², Boyang Zhang¹ & Heng Fan³

Received: 08 September 2015

Accepted: 11 December 2015

Published: 19 January 2016

Lying at the heart of quantum mechanics, coherence has recently been studied as a key resource in quantum information theory. Quantum steering, a fundamental notion originally considered by Schrödinger, has also recently received much attention. When Alice and Bob share a correlated quantum system, Alice can perform a local measurement to 'steer' Bob's reduced state. We introduce the maximal steered coherence as a measure describing the extent to which steering can remotely create coherence; more precisely, we find the maximal coherence of Bob's steered state in the eigenbasis of his original reduced state, where maximization is performed over all positive-operator valued measurements for Alice. We prove that maximal steered coherence vanishes for quantum-classical states whilst reaching a maximum for pure entangled states with full Schmidt rank. Although invariant under local unitary operations, maximal steered coherence may be increased when Bob performs a channel. For a two-qubit state we find that Bob's channel can increase maximal steered coherence if and only if it is neither unital nor semi-classical, which coincides with the condition for increasing discord. Our results show that the power of steering for coherence generation, though related to discord, is distinct from existing measures of quantum correlation.

Quantum coherence, originating from the quantum pure state superposition principle, is one of the most fundamental properties of quantum mechanics. It is increasingly recognized as a vital resource in a range of scenarios, including quantum reference frames^{1–3}, transport in biological systems^{4–6} and quantum thermodynamics^{7–9}. How to measure coherence is an essential problem in both quantum theory and quantum information and has recently attracted much attention^{10–14}. The quantification of coherence in a single quantum system depends on both the quantum state and a fixed basis for the density matrix of the system^{10,11}. The fixed basis is usually chosen as the eigenbasis of the Hamiltonian or another observable. In either case, the quantified coherence is not an intrinsic property of the single-party quantum state itself. The dynamics of quantum coherence under certain noisy channels has also attracted a lot of research attention^{15,16} and is connected to the dynamics of quantum correlations¹⁷.

When Alice and Bob share a correlated quantum system, a measurement by Alice can 'steer' the quantum state of Bob. Quantum steering, especially Einstein-Podolsky-Rosen (EPR) steering, has long been noted as a distinct nonlocal quantum effect¹⁸ and has attracted recent research interest both theoretically and experimentally^{19–22}. The quantum steering ellipsoid (QSE)^{23–27}, defined as the whole set of Bloch vectors to which Bob's qubit can be steered by a positive-operator valued measurement (POVM) on Alice's qubit, provides a faithful geometric presentation for two-qubit states. Using the QSE formalism we have studied a class of two-qubit states whose quantum discord can be increased by local operations²⁸. Interestingly, arbitrarily small mutual information is sufficient for the QSE of a pure two-qubit state to be the whole Bloch ball. Since mutual information is an upper bound of quantum correlation measures such as entanglement and discord, the power that one qubit has to steer another cannot be fully characterized by the quantum correlation between the two qubits. A measure that quantifies the power of generating quantum coherence by steering is therefore necessary.

In this paper we consider a bipartite quantum state ρ with non-degenerate reduced state ρ_B and study the coherence of Bob's steered state, which is obtained by Alice's POVM. Here the eigenbasis of ρ_B is employed as the fixed basis in which to calculate the coherence of the steered state. The significance of this choice of basis is that Bob's initial state is incoherent. When Alice performs a local measurement, she can steer Bob's state to one that is coherent in the eigenbasis of ρ_B , i.e. Alice generates Bob's coherence. By $\mathcal{C}(\rho)$ we denote the maximum coherence that Alice can generate through local measurement and classical communication. In contrast to existing quantifiers of coherence, \mathcal{C} is an intrinsic property of the bipartite quantum state ρ , because the reference basis of coherence

¹School of Information Science and Engineering, and Shandong Provincial Key Laboratory of Laser Technology and Application, Shandong University, Jinan, 250100, P. R. China. ²Controlled Quantum Dynamics Theory, Department of Physics, Imperial College London, London SW7 2AZ, UK. ³Institute of Physics, Chinese Academy of Sciences, Beijing 100190, China. Correspondence and requests for materials should be addressed to X.H. (email: xyhu@sdu.edu.cn)

ence, chosen as the eigenbasis of ρ_B , is inherent to the bipartite state. Furthermore, we find that \mathcal{C} gives a different ordering of states compared to quantum entanglement or discord; this indicates that \mathcal{C} describes remote quantum properties distinct from these measures of quantum correlation. Properties of \mathcal{C} are also studied. The maximal steered coherence is found to vanish only for classical states and can be created and increased by local quantum channels. Given that coherence plays a central role in a diverse range of quantum information processing tasks, we can also consider how steered coherence might be used as a resource. We close our discussion by presenting one such scenario.

We note that, shortly after this paper first appeared, Mondal *et al.* presented a study on the steerability of local quantum coherence²⁹. We consider our works to be complementary: though examining a similar topic, our approaches are very different (Mondal *et al.* consider steering from the existence of a local hidden state model rather than from the perspective of the QSE formalism).

Results

Definition. We consider a bipartite quantum state ρ , where the reduced state ρ_B is non-degenerate with eigenstates $\Xi = \{|\xi_i\rangle\}$. When Alice obtains the POVM element M as a measurement outcome, Bob's state is steered to $\rho_B^M := \text{tr}_A(M \otimes \mathbb{1}\rho)/p_M$ with probability $p_M := \text{tr}(M \otimes \mathbb{1}\rho)$, where $\mathbb{1}$ denotes the single qubit identity operator. Baumgratz *et al.*¹⁰ gives the the quantum coherence C of ρ_B^M in the basis $\{|\xi_i\rangle\}$ as the summation of the absolute values of off-diagonal elements:

$$C(\rho_B^M, \{|\xi_i\rangle\}) = \frac{1}{p_M} \sum_{i \neq j} |\langle \xi_i | \text{tr}_A(M \otimes \mathbb{1}\rho) |\xi_j \rangle|. \quad (1)$$

Here we maximize the coherence $C(\rho_B^M, \{|\xi_i\rangle\})$ over all possible POVM operators M and define the maximal steered coherence as

$$\mathcal{C}(\rho) := \max_{M \in \text{POVM}} \left[\frac{1}{p_M} \sum_{i \neq j} |\langle \xi_i | \text{tr}_A(M \otimes \mathbb{1}\rho) |\xi_j \rangle| \right]. \quad (2)$$

When ρ_B is degenerate, Ξ is not uniquely defined; however, we can take the infimum over all possible eigenbases for Bob and define the maximal steered coherence as

$$\mathcal{C}(\rho) := \inf_{\Xi} \left\{ \max_{M \in \text{POVM}} \left[\frac{1}{p_M} \sum_{i \neq j} |\langle \xi_i | \text{tr}_A(M \otimes \mathbb{1}\rho) |\xi_j \rangle| \right] \right\}. \quad (3)$$

It is worth noting that \mathcal{C} is an intrinsic property of the bipartite quantum state ρ . When fixing the basis in which to calculate the coherence, we need not choose an observable that is independent of the state; the basis $\{|\xi_i\rangle\}$ we choose here is inherent to the state ρ .

Properties. We prove that the following important properties hold for maximal steered coherence.

(E1) \mathcal{C} vanishes if and only if ρ is a classical state (zero discord for Bob), i.e. can be written as

$$\rho = \sum_{i=1}^d p_i \rho_i^A \otimes |\xi_i\rangle \langle \xi_i|. \quad (4)$$

The proof of this is given in Methods.

(E2) \mathcal{C} reaches a maximum for all pure entangled states with full Schmidt rank, i.e. states that can be written in as $|\Psi\rangle = \sum_{i=1}^{d_B} \lambda_i |\phi_i^A\rangle \otimes |\xi_i^B\rangle$ with $\lambda_i \neq 0 \forall i$. Here d_B is the dimension of Bob's state. For a single quantum system of dimension d_B , the maximally coherent state is $|\psi_m^B\rangle = \frac{1}{\sqrt{d_B}} \sum_{i=1}^{d_B} |xi_i^B\rangle$ ¹⁰; Bob is steered to this when Alice obtains the measurement outcome $M = |\psi^A\rangle \langle \psi^A|$, where $|\psi^A\rangle$ is the state $\sum_{i=1}^{d_B} \frac{1}{\lambda_i} |\phi_i^A\rangle$ after normalisation.

(E3) \mathcal{C} is invariant under local unitary operations. When the unitary operator $U = U_A \otimes U_B$ acts on a bipartite state ρ , the eigenbasis of ρ_B is rotated by U_B , so that the off-diagonal elements of ρ_B^M become

$$\langle \xi_i | U_B^\dagger \text{tr}_A[M \otimes \mathbb{1}(U\rho U^\dagger)] U_B |\xi_j \rangle = \langle \xi_i | \text{tr}_A[U_A^\dagger M U_A \otimes \mathbb{1}\rho] |\xi_j \rangle. \quad (5)$$

From Eq. (2) it is clear that $\mathcal{C}(U\rho U^\dagger) = \mathcal{C}(\rho)$.

(E4) \mathcal{C} can be increased by Bob performing a local quantum channel prior to Alice's steering. Property (E4) holds owing to the fact that a local channel Λ_B , under certain conditions³⁰, can transform a classical state with vanishing \mathcal{C} into a discordant state with strictly positive \mathcal{C} . Note, however, that a channel Λ_A performed by Alice prior to steering cannot increase \mathcal{C} . (This follows because Λ_A can be performed by applying a unitary operation to A and an ancilla A' and then discarding A' ; the unitary operation does not affect the set of Bob's steered states, while discarding A' may limit Alice's ability to steer Bob's state. Thus Λ_A performed by Alice does not alter Bob's reduced state ρ_B but shrinks the set of his steered states; such a channel cannot increase \mathcal{C} .)

Let us also note an important consequence of property (E2): \mathcal{C} is distinct from the entanglement E ³¹ and discord-type quantum correlations \mathcal{D} ³². In fact, \mathcal{C} gives a different ordering of states from E or \mathcal{D} . We demonstrate this by considering states $\rho_1 = |\Psi_1\rangle \langle \Psi_1|$ and $\rho_2 = (1 - \delta)|\Psi_2\rangle \langle \Psi_2| + \delta \frac{\mathbb{1}_d \otimes \mathbb{1}_d}{d^2}$, where

$|\Psi_1\rangle = \sqrt{1 - (d-1)\delta}|00\rangle + \sqrt{\delta}\sum_{j=1}^{d-1}|jj\rangle$ and $|\Psi_2\rangle = \frac{1}{\sqrt{d}}\sum_{j=0}^{d-1}|jj\rangle$ are both pure entangled states with full Schmidt rank of dimension d , and $0 < \delta \ll 1$. According to (E2), $\mathcal{C}(\rho_1)$ reaches the maximum; whereas for ρ_2 , Bob's steered state is always mixed and hence not maximally coherent state in any given basis. We therefore have $\mathcal{C}(\rho_1) > \mathcal{C}(\rho_2)$. Meanwhile, $\mathcal{D}(\rho_1)$ and $E(\rho_1)$ can be made arbitrarily small by taking δ to be small enough, whilst $\mathcal{D}(\rho_2)$ and $E(\rho_2)$ approach 1 for small δ . Hence δ exists such that $\mathcal{D}(\rho_1) < \mathcal{D}(\rho_2)$ and $E(\rho_1) < E(\rho_2)$.

General expression for two-qubit states. We now derive the general form of \mathcal{C} for two-qubit states. The state of a single qubit can be written as $\rho_B = \frac{1}{2}\sum_{i=0}^3 b_i \sigma_i$, where $\sigma_0 = \mathbf{1}$, σ_i with $i = 1, 2, 3$ are Pauli matrices, $b_i = \text{tr}(\rho_B \sigma_i)$, and $\mathbf{b} = (b_1, b_2, b_3)^T$ is Bob's Bloch vector. The norm of the vector \mathbf{b} is denoted by b . The quantum coherence of ρ_B in a given basis $\{\mathbf{n}, -\mathbf{n}\}$, where $|\mathbf{n}| = 1$, is

$$C(\rho_B, \mathbf{n}) = |\mathbf{b} \times \mathbf{n}|. \quad (6)$$

Let B and N be the points associated with the vectors \mathbf{b} and \mathbf{n} respectively, and let O be the origin. Since $\frac{1}{2}|\mathbf{b} \times \mathbf{n}|$ is the area of ΔOBN and the line segment \overline{ON} is unit length, $C(\rho_B, \mathbf{n})$ is simply the perpendicular distance between the point B and the line \overline{ON} .

Similarly, we can write a two-qubit state in the Pauli basis as $\rho = \frac{1}{4}\sum_{i,j=0}^3 \Theta_{ij} \sigma_i^A \otimes \sigma_j^B$, where the coefficients $\Theta_{ij} = \text{tr}(\rho \sigma_i^A \otimes \sigma_j^B)$ form a block matrix $\Theta = \begin{pmatrix} \mathbf{1} & \mathbf{b}^T \\ \mathbf{a} & T \end{pmatrix}$. Here \mathbf{a} and \mathbf{b} are Alice and Bob's Bloch vectors respectively, and T is a 3×3 matrix. Note that when ρ_B is non-degenerate we have $\mathbf{b} \neq \mathbf{0}$. We ignore the trivial case that $\mathbf{a} = \mathbf{1}$, when ρ_A is pure and hence ρ is a product state.

When the POVM operator $M = \frac{1}{2}(\mathbf{1} + \mathbf{m} \cdot \boldsymbol{\sigma})$ is obtained on Alice's qubit, Bob's state becomes

$$\mathbf{b}_M = \frac{\mathbf{b} + T^T \mathbf{m}}{1 + \mathbf{a} \cdot \mathbf{m}}. \quad (7)$$

Here $\boldsymbol{\sigma} = (\sigma_1, \sigma_2, \sigma_3)^T$, and $\mathbf{m} = (m_1, m_2, m_3)^T$ can be any point on or inside the Bloch sphere. The set of \mathbf{b}_M forms the QSE \mathcal{E}_B . When ρ_B is non-degenerate, we have $\mathbf{b} \neq \mathbf{0}$. According to Eq. (6), the coherence of \mathbf{b}_M in the basis $\{|\xi_i\rangle\}$ is $C(\rho_B^M, \mathbf{n}_B) = |\mathbf{b}_M \times \mathbf{n}_B|$, with $\mathbf{n}_B = \mathbf{b}/b$; this represents the perpendicular distance from the point B_M to the line \overline{OB} (Fig. 1a). Hence the maximal steered coherence $\mathcal{C}(\rho)$, as defined in Eq. (2), is the maximal perpendicular distance between a point on the surface of \mathcal{E}_B and \overline{OB} . Explicitly, we have $C(\rho_B^M, \mathbf{n}_B) = |(T^T \mathbf{m}) \times \mathbf{n}_B|/|1 + \mathbf{a} \cdot \mathbf{m}|$ and

$$\mathcal{C}(\rho) = \max_{\mathbf{m} \in \mathbb{R}^3, m=1} \left| \frac{T^T \mathbf{m} \times \mathbf{n}_B}{1 + \mathbf{a} \cdot \mathbf{m}} \right|. \quad (8)$$

The maximization needs to be performed only over all projective measurements with $m = 1$ because steered states on the surface of \mathcal{E}_B correspond to measurements \mathbf{m} on the surface of the Bloch sphere.

When ρ_B is degenerate, $\mathbf{b} = \mathbf{0}$ and \mathbf{n}_B is arbitrary; the infimum can then be taken over all \mathbf{n}_B to give the maximal steered coherence of a two-qubit state as

$$\mathcal{C}(\rho) = \inf_{\mathbf{n}_B \in \mathbb{R}^3, n_B=1} \left(\max_{\mathbf{m} \in \mathbb{R}^3, m=1} \left| \frac{T^T \mathbf{m} \times \mathbf{n}_B}{1 + \mathbf{a} \cdot \mathbf{m}} \right| \right). \quad (9)$$

Properties for two-qubit states. We now study two-qubit states in more detail; this allows us to identify some important features of the maximal steered coherence, as well as giving a clear geometric interpretation of \mathcal{C} using the steering ellipsoid formalism.

As demonstrated by property (E4), a trace-preserving channel Λ_B performed by Bob may increase \mathcal{C} ; we now study an explicit example. Say that Alice and Bob share the classical two-qubit state

$$\rho_c = t|++\rangle\langle++| + (1-t)|--\rangle\langle--|, \quad (10)$$

with $\frac{1}{2} < t < 1$ and $|\pm\rangle = \frac{1}{\sqrt{2}}(|0\rangle \pm |1\rangle)$. When Bob applies the single-qubit amplitude damping channel, the state transforms as $\rho'_c = \mathbf{1}_A \otimes \Lambda_B^{\text{AD}}(\rho_c)$, where $\Lambda^{\text{AD}}(\cdot) = \sum_i E_i(\cdot) E_i^\dagger$ with $E_0 = |0\rangle\langle 0| + \sqrt{1-\gamma}|1\rangle\langle 1|$ and $E_1 = \sqrt{\gamma}|0\rangle\langle 1|$. We then find that the maximal steered coherence of the transformed state is

$$\mathcal{C}(\rho'_c) = \frac{2t\gamma\sqrt{1-\gamma}}{\sqrt{(1-2t)^2(1-\gamma) + \gamma^2}}. \quad (11)$$

$\mathcal{C}(\rho'_c)$ vanishes when $\gamma = 0$, becomes positive for $0 < \gamma < 1$, and then vanishes again at $\gamma = 1$.

Maximal steered coherence can be increased by Bob's local amplitude damping channel even when Alice and Bob share a non-classical state. Consider the two-qubit state

$$\rho_p = p|\Psi\rangle\langle\Psi| + \frac{1-p}{4}\sigma_0 \otimes \sigma_0, \quad (12)$$




Figure 1. An illustration of the geometric interpretation of maximal steered coherence $\mathcal{C}(\rho)$ for two-qubit states ρ using the QSE \mathcal{E}_B . For simplicity we take $a = 0$. The point B representing Bob's Bloch vector is indicated by a green blob, and the line \overline{OB} is also shown in green; states lying along this line are incoherent in the basis ρ_B . $\mathcal{C}(\rho)$ is given by the maximal perpendicular distance between a point on \mathcal{E}_B and \overline{OB} ; this is shown by the red arrow. (a) Theorem 2 shows that for any canonical state, $\mathcal{C}(\rho)$ is bounded by the longest semiaxis of the QSE. (b) A state of the form (15), which achieves maximal $\mathcal{C}(\rho)$ for a given b . The QSE is a chord perpendicular to \overline{OB} . (c) When ρ is an X state, \overline{OB} lies along an axis of the QSE, and $\mathcal{C}(\rho)$ is the length of the longest of the other two semiaxes. (d) When ρ is a Werner state, \mathcal{E}_B is a ball centred on the origin. In this case, even though ρ_B is degenerate, $\mathcal{C}(\rho)$ is well-defined as the radius of the ball.

where $0 < p < 1$ and $|\Psi\rangle = \cos \frac{\theta}{2} |++\rangle + \sin \frac{\theta}{2} |--\rangle$. The QSE for such a state is an ellipsoid centered at $\left(\frac{p(1-p)\cos\theta}{1-(p\cos\theta)^2}, 0, 0 \right)$ with semiaxes of length $c_1 = \frac{p(1-p\cos^2\theta)}{1-(p\cos\theta)^2}$, $c_2 = c_3 = \frac{p\sin\theta}{\sqrt{1-(p\cos\theta)^2}}$ aligned with the coordinate axes (\mathcal{E}_B is in fact a prolate spheroid as $c_1 > c_2 = c_3$). Bob's Bloch vector is $\mathbf{b} = (p\cos\theta, 0, 0)$, which lies on the x axis. The maximal steered coherence is therefore

$$\mathcal{C}(\rho_p) = \frac{p\sin\theta}{\sqrt{1-(p\cos\theta)^2}}. \quad (13)$$

For $p < \frac{1}{2\sin\theta+1}$, the state ρ_p has zero entanglement but nonzero \mathcal{C} . Note also that $\mathcal{C}(\rho'_p)$ is related to both the fraction of $|\Psi\rangle$ and the entanglement associated with $|\Psi\rangle$.

Figure 2 shows the evolution of \mathcal{C} under the channel Λ_B^{AB} , i.e. $\mathcal{C}(\rho'_p)$, where $\rho'_p = \mathbb{1}_A \otimes \Lambda_B^{AD}(\rho_p)$. By altering p and θ we alter the ratio of the axes c_3/c_1 . The results indicate that the potential for increasing \mathcal{C} under Bob's local amplitude damping is related to the ratio c_3/c_1 : the smaller the ratio, the stronger the local increase of \mathcal{C} . In other words the effect is strongest when the QSE \mathcal{E}_B is highly prolate ('baguette-shaped').

In fact, it is possible to formulate a necessary and sufficient condition for the increase of maximal steered coherence for two-qubit states.

Theorem 1. Bob's local qubit channel Λ_B can increase maximal steered coherence for some input two-qubit state if and only if Λ_B is neither unital nor semi-classical.

The proof is given in Methods. We therefore see that the behavior of maximal steered coherence \mathcal{C} under local operations is similar to that of quantum discord \mathcal{D} . The set of local channels that can increase \mathcal{C} for some two-qubit state is the same as the set of local channels that can increase \mathcal{D} . Moreover, \mathcal{C} can be increased when the QSE \mathcal{E}_B is very prolate; we showed in Reference²⁸ that the quantum discord of Bell-diagonal states with such




Figure 2. The evolution of maximal steered coherence under Bob's local amplitude damping channel: $\mathcal{C}(\mathbf{1}_A \otimes \Lambda_B^{\text{AD}}(\rho_p))$, with ρ_p given by Eq. (12). The parameters for the four curves are $p = 0.9, \theta = 0.2\pi$ for the cyan dashed line; $p = 0.9, \theta = 0.1\pi$ for the red dotted line; $p = 0.7, \theta = 0.1\pi$ for the green solid line; and $p = 0.5, \theta = 0.1\pi$ for the blue dash-dotted line. The corresponding semiaxes ratios, which give a measure of the prolateness of \mathcal{E}_B , are $c_3/c_1 = 0.980, 0.859, 0.629$ and 0.496 respectively. The effect of locally increasing \mathcal{C} is stronger for more prolate \mathcal{E}_B .

baguette-shaped \mathcal{E}_B can be increased by the local amplitude damping channel. We therefore conjecture the local increase in quantum correlations originates from the increase in steered coherence.

We now investigate the set of so-called canonical states, which have particular significance in the steering ellipsoid formalism^{24,25,33}. Here, a canonical state ρ_{can} corresponds to one for which Alice's marginal is maximally mixed ($\mathbf{a} = \mathbf{0}$). This implies that the QSE \mathcal{E}_B is centered at B (Fig. 1a). Let c_1, c_2 and c_3 be the lengths of the semiaxes of \mathcal{E}_B ordered such that $c_1 \geq c_2 \geq c_3$.

Theorem 2. *For any canonical state ρ_{can} the maximal steered coherence is bounded by the longest semiaxis. This in turn is bounded by the length of Bob's Bloch vector as*

$$\mathcal{C}(\rho_{\text{can}}) \leq c_1 \leq \sqrt{1 - b^2}. \quad (14)$$

The bound is saturated if and only if \mathcal{E}_B is a chord perpendicular to \mathbf{b} meeting the surface of the Bloch sphere at $|\chi\rangle$ and $|\chi'\rangle$. This represents a canonical state of the form

$$\rho_{\text{can}} = \frac{1}{2}|\psi\rangle\langle\psi| \otimes |\chi\rangle\langle\chi| + \frac{1}{2}|\bar{\psi}\rangle\langle\bar{\psi}| \otimes |\chi'\rangle\langle\chi'|, \quad (15)$$

where $\langle\psi|\bar{\psi}\rangle = 0$.

The proof is given in Methods, and an example QSE for an optimal state of the form (15) is shown in Fig. 1b). Note that this bound is remarkably simple and geometrically intuitive: it depends only on the longest semiaxis of \mathcal{E}_B and not on the orientation or position of the QSE. Theorem 2 is in the same vein as bounds presented in Reference³³ that relate several other measures of quantum correlation to geometric features of QSEs.

We also note that optimal states of the form (15) have the highest quantum discord among discordant states with a given \mathbf{b} that are obtained from classical states by a local trace-preserving channel. As shown in Reference³⁴, when we take a two-qubit B -side classical (zero discord) state and apply a channel Λ_B to Bob's qubit, in order to create maximal B -side quantum discord in the output state, the optimal input state is of the form $\frac{1}{2}|\psi\rangle\langle\psi| \otimes |\phi\rangle\langle\phi| + \frac{1}{2}|\bar{\psi}\rangle\langle\bar{\psi}| \otimes |\bar{\phi}\rangle\langle\bar{\phi}|$, and the channel Λ_B should have Kraus operators $E_0 = |\chi\rangle\langle\phi|$, $E_1 = |\chi'\rangle\langle\bar{\phi}|$, where $|\chi\rangle$ and $|\chi'\rangle$ are determined by \mathbf{b} .

Examples. Let us now examine some interesting classes of two-qubit states for which maximal steered coherence is easy both to find analytically and to interpret geometrically using QSEs.

X states. When \mathbf{n}_B lies along an axis of the QSE \mathcal{E}_B it is straightforward to see that $\mathcal{C}(\rho)$ is simply the length of the longest of the other two semiaxes (Fig. 1c). All ρ which are X states, i.e. have non-zero entries only in the characteristic X shape in the computational basis³⁵, will have such QSEs²⁶.

Werner states. As a special case of the above, when \mathcal{E}_B is a ball of radius r centered on O' and \mathbf{n}_B is collinear with OO' , we have $\mathcal{C}(\rho) = r$. Furthermore, when \mathcal{E}_B is an origin-centered ball, we have $\mathcal{C}(\rho) = r$ regardless of the value of \mathbf{b} . This allows us to evaluate \mathcal{C} for Werner states³⁶, which do not in fact satisfy the non-degenerate

condition $\mathbf{b} \neq \mathbf{0}$. For a Werner state $\rho_W = p|\Psi^-\rangle\langle\Psi^-| + \frac{1-p}{4}\sigma_0 \otimes \sigma_0$ with $|\Psi^-\rangle = \frac{1}{\sqrt{2}}(|01\rangle - |10\rangle)$, \mathcal{C}_B is an origin-centered ball of radius p and hence $\mathcal{C}(\rho_W) = p$ (Fig. 1d).

Discordant states locally created from a classical state. We know from property (E1) that \mathcal{C} vanishes for classical states; for a classical two-qubit state, all steered states must have the same orientation, and the QSE \mathcal{C}_B is therefore a radial line segment. For a state obtained locally from a classical state, ρ_{dlc} , the QSE is a nonradial line segment²⁵. \mathbf{b} can be any point on this segment except for the two ends of \mathcal{C}_B , which we call \mathbf{b}_1 and \mathbf{b}_2 , where $b_1 \geq b_2$. By definition $\mathcal{C}(\rho_{\text{dlc}})$ varies for different \mathbf{b} ; in general, we find that

$$b_1 \sin \theta_1 \leq \mathcal{C}(\rho_{\text{dlc}}) \begin{cases} < b_1 \sin \theta, & \text{for } \theta \leq \frac{\pi}{2}, \\ \leq b_1, & \text{for } \theta > \frac{\pi}{2}, \end{cases} \quad (16)$$

where θ is the angle between \mathbf{b}_1 and \mathbf{b}_2 and θ_1 is determined by $b_1 \sin \theta_1 = b_2 \sin(\theta - \theta_1)$. From Eq. (16), we see that $\mathcal{C}(\rho_{\text{dlc}})$ is strictly larger than zero. In fact, $\mathcal{C}(\rho_{\text{dlc}})$ can reach unity when $b_1 = 1$ and $\theta > \frac{\pi}{2}$.

Maximally obese states. The general form of a maximally obese state is given by³³

$$\rho_{\text{mo}} = \left(1 - \frac{b}{2}\right)|\Psi_b\rangle\langle\Psi_b| + \frac{b}{2}|00\rangle\langle00|, \quad (17)$$

where $|\Psi_b\rangle = \frac{1}{\sqrt{2-b}}(\sqrt{1-b}|01\rangle + |10\rangle)$. This is a canonical state ($\mathbf{a} = \mathbf{0}$) with \mathcal{C}_B centered at $(0, 0, b)$ and semiaxes of length $c_1 = c_2 = \sqrt{1-b}$, $c_3 = 1 - b$ aligned with the coordinate axes. We therefore have $\mathcal{C}(\rho_{\text{mo}}) = \sqrt{1-b}$. It should be noted that maximally obese states maximize several measures of quantum correlation (CHSH nonlocality, singlet fraction, concurrence and negativity) over the set of all canonical states with a given marginal for Bob³³. Interestingly, however, they do not achieve the maximum possible $\mathcal{C} = \sqrt{1-b^2}$.

Discussion

We have studied the maximal steered coherence $\mathcal{C}(\rho)$ for a bipartite state ρ . When Alice obtains a POVM outcome M , Bob's state is steered to ρ_B^M ; $\mathcal{C}(\rho)$ is defined as the coherence of the steered state in Bob's original basis, maximized over all possible M . The general form of \mathcal{C} is derived for two-qubit states. By calculating the maximal steered coherence for some important classes of two-qubit state, we find that \mathcal{C} gives a different ordering of states from quantum entanglement or discord-like correlations. This means that \mathcal{C} is a distinct and new measure for characterizing the remote quantum properties of bipartite states.

The maximal steered coherence vanishes only when ρ is a classical state, and $\mathcal{C}(\rho)$ can be increased by local trace-preserving channels. For a two-qubit state ρ we derive a necessary and sufficient condition for a local qubit channel to be capable of increasing $\mathcal{C}(\rho)$. This is in fact identical to the condition for increasing quantum discord, suggesting that local increase of quantum discord might be used in a protocol for increasing steered coherence.

Finally, we consider the relevance of \mathcal{C} from a more physical perspective by presenting a concrete example in which steered coherence can be exploited. Say that Alice and Bob share a two-qubit state of the form (15) with $|\psi\rangle = |0\rangle, |\bar{\psi}\rangle = |1\rangle$ and $\mathbf{b} = (0, 0, b)$ (which, as illustrated in Fig. 1b, will lie at the midpoint of the chord \mathcal{C}_B joining $|\chi\rangle$ and $|\chi'\rangle$). Suppose also that Alice's and Bob's systems are described by the local Hamiltonian $H = -\hbar\omega\sigma_3$. Let us restrict Alice's and Bob's local operations to those which are covariant with respect to time-translation symmetry³⁷. For Alice, these operations are the ones for which $\Phi(e^{-iHt}\rho_A^M e^{iHt}) = e^{-iHt}\Phi(\rho_A) e^{iHt}$; and similarly for Bob. Physically, this restriction corresponds to local energy-conserving unitaries with the assistance of incoherent environmental ancillas³⁸: Alice's operation is covariant if and only if it can be written as $\Phi(\rho_A) = \text{tr}_E[U(\rho_A \otimes \xi_E)U^\dagger]$, where U is a unitary, H_E is the Hamiltonian of the ancilla, $[U, H + H_E] = 0$ and $[\xi_E, H_E] = 0$; and similarly for Bob. The set of covariant operations is a strict subset of incoherent operations¹⁰ and a strict superset of thermal operations³⁹.

Bob's reduced state $\rho_B = \frac{1}{2}(1 + b\sigma_3)$ is incoherent in his energy eigenbasis $\{|0\rangle, |1\rangle\}$, and his local covariant operations alone cannot generate any coherence. However, by performing a σ_3 measurement, which is a covariant operation, and classically communicating the result to Bob, Alice steers him to either $|\chi\rangle$ or $|\chi'\rangle$, states that are manifestly coherent in the energy eigenbasis. $\mathcal{C}(\rho)$ gives a measure of the maximal coherence that Alice can induce on Bob's system by steering. In this way, Alice remotely 'activates' a coherent state for Bob that he was unable to produce himself. Bob may now use this coherence as a resource for quantum information processing tasks, e.g. work extraction by a thermal machine, which is known to be enhanced in the presence of a coherent reference system⁴⁰. Given the ever-increasing number of applications for coherence found throughout quantum information science, one can envisage a range of such scenarios in which steered coherence could be used as a resource.

Methods

Proof of property (E1). The 'if' part is obvious: Bob's reduced state is $\rho_B = \sum_i p_i |\xi_i\rangle\langle\xi_i|$, and the steered state is $\rho_B^M = \sum_i p_i \text{tr}(\rho_i^A M) |\xi_i\rangle\langle\xi_i|$. These are both diagonal in the basis $\{|\xi_i\rangle\}$, and hence $\mathcal{C}(\rho) = 0$.

For the 'only if' part, first consider a separable state $\rho_s = \sum_i p_i \rho_i^A \otimes \rho_i^B$. When $\mathcal{C}(\rho_s) = 0$, the steered states $\rho_B^M = \sum_i p_i \text{tr}(\rho_i^A M) \rho_i^B$ for different POVM operators M should commute with each other, which is equivalent

to all ρ_i^B commuting with each other. So $\mathcal{C}(\rho_s)$ vanishes only if it is in the form (4). For an entangled state ρ_e , we express ρ_e in the optimal pure state decomposition form as $\rho_e = \sum_i p_i |\Phi_i\rangle\langle\Phi_i|$, so that the entanglement of formation is $E_F(\rho_e) = \sum_i p_i E_F(|\Phi_i\rangle\langle\Phi_i|)$. Since ρ_e is entangled, at least one of the $|\Phi_i\rangle$ is entangled. Hence, for ρ_e , it is not possible for all of Bob's steered states to share the same eigenbasis; this means that $\mathcal{C}(\rho_e) \neq 0$ for any entangled ρ_e .

Proof of Theorem 1. A channel Λ_B that is neither unital nor semi-classical can increase \mathcal{C} , because such channels can transform a classical state with vanishing \mathcal{C} into a discordant state with nonzero \mathcal{C} ^{30,41}. We now focus on the 'only if' part, and prove that a local unital channel or a local semi-classical channel cannot increase \mathcal{C} for any two-qubit input state.

A semi-classical channel Λ^{sc41} , which maps any input state ρ to a state with zero coherence in a given basis $\Lambda^c(\rho) = \sum_k p_k(\rho) |k\rangle\langle k|$, yields $\mathcal{C} = 0$ for any input state. As proved by King and Ruskai⁴², any unital channel is equivalent to $\Lambda^u(\cdot) = \sum_{i=0}^3 e_i \sigma_i(\cdot) \sigma_i$, where $0 \leq e_i \leq 1$ and $\sum_i e_i = 1$. The effect of this channel on a qubit state is to shrink the Bloch vector as $\Lambda^u: \mathbf{b} \rightarrow \mathbf{b}' = (p_1 b_1, p_2 b_2, p_3 b_3)^T$, where $p_1 = e_0 + e_1 - e_2 - e_3$, and $p_{2,3}$ are related to e_i in a similar way. Let \mathbf{b}_M be a steered state for the input state ρ . Then the coherence of \mathbf{b}_M is $C(\mathbf{b}_M, \mathbf{b}) = |\mathbf{b}_M \times \mathbf{b}|/b$. Under the action of Λ^u , the steered state and Bob's reduced state become \mathbf{b}'_M and \mathbf{b}' respectively, and the coherence of \mathbf{b}'_M in the eigenbasis of \mathbf{b}' is

$$C^2(\mathbf{b}'_M, \mathbf{b}') = \frac{(b_{M1}b_3 - b_{M3}b_1)^2 p_1^2 p_3^2}{(b_1 p_1)^2 + (b_2 p_2)^2 + (b_3 p_3)^2} + (1 \rightarrow 2 \rightarrow 3) + (1 \rightarrow 3 \rightarrow 2). \quad (18)$$

If the inequality

$$C(\mathbf{b}'_M, \mathbf{b}') \leq C(\mathbf{b}_M, \mathbf{b}) \quad (19)$$

holds then the maximal steered coherence for the output state $\mathcal{C}(1_A \otimes \Lambda_B^u(\rho)) = \mathcal{C}(\mathbf{b}'_{M_{opt}}, \mathbf{b}') \leq \mathcal{C}(\mathbf{b}_{M_{opt}}, \mathbf{b}) \leq \mathcal{C}(\rho)$, where M_{opt} is the optimal POVM operator to maximize (2) for the output state and $\mathbf{b}_{M_{opt}}$ is the corresponding input state for $\mathbf{b}'_{M_{opt}}$. Hence it is sufficient to prove that (19) holds for some \mathbf{b}_M and \mathbf{b} . Note that $\frac{(b_{M1}b_3 - b_{M3}b_1)^2 p_1^2 p_3^2}{(b_1 p_1)^2 + (b_3 p_3)^2} \leq \frac{(b_{M1}b_3 - b_{M3}b_1)^2}{b_1^2 + b_3^2}$. By using the fact that $\frac{A+B}{C+D} < \frac{A'+B}{C'+D}$ for $0 < B < D$, $0 < A < C$, $0 < A' < C'$ and $\frac{A}{C} > \frac{A'}{C'}$, we arrive at $\mathcal{C}^2(\mathbf{b}'_M, \mathbf{b}') \leq C^2(\mathbf{b}_M, \mathbf{b})$, which is equivalent to (19).

Proof of Theorem 2. The steered state \mathbf{b}_M which achieves the maximum in Eq. (8) corresponds to a point B_M on the surface of \mathcal{E}_B . We have $\mathcal{C}(\rho_{can}) = C(\mathbf{b}_M, \mathbf{n}_B) = D(B_M, \overline{OB}) \leq \overline{BB_M} \leq c_1$, where $D(B_M, \overline{OB})$ is the perpendicular distance between B_M and \overline{OB} . To ensure that \mathcal{E}_B lies inside the Bloch sphere we require that $c_1 \leq \sqrt{1 - b^2}$.

To saturate the bound we take $c_1 = \sqrt{1 - b^2}$, but we must also demonstrate that $c_2 = c_3 = 0$, i.e. that \mathcal{E}_B cannot be an ellipsoid or an ellipse. We know that \mathcal{E}_B must meet the Bloch sphere at two points, corresponding to the pure states $|\chi\rangle$ and $|\chi'\rangle$. Firstly suppose that \mathcal{E}_B is a three-dimensional ellipsoid. Elementary geometry tells us that the surface of an ellipsoid at the end of any axis must be perpendicular to that axis. The points at the ends of the c_1 axis on \mathcal{E}_B lie on the surface of the Bloch sphere. Since the surface of \mathcal{E}_B must lie perpendicular to the c_1 axis at these points, \mathcal{E}_B must puncture the surface of the Bloch sphere. Such \mathcal{E}_B cannot represent a physical two-qubit state and so \mathcal{E}_B cannot be an ellipsoid. Now consider the case that \mathcal{E}_B is an ellipse. The nested tetrahedron condition tells us that any degenerate \mathcal{E}_B describing a physical state must fit inside a triangle inside the Bloch sphere^{25,26}. Geometrically, no ellipse that touches the Bloch sphere at two points can satisfy this, and so \mathcal{E}_B cannot be an ellipse. \mathcal{E}_B must therefore be a line, i.e. the chord going between $|\chi\rangle$ and $|\chi'\rangle$; this corresponds to the state (15).

References

1. Bartlett, S. D., Rudolph, T. & Spekkens, R. W. Reference frames, superselection rules, and quantum information. *Rev. Mod. Phys.* **79**, 555–609 (2007).
2. Marvian, I. & Spekkens, R. W. The theory of manipulations of pure state asymmetry: basic tools and equivalence classes of states under symmetric operations. *New J. Phys.* **15**, 033001 (2013).
3. Marvian, I. & Spekkens, R. W. Modes of asymmetry: the application of harmonic analysis to symmetric quantum dynamics and quantum reference frames. *Phys. Rev. A* **90**, 062110 (2014).
4. Lloyd, S. Quantum coherence in biological systems. *J. Phys. Conf. Ser.* **302**, 012037 (2011).
5. Li, C.-M., Lambert, N., Chen, Y.-N., Chen, G.-Y. & Nori, F. Examining non-locality and quantum coherent dynamics induced by a common reservoir. *Sci. Rep.* **2**, 885 (2012).
6. Lambert, N. *et al.* Quantum biology. *Nat. Phys.* **9**, 10–18 (2013).
7. Narasimhachar, V. & Gour, G. Low-temperature thermodynamics with quantum coherence. *arXiv:1409.7740* (2014).
8. Åberg, J. Catalytic coherence. *Phys. Rev. Lett.* **113**, 150402 (2014).
9. Ćwikliński, P., Studziński, M., Horodecki, M. & Oppenheim, J. Towards fully quantum second laws of thermodynamics: limitations on the evolution of quantum coherences. *arXiv:1405.5029* (2014).
10. Baumgratz, T., Cramer, M. & Plenio, M. B. Quantifying coherence. *Phys. Rev. Lett.* **113**, 140401 (2014).
11. Girolami, D. Observable measure of quantum coherence in finite dimensional systems. *Phys. Rev. Lett.* **113**, 170401 (2014).
12. Shao, L.-H., Xi, Z., Fan, H. & Li, Y. Fidelity and trace-norm distances for quantifying coherence. *Phys. Rev. A* **91**, 042120 (2015).
13. Pires, D. P., Céleri, L. C. & Soares-Pinto, D. O. Geometric lower bound for a quantum coherence measure. *Phys. Rev. A* **91**, 042330 (2015).

14. Singh, U., Bera, M. N., Dhar, H. S. & Pati, A. K. Maximally coherent mixed states: Complementarity between maximal coherence and mixedness. *Phys. Rev. A* **91**, 052115 (2015).
15. Zhang, Y.-J., Han, W., Xia, Y.-J., Yu, Y.-M. & Fan, H. Role of initial system-bath correlation on coherence trapping. *Sci. Rep.* **5**, 13359 (2015).
16. Chin, A. W. *et al.* The role of non-equilibrium vibrational structures in electronic coherence and recoherence in pigmentprotein complexes. *Nat. Phys.* **9**, 113 (2013).
17. Bromley, T. R., Cianciaruso, M. & Adesso, G. Frozen quantum coherence. *Phys. Rev. Lett.* **114**, 210401 (2015).
18. Schrödinger, E. Discussion of probability relations between separated systems. *Proc. Cambridge Philos. Soc.* **31**, 555 (1935).
19. Wiseman, H. M., Jones, S. J. & Doherty, A. C. Steering, entanglement, nonlocality, and the einstein-podolsky-rosen paradox. *Phys. Rev. Lett.* **98**, 140402 (2007).
20. Saunders, D. J., Jones, S. J., Wiseman, H. M. & Pryde, G. J. Experimental epr-steering using bell-local states. *Nat. Phys.* **6**, 845 (2010).
21. Händchen, V. *et al.* Observation of one-way einstein-podolsky-rosen steering. *Nat. Photon.* **6**, 596 (2012).
22. Skrzypczyk, P., Navascués, M. & Cavalcanti, D. Quantifying einstein-podolsky-rosen steering. *Phys. Rev. Lett.* **112**, 180404 (2014).
23. Verstraete, F. *A study of entanglement in quantum information theory*. Ph.D. thesis, Katholieke Universiteit Leuven (2002).
24. Shi, M., Yang, W., Jiang, F. & Du, J. Geometric picture of quantum discord for two-qubit quantum states. *J. Phys. A: Math. Theor.* **44**, 415304 (2011).
25. Jevtic, S., Pusey, M., Jennings, D. & Rudolph, T. Quantum steering ellipsoids. *Phys. Rev. Lett.* **113**, 020402 (2014).
26. Milne, A., Jevtic, S., Jennings, D., Wiseman, H. & Rudolph, T. Quantum steering ellipsoids, extremal physical states and monogamy. *New J. Phys.* **16**, 083017 (2014).
27. Milne, A., Jennings, D. & Rudolph, T. Geometric representation of two-qubit entanglement witnesses. *Phys. Rev. A* **92**, 012311 (2015).
28. Hu, X. & Fan, H. Effect of local channels on quantum steering ellipsoids. *Phys. Rev. A* **91**, 022301 (2015).
29. Debasis Mondal, A. K. P. Tanumoy Pramanik. Steerability of local quantum coherence. *arXiv:1508.03770* (2015).
30. Hu, X., Fan, H., Zhou, D. L. & Liu, W.-M. Necessary and sufficient conditions for local creation of quantum correlation. *Phys. Rev. A* **85**, 032102 (2012).
31. Wootters, W. K. Entanglement of formation of an arbitrary state of two qubits. *Phys. Rev. Lett.* **80**, 2245–2248 (1998).
32. Zurek, W. H. Quantum discord and maxwell's demons. *Phys. Rev. A* **67**, 012320 (2003).
33. Milne, A., Jennings, D., Jevtic, S. & Rudolph, T. Quantum correlations of two-qubit states with one maximally mixed marginal. *Phys. Rev. A* **90**, 024302 (2014).
34. Hu, X., Fan, H., Zhou, D. L. & Liu, W.-M. Quantum correlating power of local quantum channels. *Phys. Rev. A* **87**, 032340 (2013).
35. Yu, T. & Eberly, J. Evolution from entanglement to decoherence of bipartite mixed “x” states. *Quantum Inf. Comput.* **7**, 459 (2007).
36. Werner, R. F. Quantum states with einstein-podolsky-rosen correlations admitting a hidden-variable model. *Phys. Rev. A* **40**, 4277–4281 (1989).
37. Marvian, I. & Spekkens, R. W. Extending Noether's theorem by quantifying the asymmetry of quantum states. *Nat. Commun.* **5**, 3821 (2014).
38. Keyl, M. & Werner, R. F. Optimal cloning of pure states, testing single clones. *J. Math. Phys.* **40**, 3283–3299 (1999).
39. Brandão, F. G. S. L., Horodecki, M., Oppenheim, J., Renes, J. M. & Spekkens, R. W. Resource theory of quantum states out of thermal equilibrium. *Phys. Rev. Lett.* **111**, 250404 (2013).
40. Korzekwa, K., Lostaglio, M., Oppenheim, J. & Jennings, D. The extraction of work from quantum coherence. *arXiv:1506.07875* (2015).
41. Streltsov, A., Kampermann, H. & Bruß, D. Behavior of quantum correlations under local noise. *Phys. Rev. Lett.* **107**, 170502 (2011).
42. King, C. & Ruskai, M. B. Minimal entropy of states emerging from noisy quantum channels. *IEEE Trans. Info. Theor.* **47**, 192 (2001).

Acknowledgements

This work was supported by NSFC under Grant No. 11447161,11504205, the Fundamental Research Funds of Shandong University under Grant No. 2014TB018, and the National Key Basic Research Program of China under Grant No. 2015CB921003. A.M. is funded by EPSRC. We thank Kamil Korzekwa, Matteo Lostaglio and Terry Rudolph for useful discussions.

Author Contributions

X.H. contributed the idea. X.H., A.M. and B.Z. performed the calculations. X.H., A.M. and H.F. wrote the paper. All authors reviewed the manuscript and agreed with the submission.

Additional Information

Competing financial interests: The authors declare no competing financial interests.

How to cite this article: Hu, X. *et al.* Quantum coherence of steered states. *Sci. Rep.* **6**, 19365; doi: 10.1038/srep19365 (2016).



This work is licensed under a Creative Commons Attribution 4.0 International License. The images or other third party material in this article are included in the article's Creative Commons license, unless indicated otherwise in the credit line; if the material is not included under the Creative Commons license, users will need to obtain permission from the license holder to reproduce the material. To view a copy of this license, visit <http://creativecommons.org/licenses/by/4.0/>